

Copyright for test papers and marking guides remains with West Australian Test Papers. Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale. Test papers should be withdrawn after use and stored securely in the school until Friday 12th June 2020.



**SEMESTER ONE**

**MATHEMATICS**

**METHODS**

**UNIT 3**

2020

## SOLUTIONS

$$\begin{aligned} \text{When } \theta &= 1.134^\circ, \quad dx = 0.559 \text{ m} \\ \Delta x &= -\frac{\cos \theta}{10} \times (-0.01) \\ x &= 30 - 10 \tan \theta \quad \leftarrow \frac{dx}{d\theta} = -\frac{1}{10} \cos^2 \theta \end{aligned}$$

$$\begin{aligned}
 & L(x) = \frac{1}{2} \left[ 16x + x^2 \right] - \frac{1}{2} \left[ (2x - 60) + (x^2 - 60x + 100) \right] \\
 & L(x) = \frac{1}{2} \left[ 16x + x^2 \right] - \frac{1}{2} \left[ x^2 - 60x + 100 \right] \\
 & L(x) = \frac{1}{2} \left[ 16x + 60x \right] - \frac{1}{2} \cdot 100 \\
 & L(x) = \frac{1}{2} \left[ 76x \right] - \frac{1}{2} \cdot 100 \\
 & L(x) = 38x - 50 \\
 & \text{min occurs when } L(x) = 0 \\
 & 38x - 50 = 0 \\
 & 38x = 50 \\
 & x = \frac{50}{38} \\
 & x = 1.34
 \end{aligned}$$

$$M(t) = 2 \cdot 10^6 \cdot 0.04463$$

2020

©WATP

$$\begin{aligned}
 & \text{Arba} = \int_{-2}^2 f(x) dx = \int_{-2}^2 f(x) dx \quad \text{since positive} \\
 & \text{Arba} = \int_{-2}^2 f(x) dx = \int_{-2}^2 f(x) dx = 0 \\
 & f'(x) = f(-2) - f(2) = 0 - 0 = 0 \\
 & \int_{-2}^2 f(x) dx = f(-2) - f(-2) = 2 - (-2) = 4
 \end{aligned}
 \tag{c)
 }$$

(e) 6

<p>8. (a) <math>\frac{30 + 30}{30} = \frac{150}{30} + \frac{240}{30} = \frac{50}{16.6}</math></p> <p><b>Calculus-Asymptote Solutions</b></p>	<p><math>a = 16</math> and <math>b = 10</math></p> <p><b>Calculus-Asymptote Solutions</b></p>
<p><b>Calculus-Asymptote Solutions</b></p>	<p><b>Calculus-Asymptote Solutions</b></p>

## Calculator-free Solutions

1. (a)  $\frac{d}{dx_L} \left[ \left( \sin\left(\frac{x}{2}\right) \right)^3 \right] = 3\sin^2\left(\frac{x}{2}\right) \times \cos\left(\frac{x}{2}\right) \times \frac{1}{2}$  ✓✓

(b)  $2t(\tan t) + \frac{t^2}{\cos^2 t}$  ✓✓

(c)  $f(y) = \cos\left(\frac{1}{(\sin y)^2}\right)$

$f'(y) = -\sin\sqrt{\sin y} \times \frac{1}{2} \left( \sin^{-\frac{1}{2}} y \right) \times \cos y$  ✓✓✓ [7]

2. (a)  $v(t) = 2e^{2t} - 2e^2 t + c$  ✓

$x(t) = e^{2t} - e^2 t^2 + ct + k$  ✓

When  $t = 0$ ,  $x = 0 \rightarrow k = -1$  ✓

When  $t = 1$ ,  $x = 0 \rightarrow c = 1$  ✓

$x(t) = e^{2t} - e^2 t^2 + t - 1$  ✓

(b) 0 ✓

(c)  $v(0) = 3$  ✓ [6]

3. (a)  $2\sin\frac{x}{2} - e^{2\cos x} + c$  ✓✓

(b)  $f'(y) = (1-2y)^{-\frac{1}{2}}$  ✓✓

$f(y) = \frac{(1-2y)^{\frac{1}{2}}}{\left(\frac{1}{2}\right) \times (-2)}$  ✓✓

$(-4, 3) \rightarrow 3 = -\sqrt{9} + c \rightarrow c = 6$  ✓

$f(y) = -\sqrt{1-2y} + 6$  ✓

$g(x) = -\int_{-1}^{2x} \frac{dt}{\tan^2 t}$  ✓

(c)  $g'(x) = -\frac{2}{\tan^2(2x)} \times 2 = -\frac{4}{\tan^2 2x}$  ✓✓ [9]

## Calculator-free Solutions

15. (a)  $f'(x) = \frac{\sqrt{3}}{4} + \frac{1}{2} \cos\left(\frac{x}{2}\right)$  ✓

TP occurs when  $f'(x) = 0$  ✓

$\frac{1}{2} \cos\left(\frac{x}{2}\right) = -\frac{\sqrt{3}}{4} \rightarrow \cos\left(\frac{x}{2}\right) = -\frac{\sqrt{3}}{2}$  ✓

$\frac{x}{2} = \frac{5\pi}{6} \rightarrow x = \frac{5\pi}{3}$  km ✓✓

and height = 2.767 m ✓✓

(b) Max gradient occurs when  $f''(x) = 0$  ✓

$-\frac{1}{4} \sin\left(\frac{x}{2}\right) = 0$  ✓

$x = 0, 6.28, 12.57, 18.85$  ✓✓

Max gradient = 0.93 ✓✓

[8]

16. (a)  $f'(x) = \sin x + x \cos x$  ✓✓

(b) (i)  $\int f'(x) dx = \int \sin x dx + \int x \cos x dx$  ✓✓

$f(x) = -\cos x + \int x \cos x dx$  ✓✓

$\int x \cos x dx = f(x) + \cos x + c$  ✓✓

$\int x \cos x dx = x \sin x + \cos x + c$  ✓✓

(ii)  $\int x \cos x dx = [x \sin x + \cos x]_0^{\pi}$  ✓✓

$= (0-1) - (0+1) = -2$  ✓✓

(c)  $\int_0^{\pi} |x \cos x| dx = 3.14$  ✓✓ [9]

## Calculator-free Solutions

(a)

TP

occurs

when

 $f'(x) = 0$ 

i.e.

 $\frac{1}{2} \cos\left(\frac{x}{2}\right) = -\frac{\sqrt{3}}{4}$ 

i.e.

 $\cos\left(\frac{x}{2}\right) = -\frac{\sqrt{3}}{2}$ 

i.e.

 $\frac{x}{2} = \frac{5\pi}{6}$ 

i.e.

 $x = \frac{5\pi}{3}$ 

km

✓✓

and

height

= 2.767

m

✓✓

[8]

(b)

We

would

need

to know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[9]

(c)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[10]

(d)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[11]

(e)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[12]

(f)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[13]

(g)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[14]

(h)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[15]

(i)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[16]

(j)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[17]

(k)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[18]

(l)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[19]

(m)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[20]

(n)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[21]

(o)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[22]

(p)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[23]

(q)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓

[24]

(r)

We

will

need

to

know

where

 $h(t)$ 

lies

below

the

x-axis

✓