

The release date for this exam and marking scheme is 14th June.

This examination is Copyright but may be freely used within the school that purchases this license.
The items that are contained in this examination are to be used solely in the school for which they
are purchased.
• They are not to be shared in any manner with a school which has not purchased their own license.
• The terms and the solutions/marketing keys are to be kept confidentially and not copied or made
available to anyone who is not a teacher at the school. Teachers may give feedback to students in
the form of showing them how the work is marked but students are not to retain a copy of the
paper or marking guide until the agreed release date stipulated in the purchasing
agreement/license.

Licence Agreement

© MAWA, 2019

MATHMATICS METHODS

MAWA Semester 1 (Unit 3) Examination 2019

Calculator-free

Marking Key

Section One: Calculator-free**(50 Marks)****Question 1(a)****(2 marks)**

Solution	
<i>Let $f(x) = xe^{3x}$</i>	
$f'(x) = e^{3x} + x3e^{3x} = e^{3x} + 3xe^{3x}$	
Mathematical behaviours	Mark
• applies product rule	1
• differentiates exponential correctly	1

Question 1(b)**(3 marks)**

Solution	
$\frac{d}{dx} \left(\frac{\cos x}{x^3} \right) = \frac{x^3(-\sin x) - \cos x(3x^2)}{(x^3)^2} = \frac{-x^2(x\sin x + 3\cos x)}{x^6} = -\frac{(x\sin x + 3\cos x)}{x^4}$	
Mathematical behaviours	Marks
• applies quotient rule	1
• differentiates $\cos x$ correctly	1
• simplifies result	1

Question 1(c)**(3 marks)**

Solution	
$g(u) = \sqrt{u} \Rightarrow \frac{dg}{du} = \frac{1}{2}u^{-\frac{1}{2}}$	
$u = 2 - 3x^2 \Rightarrow \frac{du}{dx} = -6x$	
$\Rightarrow \frac{dg}{dx} = \frac{1}{2}u^{-\frac{1}{2}} \times -6x = \frac{-3x}{\sqrt{2-3x^2}}$	
Mathematical behaviours	Marks
• states $\frac{dg}{du}$	1
• states $\frac{du}{dx}$	1
• states $\frac{dg}{dx}$	1
• states $\frac{dg}{dx}$ in terms of x .	

Question 1(d)**(3 marks)**


Solution	
$x(t) = 3\sin 2t \Rightarrow v(t) = 3 \times 2\cos 2t$	
$v(t) = 0 \Rightarrow \cos 2t = 0$	
$ie 2t = \frac{\pi}{2} \Rightarrow t = \frac{\pi}{4}s$	
Mathematical behaviours	Marks
• differentiates to obtain $v(t)$	1
• equates $v(t) = 0$	1
	1

- circles the 2nd graph

Question 3 (4 marks)

Solution	
$\text{Area} = \int_0^{2\pi} [\sin x - x(x - 2\pi)] dx$ $= \int_0^{2\pi} (\sin x - x^2 + 2\pi x) dx$ $= \left[-\cos x - \frac{x^3}{3} + \pi x^2 \right]_0^{2\pi}$ $= \left[-\cos 2\pi - \frac{(2\pi)^3}{3} + \pi(2\pi)^2 \right] - [-\cos 0]$ $= -1 - \frac{8\pi^3}{3} + 4\pi^3 + 1$ $= \frac{4\pi^3}{3}$	

Mathematical behaviours	Marks
• states a correct expression using integrals to determine the area	1
• anti-differentiates each part correctly	1
• substitutes in limits of integration	1
• evaluates result	1



- rearranges to get required result

Question 7(c) (3 marks)

Solution	
$\int_0^{\frac{\pi}{6}} (\sin x \cos x + 2) dx = \left[\frac{1}{2} \sin^2 x + 2x \right]_0^{\frac{\pi}{6}}$ $= \frac{1}{2} \left[\left(\frac{1}{2} \right)^2 - 0^2 \right] + 2 \left[\frac{\pi}{6} - 0 \right]$ $= \frac{1}{8} + \frac{\pi}{3}$	

Mathematical behaviours	Marks
• recognises $\sin^2 x$ term is to be involved	1
• states correct integral and bounds of integration	1
• substitutes bounds of integration and simplifies	1

Marks	Mathematical behaviours	$\int_1^x dt = \int_1^{\sqrt{1-t}} dt$	• uses the relationship $x = \sqrt{1-t}$	• applies Fundamental Theorem of Calculus	1
Marks	Mathematical behaviours	$\int_1^x dt = \int_1^{\sqrt{1-t}} dt$	• uses the relationship $x = \sqrt{1-t}$	• applies Fundamental Theorem of Calculus	1
Solution					(2 marks)
Question 4(c)					(2 marks)

Marks	Mathematical behaviours	$\int_1^3 (3 - 2x)^2 dx = \int_1^3 (1 - 3x)^2 dx = \frac{1}{3} [1 - 3x]^3 \Big _1^3 = \frac{13}{6} = \frac{26}{6} = \frac{13}{3}$	• substitutes limits of integration and evaluates	• anti-differentiates correctly	1
Marks	Mathematical behaviours	$\int_1^3 (3 - 2x)^2 dx = \int_1^3 (1 - 3x)^2 dx = \frac{1}{3} [1 - 3x]^3 \Big _1^3 = \frac{13}{6} = \frac{26}{6} = \frac{13}{3}$	• substitutes limits of integration and evaluates	• anti-differentiates correctly	1
Solution					(2 marks)
Question 4(b)					(2 marks)

Marks	Mathematical behaviours	$\int_{-1}^2 e^{2x} - 3x^2 dx = \int_{-1}^2 2e^{2x} - 3x^2 dx = \frac{1}{2} e^{2x} - 3x^2 \Big _{-1}^2 = \frac{1}{2} e^4 - 3(2)^2 - \left(\frac{1}{2} e^{-2} - 3(-1)^2 \right) = e^4 - 6\sqrt{e^2} + C$	• anti-differentiates the square root function correctly	• anti-differentiates the exponential function correctly	1
Marks	Mathematical behaviours	$\int_{-1}^2 e^{2x} - 3x^2 dx = \int_{-1}^2 2e^{2x} - 3x^2 dx = \frac{1}{2} e^{2x} - 3x^2 \Big _{-1}^2 = \frac{1}{2} e^4 - 3(2)^2 - \left(\frac{1}{2} e^{-2} - 3(-1)^2 \right) = e^4 - 6\sqrt{e^2} + C$	• anti-differentiates the square root function correctly	• anti-differentiates the exponential function correctly	1
Solution					(2 marks)
Question 4(a)					(2 marks)

MATHEMATICS METHODS	CALCULATOR-FREE	SEMESTER 1 (UNIT 3) EXAMINATION
5		SEMINAR

Solution	$y = \sin^2 x$	$\frac{dy}{dx} = 2 \sin x \cos x$	$y = \int 2 \sin x \cos x dx + C$	$\int 2 \sin x \cos x dx = \frac{d}{dx} (\sin^2 x) + C$	• applies fundamental theorem
Solution	$y = \sin^2 x$	$\frac{dy}{dx} = 2 \sin x \cos x$	$y = \int 2 \sin x \cos x dx + C$	$\int 2 \sin x \cos x dx = \frac{d}{dx} (\sin^2 x) + C$	• applies fundamental theorem
Question 7(b)					(3 marks)
Solution					(3 marks)

Solution	$y = \frac{6}{2 + \sqrt{3}}$	$= \frac{6}{\sqrt{4 + 2\sqrt{3}}} \div 2$	$= \frac{6}{\sqrt{2 + \sqrt{3}}} \div 2$	$i.e. \text{Estimated area under } f(x) = \sin x \text{ from } x = 0 \text{ to } x = \frac{\pi}{2} \text{ is}$	• simplifies to deduce the required result
Solution	$y = \frac{6}{2 + \sqrt{3}}$	$= \frac{6}{\sqrt{2 + \sqrt{3}}} \div 2$	$= \frac{6}{\sqrt{2 + \sqrt{3}}} \div 2$	$i.e. \text{Estimated area under } f(x) = \sin x \text{ from } x = 0 \text{ to } x = \frac{\pi}{2} \text{ is}$	• determines the average of the two areas obtained in part (a)
Question 7(a)					(1 mark)
Solution					(1 mark)

Solution	$\frac{1}{3} + \frac{\sqrt{3}}{2}$	$= \frac{6}{\sqrt{2 + \sqrt{3}}} \div 2$	$= \frac{6}{\sqrt{2 + \sqrt{3}}} \div 2$	$= \frac{6}{\sqrt{2 + \sqrt{3}}} \div 2$	• states the sum of the three rectangles and simplifies correctly
Solution	$\frac{1}{3} + \frac{\sqrt{3}}{2}$	$= \frac{6}{\sqrt{2 + \sqrt{3}}} \div 2$	$= \frac{6}{\sqrt{2 + \sqrt{3}}} \div 2$	$= \frac{6}{\sqrt{2 + \sqrt{3}}} \div 2$	• states the sum of the three rectangles and simplifies correctly
Question 6(b)					(2 marks)
Solution					(2 marks)

MATHEMATICS METHODS	CALCULATOR-FREE	SEMESTER 1 (UNIT 3) EXAMINATION
8		SEMINAR

Question 4(d)

(3 marks)

Solution	
$\int_{-m}^m (m^3 - x^3) dx = 1250$	
$\left[m^3 x - \frac{x^4}{4} \right]_{-m}^m = 1250$	
$\left[m^4 - \frac{m^4}{4} \right] - \left[-m^4 - \frac{m^4}{4} \right] = 1250$	
$\frac{3m^4}{4} + \frac{5m^4}{4} = 1250$	
$\frac{8m^4}{4} = 1250$	
$2m^4 = 1250$	
$m^4 = 625$	
$m = \pm 5$	
Mathematical behaviours	Marks
• anti-differentiates integral correctly	1
• substitutes in limits of integration correctly and simplifies to obtain correct expression on the LHS	1
• determines correct answers for m .	1

Question 5(a)

(3 marks)

Solution	
$\mu = \frac{1}{36}, \sigma^2 = \frac{1}{36} \times \frac{35}{36} = \frac{35}{36^2}$	
Bernoulli distribution with	
Mathematical behaviours	Marks
• states Bernoulli	1
• states mean	1
• states variance	1

Question 5(b)

(3 marks)

Solution	
This represents a Binomial with $n=15$ and $p = \frac{1}{36}$.	
Mathematical behaviours	Marks
• states Binomial	1
• states n	1
• states p	1

Question 5(c)

(1 mark)

Solution	
$W \sim \text{Bin}\left(15, \frac{1}{36}\right)$	
$P(W=1) = {}^{15}C_1 \left(\frac{1}{36}\right)^1 \left(\frac{35}{36}\right)^{14} = 15 \left(\frac{1}{36}\right)^1 \left(\frac{35}{36}\right)^{14}$	
Mathematical behaviours	Marks
• states correct expression	1

Question 5(d)

(3 marks)

Solution	
$Let Z \sim \text{Bin}\left(30, \frac{1}{36}\right)$	
$P(Z=2 Z \geq 1) = \frac{P(Z=2)}{P(Z \geq 1)} = \frac{P(Z=2)}{1 - P(Z=0)} = \frac{{}^{30}C_2 \left(\frac{1}{36}\right)^2 \left(\frac{35}{36}\right)^{28}}{1 - {}^{30}C_0 \left(\frac{1}{36}\right)^0 \left(\frac{35}{36}\right)^{30}}$	
Mathematical behaviours	Marks
• recognises the situation involves a binomial $\left(\frac{1}{36}\right)$ and conditional probability	1
• states correct expression for numerator	1
• states correct expression for denominator	1

Question 6(a)

(1 mark)

Solution	
(i) Under-estimated Area = $\left(\frac{\pi}{6} \times \frac{1}{2}\right) + \left(\frac{\pi}{6} \times \frac{\sqrt{3}}{2}\right)$	
$= \frac{\pi}{6} \left(\frac{1+\sqrt{3}}{2}\right)$	
(ii) Over-estimated Area = $\left(\frac{\pi}{6} \times \frac{1}{2}\right) + \left(\frac{\pi}{6} \times \frac{\sqrt{3}}{2}\right) + \left(\frac{\pi}{6} \times 1\right)$	
$= \frac{\pi}{6} \left(\frac{1+\sqrt{3}+2}{2}\right)$	
$= \frac{\pi}{6} \left(\frac{3+\sqrt{3}}{2}\right)$	
Mathematical behaviours	Marks
(i)	
• states the sum of the area of the two rectangles and simplifies correctly	1
(ii)	