


Note on marking:
-1 mark at most in Section Two for missing units
-1 mark at most in Section Two for incorrect rounding



Semester One Examination, 2018

Question/Answer booklet

SOLUTIONS

MATHEMATICS

METHODS

UNIT 1

Section Two:

Calculator-assumed

In words

Your name

**Time allowed for this section**

Reading time before commencing work: ten minutes

Working time: one hundred minutes

**Materials required/recommended for this section**

This Question/Answer booklet

Formula sheet (retained from Section One)

*To be provided by the supervisor*

**Standard items:** pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

**Special items:** drawing instruments, templates, notes on one unfolded sheet of A4 paper (both sides), and up to three calculators approved for use in this examination

**Important note to candidates**

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised material. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

**Structure of this paper**

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	8	8	50	52	35
Section Two: Calculator-assumed	14	14	100	98	65
<b>Total</b>					100

**Instructions to candidates**

1. The rules for the conduct of examinations are detailed in the school handbook. Sitting this examination implies that you agree to abide by these rules.
2. Write your answers in this Question/Answer booklet.
3. You must be careful to confine your response to the specific question asked and to follow any instructions that are specified to a particular question.
4. Supplementary pages for the use of planning/continuing your answer to a question have been provided at the end of this Question/Answer booklet. If you use these pages to continue an answer, indicate at the original answer where the answer is continued, i.e. give the page number.
5. Show all your working clearly. Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
6. It is recommended that you do not use pencil, except in diagrams.
7. The Formula sheet is not to be handed in with your Question/Answer booklet.

Additional working space

Question number: \_\_\_\_\_

Additional working space

Question number: \_\_\_\_\_


65% (98 Marks)

**Section Two: Calculator-assumed**  
This section has **fourteen (14)** questions. Answer **all** questions. Write your answers in the spaces provided.

Working time: 100 minutes.

**Question 9** (3 marks)

The graphs of  $y = 2x^2 + 2x + c$ ,  $y = a(x - 2)^2 + 1$  and  $y = (x + b)(x + 3)$  are shown below.



Determine the values of the constants  $a$ ,  $b$  and  $c$ .

Solution	
$x = 4 \Rightarrow 0 = a(4 - 2)^2 + 1$	$a = -\frac{1}{4}$
$b = -1$ (Other root at -3)	
$c = -1$ (y-intercept)	
Specific behaviours	
✓ value of $a$	
✓ value of $b$	
✓ value of $c$	

SN018-112-8

See next page

**Question 10****(5 marks)**

An online grocery is offering new customers the opportunity to select 8 different products for just \$2 each. They can select from a range of 12 different canned items, 14 different snacks and 13 different drinks.

- (a) Determine how many different selections can be made. (1 mark)

Solution
$12 + 14 + 13 = 39$
${}^{39}C_8 = 61\,523\,748$
Specific behaviours
✓ correct number

- (b) Determine how many different selections can be made that just include drinks. (1 mark)

Solution
${}^{13}C_8 = 1\,287$
Specific behaviours
✓ correct number

In a separate offer, the online grocery forms a special bargain pack containing 4 different canned items and 3 different snacks.

- (c) How many different special bargain packs can be formed? (3 marks)


Solution
${}^{12}C_4 \times {}^{14}C_3 = 180\,180$
Specific behaviours
✓ indicates choosing 4 from 12
✓ indicates choosing 3 from 14
✓ calculates product of these

Additional working space

Question number: \_\_\_\_\_

Question 11

Calculate the value of the distance  $x$  in the following triangle.



Solution
using cosine rule: $4^2 = x^2 + 5^2 - 2 \times (5) \times (x) \times \cos(35)$ CAS SOLVE: $x = 1.31 \text{ cm}$ or $6.88 \text{ cm}$
Specific behaviours
✓ gives appropriate equation for $x$ ✓ gives first possible value of distance $x$ , 1.31 cm ✓ gives second possible value of distance $x$ , 6.88 cm

See next page

SN018-112-8

Question 22

(a) Use your calculator to

(i) determine the exact value of  $\cos 36^\circ$ .

(ii) determine the exact value of  $\sin 105^\circ$ .

(1 mark)

(1 mark)

Solution
(i) $\frac{\sqrt{5} + 1}{4}$ (ii) $\frac{\sqrt{2}(\sqrt{3} + 1)}{4}$ ✓ each exact value
Specific behaviours

(iii) solve  $\cos(x + 60^\circ) = \sin x$  for  $-270^\circ \leq x \leq 270^\circ$ .

(1 mark)

Solution
$x = -165^\circ$ , $x = 15^\circ$ , $x = 195^\circ$
Specific behaviours
✓ all three correct solutions for $x$ <i>NB Graphical or numerical solve quickest - use of exact solve slow</i>

(b) Using suitable exact values of acute angles and an angle sum and difference identity, justify your above value of  $\sin 105^\circ$ . (5 marks)

Solution
$\sin(105) = \sin(45 + 60)$ $= \sin 45 \cos 60 + \cos 45 \sin 60$ $= \frac{\sqrt{2}}{2} \times \frac{1}{2} + \frac{\sqrt{2}}{2} \times \frac{\sqrt{3}}{2}$ $= \frac{\sqrt{2}(1 + \sqrt{3})}{4}$ ✓ chooses suitable pair of angles ✓ chooses correct identity ✓ first pair of exact values ✓ second pair of exact values ✓ simplifies to match previous answer
Specific behaviours

End of questions

SN018-112-8

## Question 12

(6 marks)

- (a) Determine the equation of the axis of symmetry for the graph of  $y = 3x^2 + 12x + 40$ .  
(2 marks)

Solution
$x = -\frac{b}{2a} = -\frac{12}{2 \times 3} = -2$ $x = -2$
Specific behaviours
<ul style="list-style-type: none"> <li>✓ indicates use of formula</li> <li>✓ correct equation</li> </ul>

- (b) The graph of  $y = ax^2 + bx + 13$  passes through the points  $(-3, -23)$  and  $(4, 5)$ .  
Determine the values of the constants  $a$  and  $b$ .  
(4 marks)

Solution
$-23 = (-3)^2 a - 3b + 13$ $-23 = 9a - 3b + 13$ $5 = 4^2 a + 4b + 13$ $5 = 16a + 4b + 13$ <p>Solve simultaneously using CAS</p> $a = -2, \quad b = 6$
Specific behaviours
<ul style="list-style-type: none"> <li>✓ substitutes first point</li> <li>✓ substitutes second point</li> <li>✓ solves for <math>a</math></li> <li>✓ solves for <math>b</math></li> </ul>

## Question 21 (continued)


- (c) Calculate the value(s) of  $x$  when  $k = -1.2$ .  
(2 marks)

Solution
$x = \frac{2 \pm \sqrt{4 - 2(-1.2)^2}}{2(-1.2)}$ $x = -1.27, -0.39$
Specific behaviours
<ul style="list-style-type: none"> <li>✓ gives <math>x = -1.27</math></li> <li>✓ gives <math>x = -0.39</math></li> </ul>

(7 marks)

Let  $f(x) = \frac{3-x}{4}$  and  $g(x) = \frac{x}{1} + q$ , where  $p$  and  $q$  are constants.

The graph of  $y = g(x)$  is shown below.



(a) Sketch the graph of  $y = f(x)$  on the axes above. (3 marks)

<b>Solution</b>
$p = 2, \quad q = -3$
<b>Specific behaviours</b>
✓ value of $p$
✓ value of $q$

(b) Determine the values of  $p$  and  $q$ . (2 marks)

(c) Solve the equation  $f(x) = g(x)$ , giving your solution(s) to one decimal place. (2 marks)

<b>Solution</b>
$x = -1.7, \quad x = 4.4$
<b>Specific behaviours</b>
✓ one solution
✓ second solution
(Rounding for guidance only but penalise answers given as coordinates)
Allow follow through marks if different $p$ and $q$ values are given in (b)

See next page

SN018-112-8

Consider the equation

Question 21

$$2kx^2 - 4x + k = 0$$

(a) Solve this equation for  $x$ , giving your answer as a simplified expression in terms of  $k$ . (3 marks)

<b>Solution</b>
$x = \frac{4 \pm \sqrt{16 - 4 \times 2k \times k}}{4k}$ $x = \frac{4 \pm \sqrt{16 - 8k^2}}{4k}$ $x = \frac{4 \pm 2\sqrt{4 - 2k^2}}{4k}$ $x = \frac{4k}{2 \pm \sqrt{4 - 2k^2}}$ $x = \frac{2k}{2 \pm \sqrt{4 - 2k^2}}$
<b>Specific behaviours</b>
✓ correct initial use of quadratic formula
✓ correct initial simplification
✓ correct simplified final expression
including taking out factor of 2 from square root

(b) Give the value(s) of  $k$  for which the equation has exactly one solution. (2 marks)

<b>Solution</b>
For one solution: $4 - 2k^2 = 0$ $k^2 = 2$ $k = \pm\sqrt{2}$
<b>Specific behaviours</b>
✓ correct initial equation
✓ both correct final values of $k$

See next page

SN018-112-8

## Question 14

(12 marks)


During 2018, the altitude of the sun,  $\theta$  degrees, at noon in Paris on the  $n^{\text{th}}$  day of the year can be modelled by the equation

$$\theta = 23.5 \sin\left(\frac{8\pi(n + 283)}{1461}\right) + 41.1$$

- (a) On the 30th of January, the altitude was  $22.7^\circ$ . Calculate the altitude twelve days later. (2 marks)

Solution	
$n = 42,$	$\theta = 26.1^\circ$
Specific behaviours	
✓ indicates $n$	
✓ correct angle	

- (b) Graph the altitude on the axes below for  $0 \leq n \leq 365$ . (4 marks)



Solution	
See graph	
Specific behaviours	
✓ endpoints, $\theta \approx 18^\circ$	
✓ maximum close to (174, 65°)	
✓ minimum close to right endpoint	
✓ smooth curve	

See next page

SN018-112-8

## Question 20 (continued)

- (d) the area of triangle  $ABC$ . (4 marks)

Solution	
$\angle A + \angle C = 180 - 80$	
$\angle A + 3\angle A = 100 \Rightarrow \angle A = 25$	
$\frac{AC}{\sin 80} = \frac{2 \times 32}{\sin 25}$	
$AC = 149.1$	
$\text{Area} = \frac{1}{2}(64)(149.1) \sin(3 \times 25) \approx 4610 \text{ cm}^2$	
Specific behaviours	
✓ indicates use of equation to find second angle	
✓ evaluates second angle and indicates use in sin rule	
✓ evaluates second side	
✓ evaluates triangle area	

See next page

SN018-112-8



Question 14 (continued)

- (c) State the maximum altitude of the sun at noon in Paris and on which day of the year this occurred. (2 marks)

<b>Solution</b>
$\theta_{MIN} = 64.6^\circ$ on day 174
<b>Specific behaviours</b>
✓ altitude, correct to 1 dp ✓ day of year, rounded to whole number

- (d) Solar panels on the roof of a Paris apartment are designed to meet its entire power needs on cloudless days when the altitude of the sun is at least  $30^\circ$  at noon. (1 mark)

<b>Solution</b>
See graph - horizontal line $\theta = 30^\circ$
<b>Specific behaviours</b>
✓ neat, straight line

- (iii) Determine the number of days the panels are not expected to achieve this aim during 2018, ignoring the possibility of cloud cover. (3 marks)

<b>Solution</b>
$\theta > 30^\circ \Rightarrow 54 \leq n \leq 293$
$53 + (294 \text{ to } 365)$
$53 + 72 = 125 \text{ days}$
<b>Specific behaviours</b>
✓ one value of $n$ ✓ second value of $n$ ✓ correct number of days


See next page

SN018-112-8

Question 20

(10 marks)

A logo with triangular outline  $ABC$  contains a shaded segment bounded by the straight line  $PM$  and the circular arc  $PM$  with centre  $B$  and radius  $BM = 32$  cm, as shown below.



Given that  $\angle ABC = \frac{4\pi}{9}$ ,  $\angle BCA = 3\angle BAC$  and  $M$  is the midpoint of  $BC$ , determine

- (a) the size of  $\angle ABC$  in degrees. (1 mark)

<b>Solution</b>
$\frac{4\pi}{180} \times \frac{\pi}{9} = 80^\circ$
<b>Specific behaviours</b>
✓ converts angle

- (b) the area of the shaded segment. (2 marks)

<b>Solution</b>
$A = \frac{1}{2}(32)^2\left(\frac{4\pi}{9} - \sin\left(\frac{4\pi}{9}\right)\right) \approx 210.7 \text{ cm}^2$
<b>Specific behaviours</b>
✓ indicates substitution into segment area formula ✓ evaluates area

- (c) the perimeter of the shaded segment. (3 marks)

<b>Solution</b>
$PM_{arc} = 32 \times \frac{4\pi}{128\pi} = \frac{9}{9} \approx 44.68$
$b = \sqrt{32^2 + 32^2 - 2(32)(32)\cos 80} \approx 41.14$
Perimeter $= 44.68 + 41.14 \approx 85.8 \text{ cm}$
<b>Specific behaviours</b>
✓ calculates arc length ✓ indicates use of cosine rule to find $PM$ ✓ evaluates $PM$ and states perimeter

See next page



SN018-112-8

Question 15 (9 marks)

(a) The graphs of the following, where  $a, b, c, d, e$  and  $f$  are constants, are shown in the two axes grids below.

$y = \sin(ax)$     $y = b \cos(x)$     $y = \tan(cx)$     $y = d \sin(x)$     $y = \cos(ex)$     $y = f \tan(x)$

Three of the graphs are in the first axes grid and three of the graphs are in the second axes grid.



State the values of  $a, b, c, d, e$  and  $f$ .


Solution	
See table	
Specific behaviours	
✓ each value	

(6 marks)

Constant	Value
$a$	2
$b$	-4
$c$	0.5
$d$	5
$e$	4
$f$	-1

Question 19 (7 marks)

Consider the points with coordinates  $(a, b)$  and  $(c, d)$  that lie in the first and second quadrants respectively of the unit circles shown below, where  $\mu$  and  $\lambda$  are acute angles.



Determine the following in terms of  $a, b, c$  and  $d$ , simplifying your answers where possible.

(a)  $\cos \lambda$ .

Solutions	
(i) $-c$	
(ii) $\frac{b}{a}$	
(iii) $-d$	
(iv) $b$	
Specific behaviours	
✓ each correct response	

(1 mark)

(b)  $\tan(180^\circ + \mu)$ .

(1 mark)

(c)  $\sin(\pi + \lambda)$ .

(1 mark)

(d)  $\cos(90^\circ - \mu)$ .

(1 mark)

(e)  $\sin(\mu - \lambda)$ .

Solution	
$\sin(\mu - \lambda) = \sin \mu \cos \lambda - \cos \mu \sin \lambda$ $= (b)(-c) - (a)(d)$ $= -bc - ad$	
Specific behaviours	
✓ uses identity ✓ at least two correct trig values ✓ correct expression	

(3 marks)

Question 15 (continued)

(b) Calculate the acute angle in degrees between the lines  $y = x + 5$  and  $y = 3x - 1$ , rounding your answer to one decimal place. (3 marks)

Solution
$\alpha = \tan^{-1}(1) = 45^\circ$
$\beta = \tan^{-1}(3) = 71.565^\circ$
$\beta - \alpha = 26.6^\circ$ (1 dp)
Specific behaviours
✓ angle of inclination of first line
✓ angle of inclination of second line
✓ acute angle, to one decimal place

See next page

SN018-112-8

Question 18

The wind speed at a weather station,  $v$  metres per second,  $t$  hours after recording began, can be modelled by the function


$$v = 20 - 5.8t + 0.75t^2 - 0.02t^3, 0 \leq t \leq 24$$

(a) Calculate the wind speed when  $t = 11$ .

Solution
$v(11) = 20.33$ m/s
Specific behaviours
✓ value

(1 mark)

(b) Sketch the graph of wind speed against time on the axes below. (4 marks)



(c) During the 24-hour period, determine

(i) the time at which the wind speed was greatest.

Solution
$t = 20.2$ h
Specific behaviours
✓ value (at least 1 dp)

(1 mark)

(ii) the minimum wind speed.

Solution
$v_{MIN} \approx 7.23$ m/s
Specific behaviours
✓ value (at least 1 dp)

(iii) the length of time, in hours and minutes, that the wind speed was increasing.

Solution
$20.219 - - 4.781 = 15.438$
$= 15$ h 26 min
Specific behaviours
✓ interval in hours
✓ interval in hours and minutes


See next page

SN018-112-8

## Question 16

(6 marks)

- (a) The graph of  $y = f(x)$  is shown in bold below. The graphs of  $y = -f(x)$ ,  $y = f(x + p)$  and  $y = f(x) + q$  are also shown, where  $p$  and  $q$  are constants.



Complete the table below giving the equation of each of the graphs A, B and C as one of  $y = -f(x)$ ,  $y = f(x + p)$  or  $y = f(x) + q$ .

Graph	Equation
A	<b>SOLUTION: <math>y = f(x) + q</math></b> ✓
B	<b>SOLUTION: <math>y = f(x + p)</math></b> ✓
C	<b>SOLUTION: <math>y = -f(x)</math></b> ✓

Solution
See table
Specific behaviours
✓ each value

(3 marks)

- (b) The one-to-one relation  $y = 9 - 2x$  has domain and range given by  $\{x: x = -4, a, 10\}$  and  $\{y: y = -11, -7, b\}$  respectively. Determine the values of constants  $a$  and  $b$ . (3 marks)

Solution
$x = 10, y = -11$
$x = -4, y = 17 = b$
$x = a, y = 9 - 2a = -7 \Rightarrow a = 8$
Specific behaviours
✓ value of $b$
✓ indicates $a$ is mapped onto $-7$
✓ solves for value of $a$


See next page

SN018-112-8

## Question 17

(6 marks)

- (a) Part of the graph of  $y = f(x)$  is shown below, where  $f(x) = 3(x + b)^3 + c$ , and  $b$  and  $c$  are constants.



- (i) State the degree of  $f(x)$ .

Solution
3
Specific behaviours
✓ correct degree

(1 mark)

- (ii) Determine the value of  $b$ .

Solution
$b = 3$
Specific behaviours
✓ correct value

(1 mark)

- (iii) Determine  $f(0)$ .

Solution
$f(x) = 3(x + 3)^3 + 1$
$f(0) = 3(3)^3 + 1 = 82$
Specific behaviours
✓ indicates value of $c$
✓ evaluates

(2 marks)

- (b) Another function,  $g(x)$ , is a transformation of  $f(x)$ , where  $g(x) = f(x - 7)$ .

Describe how to obtain the graph of  $y = g(x)$  from the graph of  $y = f(x)$ .

(2 marks)

Solution
Translate graph 7 units to the right.
Specific behaviours
✓ uses translation
✓ indicates distance and direction

See next page

SN018-112-8