

Course Methods Test 3 Year 12

Student name: _____ Teacher name: _____

Task type: Response/Investigation

Reading time for this test : 5 mins

Working time allowed for this task: 40 mins

Number of questions: _____6_____

Materials required: No classpads

Standard items: Pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

Special items: Drawing instruments, templates, notes on one unfolded sheet of A4 paper

Marks available: _____38_____ marks

Task weighting: _____14_____ %

Formula sheet provided: No but some formulae given on page 2

Note: All part questions worth more than 2 marks require working to obtain full marks.

End of test
Working out space

Useful formulae

Logarithms

$x = \log_a b \Leftrightarrow a^x = b$	$a^{\log_a b} = b$ and $\log_a(a^b) = b$
$\log_a mn = \log_a m + \log_a n$	$\log_a \frac{m}{n} = \log_a m - \log_a n$
$\log_a(m^k) = k \log_a m$	$\log_e x = \ln x$

$\frac{d}{dx} \ln x = \frac{1}{x}$	$\int \frac{1}{x} dx = \ln x + c, \quad x > 0$
$\frac{d}{dx} \ln f(x) = \frac{f'(x)}{f(x)}$	$\int \frac{f'(x)}{f(x)} dx = \ln f(x) + c, \quad f(x) > 0$
Product rule	<div> <div>If $y = uv$</div> <div>then</div> <div>$\frac{d}{dx}(uv) = v \frac{du}{dx} + u \frac{dv}{dx}$</div> </div> <div> <div>If $y = f(x) g(x)$</div> <div>then</div> <div>$y' = f'(x) g(x) + f(x) g'(x)$</div> </div>
Quotient rule	<div> <div>If $y = \frac{u}{v}$</div> <div>then</div> <div>$\frac{d}{dx} \left(\frac{u}{v} \right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$</div> </div> <div> <div>If $y = \frac{f(x)}{g(x)}$</div> <div>then</div> <div>$y' = \frac{f'(x) g(x) - f(x) g'(x)}{(g(x))^2}$</div> </div>
Chain rule	<div> <div>If $y = f(u)$ and $u = g(x)$</div> <div>then</div> <div>$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$</div> </div> <div> <div>If $y = f(g(x))$</div> <div>then</div> <div>$y' = f'(g(x)) g'(x)$</div> </div>
Fundamental theorem	$\frac{d}{dx} \left(\int_a^x f(t) dt \right) = f(x)$ and $\int_a^b f(x) dx = f(b) - f(a)$

Q6 continued

iii) $E(X)$ i.e the mean. (No need to simplify)b) Derive the cumulative probability function $P(X \leq x)$ for $0 \leq x \leq 10$.


Q1 (2 & 2 = 4 marks)
Express each of the following as a **single logarithm**.
a) $\log_a b + 3\log_a(ab) - 4\log_a b$.

b) $5 + 3\log_5 c - \log_5(c^3) + \log_5 b$.

Q2 (2 & 2 = 4 marks)
Solve each of the following, giving your answer in **exact** form.
a) $2^{2x} - 12(2^x) + 32 = 0$

b) $7^x + 3(7^{x+2}) = 31$

Q6 (3, 3, 3 & 3 = 12 marks)
Consider the continuous random variable X and its probability density function which is graphed below.



a) Determine the following **exactly**.
i) $P(2 < X < 7)$. (Simplify)

iii) $P(X < 3 | X < 5)$. (No need to simplify)

Q3 (1, 3 & 3 = 7 marks)

The Richter scale, R , of an earthquake of intensity I is given by $R = \log_{10} \left(\frac{I}{I_o} \right)$ where I_o is a

minimum intensity level used for comparison.


a) Determine R for an earthquake with intensity $10000I_o$.

b) An earthquake measuring 5 on the Richter scale is how many times as intense as that of one measuring 4 on the Richter scale?

c) If an earthquake registers x on the Richter scale and a second earthquake registers $x + 4$ on the Richter scale, how many more times as intense is the second earthquake?

Q4 (3 marks)

Consider the function $f(x) = \log_2 x$ which undergoes a transformation $f(ax+b)$ where a & b are constants. The graph $y = f(ax+b)$ is plotted below, determine the values of a & b showing reasoning.



Q5 (3 & 5 = 8 marks)

Consider the function $g(x) = (x^2 + 3) \ln(x^3 + 3x)$.

a) Determine $g'(x)$. (Simplify)

b) Use the result from part a to determine $\int 2x \ln(x^3 + x) dx$.