

Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until Wednesday June 22nd 2018.



MATHEMATICS SPECIALIST UNIT 1

Semester One

2018

SOLUTIONS

Calculator-free Solutions

$$1. \quad (a) \quad b - a = \begin{pmatrix} 1 \\ 3 \end{pmatrix} - \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} -1 \\ 2 \end{pmatrix} \quad \checkmark$$

$$\therefore a \cdot (b - a) = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} -1 \\ 2 \end{pmatrix} = -2 + 2 = 0 \quad \checkmark\checkmark$$

$$\therefore a \perp (b - a)$$

$$(b) \quad (i) \quad \vec{OC} = \vec{AB} = b - a = -i + 2j \quad \checkmark$$

$$(ii) \quad \vec{AC} = \vec{OC} - \vec{OA} = -3i + j \quad \checkmark$$

$$\therefore |\vec{AC}| = \sqrt{10} \quad \checkmark$$

OR

$$|\vec{AC}| = |\vec{OB}| \quad \checkmark$$

$$\therefore |\vec{AC}| = \left| \begin{pmatrix} 1 \\ 3 \end{pmatrix} \right| = \sqrt{10} \quad \checkmark$$

[6]

$$2. \quad (a) \quad (i) \quad \frac{13! + 12!}{13! - 12!} = \frac{13 \times 12! + 12!}{13 \times 12! - 12!} = \frac{12!(13+1)}{12!(13-1)} \quad \checkmark$$

$$\therefore \frac{14}{12} = \frac{7}{6} \quad \checkmark$$

$$(ii) \quad \frac{{}^{10}C_4}{{}^8C_4} = \frac{10!}{4! \times 6!} \div \frac{8!}{4! \times 4!} \quad \checkmark$$

$$\therefore \frac{10 \times 9 \times 8! \times 4! \times 4!}{8! \times 6 \times 5 \times 4!} = \frac{10}{5} \times \frac{9}{6} \therefore 2 \times \frac{3}{2} = 3 \quad \checkmark$$

$$(b) \quad \text{LHS} \quad \therefore k \binom{n}{k} = k \times \frac{n!}{k!(n-k)!} \quad \checkmark$$

$$\therefore \frac{k}{k} \times \frac{n!}{(k-1)![n-k]!} \quad \checkmark$$

$$\therefore n \times \frac{(n-1)!}{(k-1)![n-1-(k-1)]!} \quad \checkmark\checkmark$$


$$\therefore n \binom{n-1}{k-1} = \therefore \text{RHS} \quad (\text{Or from RHS to LHS})$$

[8]

3. (a) $-3 > -5$ but $(-3)^2 = 9 > (-5)^2 = 25$ is false ✓
- (b) "If the triangle is not equilateral, then the triangle does not have three equal sides." ✓
 Yes it is always true since the original implication is always true by definition of equilateral triangles. ✓
- (c) "If n is divisible by 3, then n is divisible by 6". ✓
 The converse is not always true, because for n to be divisible by 6 it must be divisible by both 2 and 3. ✓
- (d) FOR ALL natural numbers p , EXISTS a real number q , such that q is the square root of p . ✓✓

[7]

4. (a) $a+b = \begin{pmatrix} 2 \\ -1 \end{pmatrix} + \begin{pmatrix} -1 \\ 4 \end{pmatrix} = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$ ✓
- $\therefore \widehat{(a+b)} = \pm \frac{1}{\sqrt{10}} \begin{pmatrix} 1 \\ 3 \end{pmatrix}$ ✓
- $\therefore u = 2\sqrt{10} \times \pm \frac{1}{\sqrt{10}} \begin{pmatrix} 1 \\ 3 \end{pmatrix} = \pm \begin{pmatrix} 2 \\ 6 \end{pmatrix}$ ✓



- (b) (i) $c = k a$ ✓
- $\begin{pmatrix} -4 \\ \alpha \end{pmatrix} = k \begin{pmatrix} 2 \\ -1 \end{pmatrix} \rightarrow k = -2 \rightarrow \alpha = 2$ ✓

$$(ii) \quad \left| \begin{pmatrix} -4 \\ \alpha \end{pmatrix} - \begin{pmatrix} -1 \\ 4 \end{pmatrix} \right| = 3 \left| \begin{pmatrix} 2 \\ -1 \end{pmatrix} \right|$$

$$\therefore \sqrt{9 + (\alpha - 4)^2} = 3\sqrt{5}$$

$$9 + (\alpha - 4)^2 = 9 \times 5$$

$$(\alpha - 4)^2 = 36$$

$$\alpha = 4 \pm 6 \rightarrow \alpha = 10 \vee -2$$

$$5. \quad (a) \quad \overrightarrow{AC} \cdot \overrightarrow{OB} = (c - a) \cdot (a + c) = 0$$

$$c \cdot a + c \cdot c - a \cdot a - a \cdot c = 0$$

$$|c|^2 - |a|^2 = 0 \rightarrow |a| = |c|$$

\therefore OABC is a rhombus

$$(b) \quad \text{LHS } |AC|^2 + |OB|^2 = |c - a|^2 + |a + c|^2$$

$$(c - a) \cdot (c - a) + (a + c) \cdot (a + c)$$

$$c \cdot c - 2a \cdot c + a \cdot a + a \cdot a + 2a \cdot c + c \cdot c$$

$$2|a|^2 + 2|c|^2$$

$$|OA|^2 + |AB|^2 + |BC|^2 + |OC|^2 \text{ as required}$$

$$6. \quad (a) \quad \angle AED = 90^\circ$$

Triangle in a semi-circle is always right angled.

$$(b) \quad \angle ABE = \angle ADE = 60^\circ$$

Angles within the same segment are congruent.

$$(c) \quad \angle CAE = 80^\circ$$

Opposite angles in a cyclic quadrilateral are supplementary

$$(d) \quad \angle TCE = \angle CBE = 100^\circ$$

The alternate segment theorem.

$$7. \quad \overrightarrow{AB} = \frac{2}{3} \overrightarrow{AC}$$

$$\therefore b - a = \frac{2}{3}(c - a)$$

$$\begin{pmatrix} 1 \\ 3 \end{pmatrix} - \begin{pmatrix} x \\ -1 \end{pmatrix} = \frac{2}{3} \begin{pmatrix} 4 - x \\ y + 1 \end{pmatrix}$$

$$\therefore x = -5 \text{ and } y = 5$$

Calculator-assumed Solutions

8. (a) Assume all non-repeated numbers are selected from
both sets: 3, 8, 4, 6 = 4 digits ✓
Plus all remaining digits from one set: 1, 2, 5, 7 = 4 digits
Plus one more digit to make the first repetition
∴ 4 + 4 + 1 = 9 digits minimum ✓
- (b) Assume the largest numbers are chosen first: ✓
8 + 7 + 7 + 6 = 28 ✓
one more digit could include the number 5, making
the sum over 30.
∴ 4 digits max ✓ [54]
9. (a) (i) ${}^{22}C_8 = 319770$ ✓
(ii) ${}^{12}C_4 \times {}^{10}C_4 = 103950$ ✓✓
(iii) ${}^{22}C_8 - {}^{20}C_6 = 281010$ ✓✓
(iv) ${}^2C_2 \times {}^{20}C_6 + {}^2C_0 \times {}^{20}C_8 = 38760 + 125970 = 164730$ ✓✓
- (b) (i) $8! = 40320$ ✓
(ii) $3! \times 6! = 4320$ ✓✓
(iii) $8! - 2! \times 7! = 30240$ ✓✓ [12]
10. (a) II and III ✓✓
(b) ${}^{10}C_1 \times {}^{60}P_6 \times {}^{42}C_1$ OR ${}^{10}C_1 \times {}^{60}C_6 \times {}^{42}C_1 \times 6!$ ✓✓✓
(c) LHS ✓

$$\frac{n!}{r! \times (n-r)!} + \frac{n!}{(r+1)! \times (n-r-1)!}$$

$$\frac{n!}{r! \times (n-r-1)!} \times \left[\frac{1}{(n-r)} + \frac{1}{(r+1)} \right]$$

$$\frac{n!}{r! \times (n-r-1)!} \times \left[\frac{r+1+n-r}{(n-r)(r+1)} \right]$$

$$\frac{n! \times (n+1)}{r! \times (r+1) \times (n-r-1)! \times (n-r)}$$

$$\frac{(n+1)!}{(r+1)! \times (n-r)!}$$


$$\frac{(n+1)!}{(r+1)! \times [(n+1) - (r+1)]!}$$

$${}^{n+1}C_{r+1} = \text{RHS}$$

[10]

11. (a) (i) $p+q$ ✓
 (ii) $2p-q$ ✓
 (iii) $-(3p+2q)$ ✓
 (iv) $2q-p$ ✓

(b)



[6]

12. (a) $\overrightarrow{AC} = \begin{pmatrix} 2 \\ k \end{pmatrix} - \begin{pmatrix} -2 \\ 0 \end{pmatrix} = \begin{pmatrix} 4 \\ k \end{pmatrix} = 4i + k j$ ✓
 $\overrightarrow{BC} = \begin{pmatrix} 2 \\ k \end{pmatrix} - \begin{pmatrix} 1 \\ -3 \end{pmatrix} = \begin{pmatrix} 1 \\ k+3 \end{pmatrix} = i + (k+3)j$ ✓

(b) $\begin{vmatrix} 4 \\ k \end{vmatrix} = \begin{vmatrix} 1 \\ k+3 \end{vmatrix}$

$\therefore \sqrt{4^2 + k^2} = \sqrt{1^2 + (k+3)^2}$ ✓

$\therefore k=1$ ✓

(c) $D = \left(\frac{-2+1}{2}, \frac{0-3}{2} \right) = (-0.5, -1.5)$ ✓

(d) $\overrightarrow{DC} = \begin{pmatrix} 2 \\ 1 \end{pmatrix} - \begin{pmatrix} -0.5 \\ -1.5 \end{pmatrix} = \begin{pmatrix} 2.5 \\ 2.5 \end{pmatrix}$

$\overrightarrow{DB} = \begin{pmatrix} 1 \\ -3 \end{pmatrix} - \begin{pmatrix} -0.5 \\ -1.5 \end{pmatrix} = \begin{pmatrix} 1.5 \\ -1.5 \end{pmatrix}$ ✓

$\therefore \overrightarrow{DC} \cdot \overrightarrow{DB} = \begin{pmatrix} 2.5 \\ 2.5 \end{pmatrix} \cdot \begin{pmatrix} 1.5 \\ -1.5 \end{pmatrix} = 2.5 \times 1.5 - 2.5 \times 1.5 = 0$ ✓

$\therefore \overrightarrow{DC} \perp \overrightarrow{DB} \rightarrow \angle CDB \text{ is right angled}$

[7]

13. (a) (i) True. ✓
 If a number is divisible by 6, then it is also divisible by both 2 and 3. ✓
- (ii) False. ✓
 It must be divisible by both 2 and 3. ✓
- (iii) True. ✓
 The conjunction AND means that it is divisible by both 2 and 3, and therefore it is also divisible by 6. ✓
- (b) Assume that n is odd AND that $3n+5$ is also odd ✓
 $\therefore \exists k \in \mathbb{N}$ such that $n=2k+1$ ✓
 $\therefore 3n+5 = 3(2k+1)+5$ ✓
 $= 6k+8 = 2(3k+4) = \text{even}$ ✓
 Since $3n+5$ is both odd and even simultaneously, this is a contradiction, implying that n must be even. ✓
- (c) $\vec{OE} + \vec{AD} + \vec{BF}$
 $= \left[b + \frac{1}{2}(a-b) \right] + \left[-a + \frac{1}{2}b \right] + \left[-b + \frac{1}{2}a \right]$ ✓✓✓
 $= \left(\frac{1}{2}a - a + \frac{1}{2}a \right) + \left(b - \frac{1}{2}b - \frac{1}{2}b \right) = 0$ ✓ [15]
14. (a) $F_1 \cos 30^\circ = F_2 \cos 45^\circ$ ✓
 $\therefore \frac{\sqrt{3}}{2} F_1 = \frac{1}{\sqrt{2}} F_2$
 $F_1 \sin 30^\circ + F_2 \sin 45^\circ = 250$ ✓
 $\therefore \frac{1}{2} F_1 + \frac{1}{\sqrt{2}} F_2 = 250$
- (b) $F_2 = \frac{\sqrt{6}}{2} F_1$
 $\therefore \frac{1}{2} F_1 + \frac{\sqrt{3}}{2} F_1 = 250$ ✓✓
 $\therefore F_1 = \frac{500}{1+\sqrt{3}} = 183.01 \text{ N}$
 $\therefore F_2 = \frac{\sqrt{6}}{2} \times \frac{500}{1+\sqrt{3}} = \frac{250\sqrt{6}}{1+\sqrt{3}} = 224.14 \text{ N}$ ✓✓
- (c) If $F_1 = 200 \text{ N}$ then $F_2 = \frac{\sqrt{6}}{2} \times 200 = 244.95 \text{ N} > 200 \text{ N}$ ✓
 \therefore Cable 2 exceeds its maximum load, hence Cable 1 must not reach its 200N maximum rating ✓

$$\text{If } F_2 = 200 \text{ N then } F_1 = \frac{2}{\sqrt{6}} \times 200 = 163.30 \text{ N} < 200 \text{ N} \quad \checkmark \checkmark$$

$$\text{Max Force } \checkmark F_1 \sin 30^\circ + F_2 \sin 45^\circ \quad \checkmark$$

$$\checkmark \frac{400}{\sqrt{6}} \times \frac{1}{2} + 200 \times \frac{1}{\sqrt{2}} = 223.07 \text{ N} \quad \checkmark \quad [12]$$

15. (a) $n(D \cup C) = n(D) + n(C) - n(D \cap C)$ ✓
 $810 = 400 + 500 - n(D \cap C)$ ✓
 $\therefore n(D \cap C) = 90$ ✓
- (b) $n(D \cup C \cup B) = n(D) + n(C) + n(B)$
 $- n(D \cap C) - n(D \cap B) - n(C \cap B)$
 $+ n(D \cap C \cap B)$ ✓
 $900 = 400 + 500 + 210 - 90 - 60 - 110 + n(D \cap C \cap B)$ ✓
 $\therefore n(D \cap C \cap B) = 50$ ✓ [6]
16. (a) $\binom{3}{C_1}^2 = {}^3C_2 + {}^4C_2$ ✓
 $\binom{5}{C_1}^2 = {}^5C_2 + {}^6C_2$ ✓
- (b) $a = 2$ ✓
 $b = n + 1$ ✓
- (c) $\binom{6}{C_1}^2 = {}^8C_3 - {}^6C_3$ ✓✓
- (d) $\binom{n}{C_1}^2 = {}^{n+2}C_3 - {}^nC_3$ ✓✓ [8]
17. (a) $\vec{OE} = a + \frac{1}{2}(b - a) = \frac{1}{2}(a + b)$ ✓
 $\vec{OF} = c + \frac{1}{2}(b - c) = \frac{1}{2}(b + c)$ ✓
- (b) $\vec{DE} = \vec{OE} - \vec{OD} = \frac{1}{2}(a + b) - \frac{1}{2}a = \frac{1}{2}b$ ✓
 $\vec{GF} = \vec{OF} - \vec{OG} = \frac{1}{2}(b + c) - \frac{1}{2}c = \frac{1}{2}b = \vec{DE}$ ✓
 $\vec{DG} = \vec{OG} - \vec{OD} = \frac{1}{2}c - \frac{1}{2}a = \frac{1}{2}(c - a)$ ✓
 $\vec{EF} = \vec{OF} - \vec{OE} = \frac{1}{2}(b + c) - \frac{1}{2}(a + b) = \frac{1}{2}(c - a) = \vec{DG}$ ✓
 $\therefore \vec{DE} = \vec{GF} \wedge \vec{DG} = \vec{EF} \Rightarrow \text{DEFG is a parallelogram}$ ✓ [7]
18. $BF \times FD = CF \times FA$ ✓
 $\therefore x(2y) = 2x(6) \rightarrow y = 6 \text{ cm}$ ✓
- $MD \times MB = MT^2$ ✓
 $\therefore 4 \times (4 + 12 + x) = \sqrt{76}^2 \rightarrow x = 3 \text{ cm}$ ✓

$$NT^2 = NC \times (NC + FC + FA)$$

✓

$$\therefore z^2 = 6 \times (6 + 6 + 6) \rightarrow z = 6\sqrt{3} \text{ cm}$$

✓

[6]

19. Let \hat{n} be a unit vector perpendicular to b

$$\text{then, } a = u + |u| \tan 60^\circ \hat{n} \quad \checkmark$$

$$|u| = \left| \begin{pmatrix} 3 \\ 1.5 \end{pmatrix} \right| = \frac{3}{2} \sqrt{5} \quad \checkmark$$

$$n \cdot \begin{pmatrix} 4 \\ 2 \end{pmatrix} = 0 \rightarrow \text{let } n = \pm \begin{pmatrix} -1 \\ 2 \end{pmatrix} \quad \checkmark$$

$$\therefore \hat{n} = \pm \frac{1}{\sqrt{5}} \begin{pmatrix} -1 \\ 2 \end{pmatrix} \quad \checkmark$$

$$\therefore a = \begin{pmatrix} 3 \\ 1.5 \end{pmatrix} + \frac{3}{2} \sqrt{5} \times \sqrt{3} \times \pm \frac{1}{\sqrt{5}} \begin{pmatrix} -1 \\ 2 \end{pmatrix} = \begin{pmatrix} 3 \\ 1.5 \end{pmatrix} \pm \frac{3}{2} \sqrt{3} \begin{pmatrix} -1 \\ 2 \end{pmatrix}$$

$$\therefore x = 3 \pm \frac{3}{2} \sqrt{3} \quad \text{and} \quad y = \frac{3}{2} \mp 3 \sqrt{3} \quad \checkmark \checkmark \quad [6]$$