
Test 4 Thursday 28 June 2018

Total marks available: 49



sətunim 02



METHODS 1&2

Student Name:

Time allowed for this assessment

ALL Calculator-assumed

Working time for this section: Formulae Sheet allowed.

Classpad assumed.

.bewolle solon to (bebis elgnis) egeq 4A 1

This study source was downloaded by 100000846012161 from CourseHero.com on 09-12-2022 04:46:57 GMT -05:00

9. [4, 4 = 8 marks]

(a) If A and B are acute angles, with $\sin(A) = \frac{3}{5}$ and $\tan(B) = \frac{12}{5}$ show that $\frac{31}{50} = |A + A| \cos A$

(b) Solve the equation
$$\pi \le x \ge 0 \quad \text{for } |x| = \cos(x) + \cos(x) = \cos(x)$$

1. [1, 1, 1, 2 = 5 marks]


Given that $U=|x|40 \le x \le 60$ | where x is an integer, A=[41,43,55,58],

 $B = [52,54,55,56] \land C = [41,43,56,60]$


- a) Find $U \lor \dot{\iota}$
- b) Find $A \cup B$
- c) Determine $n(C \cap \dot{B})$
- d) Determine $P(A \cup B \cup C)$
- 2. [1, 1 = 2 marks]

In the Venn diagrams below, shade the region representing:

a) $A' \cap B'$



b) $(B \cup C) \cap A$



8. [1, 1, 2 = 4 marks]

A committee consisting of 10 senior members and 12 junior members has decided to select five of its members to form a subcommittee.

- (a) Determine the number of different subcommittee combinations that can be selected.
- (b) Determine the number of different subcommittees that can be selected that contain only senior members
- (c) Determine the probability that a randomly chosen subcommittee contains at least one junior member

7

; [Ţ, Ţ, 3 = 5 marks]

Evaluate each of the following, show your working for full marks to be awarded:

- (a) 6!
- * d 9 (q)

(c)
$$\binom{2}{9} \times 3i$$

4. [4 marks]

Expand and simplify completely using an appropriate method that you have learnt:

$$s(x-z)$$

2. [1, 2 = 3 marks]

Consider the expansion for: $(x+3y)^{11}$ in descending powers of x.

(a) How many terms are there in this expansion?

https://www.coursehero.com/file/97147793/Year-11-Methods-Test-4-FLP-Finaldocx/

(b) Find the $3^{\rm rd}$ term in this expansion using Combination Theory $\binom{n}{r}$

3 This study source was downloaded by 100000846012161 from CourseHero.com on 09-12-2022 04:46:46:46:70 GMT -05:00

Given P(A)=m+0.2 and P(B)=m+0.3 and $P(A\cap B)=m$, calculate the value of m if:

(a) A and B are mutually exclusive

 $9.0 = (\mathbf{A} \cup \mathbf{A})\mathbf{q} \tag{d}$


(c) A and B are independent events

 $8.0 = (\mathbf{A} \vee \mathbf{A}) \mathbf{q} \tag{b}$

5) 25-91-10 660


This study source was downloaded by 100000846012161 from CourseHero.com on 09-12-2022 04:46:57 GMT -05:00

https://www.coursehero.com/file/97143/Year-11-Methods-Test-4-FLP-Finaldocx/



An analysis of the 210 students in their final year of school determined that 35 chose to study Physics, 45 chose to study Chemistry and 151 chose neither of these subjects.

(a) Complete the Venn diagram below using the information given.



- (b) Determine the number of students who chose to study both Physics and Chemistry.
- (c) Determine the probability that a randomly selected student chose to study
 - (i) Chemistry.
 - (ii) Physics but did not choose Chemistry.
 - (iii) Chemistry given that they chose to study Physics.
- (d) Is there any indication that choosing to study Chemistry is independent of choosing to study Physics? Justify your answer.