



YEAR 12 MATHEMATICS

3C/D/MAT

Section Two
(Calculator-Assumed)

Student Name: _____

Circle your teacher's name

S. ROWDEN

N. EDMUNDS

Reading time before commencing work: 10 minutes
Working time for section: 100 minutes

To be provided by the supervisor
Questions/answer booklet for Section Two. Candidates may use the removable formula sheet from Section One.

Material required/recommended for this section

Standard items: pens, pencils, pencil sharpener, highlighter, eraser, ruler
Special items: drawing instruments, templates, notes on up to two unfolded sheets of A4 paper, and up to three calculators, CAS, graphic or scientific, which satisfy the conditions set by the Curriculum Council for this course.

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor before reading any further.

Important note to candidates

Structure of this examination


	Number of questions available	Number of questions to be attempted	Suggested working time (minutes)	Marks available
Section One: Calculator—free	10	10	50 minutes	40
Section Two: Calculator—assumed	15	15	100 minutes	80
Total marks				120

Question number: _____

Instructions to candidates

1. Answer the questions in the spaces provided.
2. Spare answer pages are provided at the end of this booklet. If you need to use them, indicate in the original answer space where the answer is continued i.e. give the page number.
3. **Show all your working clearly.**
Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks.
For any question or part question worth more than two marks, valid working or justification is required to receive full marks.
If you repeat an answer to any question, ensure that you cancel the answer you do not wish to have marked.
4. It is recommended that you **do not use pencil** except in diagrams.

Question 25 (5 marks)
Consider the diagram below.



Question 25 (5 marks)
Consider the diagram below.

This section has Sixteen (16) questions. Attempt all questions.

Question 11 (4 marks)

(a) $P(E)$

(b) $P(E|R)$

(c) $P(R|E)$

[2]

(a) Show that $A = \frac{2}{X^2} + 5x$

Let A denote the area of the above figure.

[3]

(b) Use differentiation to find the approximate percentage change in A , if the percentage change in x is 5%, given that $x = 4$.

[1]

Pennros College
SEE NEXT PAGE

Semester I 2011


END OF QUESTIONS

Question 12 (4 marks)

Three types of components, X, Y, and Z, must each be processed in three separate machines. The respective processing times in machine A are 9, 10 and 16 minutes. For machine B, the corresponding times are 12, 14 and 8 minutes, while, for machine C, the times are 2, 15 and 12. How many of each type of component should be manufactured in an 8-hour shift in order to keep all the machines fully occupied?

Question 24 (5 marks)

The diagram below shows the 'velocity/time' graph for a particle which moves in a horizontal straight line for $0 \leq t \leq 8$ seconds. At time $t = 0$ seconds the particle is at a point O on the line; the initial velocity is 20 ms^{-1} .



Find:

- (a) the distance of the particle from O when $t = 8$.

[3]

- (b) the maximum distance of the particle from O.

[1]

- (c) the acceleration of the particle when $t = 2$.

[1]

[SEE NEXT PAGE](#)[SEE NEXT PAGE](#)[SEE NEXT PAGE](#)

[3]

- (b) Find using calculus when most of the population in the suburb is infected and the maximum population affected.

[1]

(a) Describe the interpretation given to $\frac{dp}{dt}$.

Given that $p(t) = \frac{4 + t^2}{t}$

During an influenza epidemic, the proportion of the population in a particular suburb who are infected is denoted $p(t)$ where t is time in weeks after the start of the epidemic.

- A: if the events are mutually exclusive, they must be independent.
 B: if the events are independent, they must be mutually exclusive.
 C: if the events are not mutually exclusive, they must be independent.
 D: if the events are not independent, they must be mutually exclusive.
 E: if the events are mutually exclusive, they cannot be independent.

Question 13 (4 marks)

Justify your answer.

Which of the following statements is true for two events, each with probability greater than 0?

Question 14 (6 marks)

The Australian Olympic Three Day Event Team must comprise of 4 elite event riders and their horses. There are 14 possible contenders and horses for this team, which will represent Australia at the 2012 London Olympics.

- (a) How many different selections are possible?

[1]

Ella and her horse Simmo, are the Australian champions and Emily, with her horse Nobby is the runner up champion.

- (b) What is the probability that of the 4 event riders chosen at random:

- (i) Ella is included?

[1]

- (ii) Ella and Emily are included?

[1]

- (iii) Ella or Emily, but not both is selected?

[1]

- (c) If Ella is selected for the Olympic team, what is the probability that Emily is also selected?


[2]

Question 22 (4 marks)

A dressmaker wishes to cut a section of cloth from a piece of material measuring two metres by one metre. The curved edges of the piece of cloth to be removed are defined as being between the following equations:

$$y_1 = 0.5x \text{ and } y_2 = x^3 - 3.1x^2 + 2.7x$$

Determine, using calculus, the area of the cloth removed correct to 2 decimal places.




[SEE NEXT PAGE](#)[SEE NEXT PAGE](#)[SEE NEXT PAGE](#)

[3]

- (b) Determine the volume of chocolate needed to make the kisses if the special gift box contains 3 kisses.

[1]

- (a) Determine the value of a given $a > 3$.



Each unit on the axes represents 1 cm.

$$g(x) = 0.125x^2 - 2x + 8 \quad \text{for } a < x \leq 8$$

$$f(x) = -0.375x^2 + 1.25x + 3 \quad \text{for } 0 \leq x \leq a$$

The kiss is to be modeled on the following equations that are rotated about the x axis for the given domains.

Hershey's are famous for their chocolate kisses and have decided to design a new size Hershey kiss and package in a special gift box.

Question 21 (4 marks)



[2]

(a) Given $f(x) = \sqrt{x} + 2$, $g(x) = \frac{1}{x+5}$ and $k(x) = \frac{1}{x} - 5$ determine:

$$g \circ f(t)$$

Question 15 (4 marks)

Question 16 (7 marks)

Gas is escaping from a spherical balloon at the rate of $0.4 \text{ m}^3/\text{min}$.

- (a) What is the change in volume during the first 10 minutes?

[2]

- (b) How fast is the surface area shrinking when the radius is 4 m?

[5]

Question 20 Continued

- (b) How many jazz and ballet classes should the dance teacher aim to have pupils for so that he can maximise his profit per week? Show all working and state clearly your optimal solution and maximum value.

[3]

- (c) Consider the vertex of the feasible region that is your solution to part (b) and the two adjacent vertices (one on either side). For each of these three vertices calculate the profit per week and per hour and comment upon your values.

[3]

- (d) There are no dance classes in the school holidays and throughout the year some classes may need to be cancelled for other reasons. Consequently, the dance teacher realizes that if he is considering his average profit over the whole year, the more realistic income from jazz dance classes is \$100 per class. What is the lowest income per ballet class that the teacher can receive so that the solution from part (b) of this question is the only combination of classes that gives maximum profit per week?

[3]

SEE NEXT PAGE

SEE NEXT PAGE

[2]

- (iii) will choose a large burger given that they have chosen a regular drink?

[1]

- (ii) will choose a regular drink given that they have chosen a regular burger

[1]

- (i) will choose both a large burger and a large drink?
 (b) Determine the probability that a randomly selected customer:


[3]

- (a) Draw a probability tree showing the possible burger and drink choices.
 Records show that two out of every three customers buy a large burger and, of these customers, one quarter of them choose a regular drink. Also, after choosing a burger, seven out of every ten customers choose a large drink.

Question 17 (7 marks)

Question 20 (3 marks)

- (a) Determine two more constraints, draw them on the above diagram and shade the feasible region.



Let x denote the number of jazz dance classes and y the number of ballet classes.
 Three of the constraints on x and y have been drawn on the diagram below.
 The teacher must make a profit (income - costs) or he will not set up his business.
 He plans to work at least 1 ballet class per week but also decides to teach no more than 5 ballet classes and no more than 8 jazz dance classes per week.
 The teacher will involve 6 hours of tuition a week and produce an income of \$360.
 Each jazz dance class will involve 3 hours of tuition a week and produce an income of \$120. Each class of dance and ballet, is writing up a business plan for a dance studio he is intends to start. He plans to teach jazz

and ballet must make a profit (income - costs) or he will not set up his business.

Let x denote the number of jazz dance classes and y the number of ballet classes.

Question 18 (8 marks)

A toy train sits at the centre of a length of track. The displacement, s , of the train from the central position, O, after t seconds is given by

$$s = 0.8t^2 - 6.4t \text{ cm}$$

- (a) Determine the displacement of the train after 3 seconds.

[1]

- (b) What speed is the train travelling after 3 seconds?

[2]

- (c) What is the acceleration after 3 seconds?


[1]

- (d) At what time(s) does the train stop in its first 20 seconds of motion?

[1]

- (e) What distance does the train travel in the first 20 seconds of motion?
Show your working.

[3]

SEE NEXT PAGE**Question 19 (3 marks)**

In each part, list the points (A-E) on the graph of $f(x)$ that satisfy the given conditions.

(a) $f'(x) > 0$ and $f''(x) > 0$

(b) $f'(x) < 0$ and $f''(x) > 0$

(c) $f'(x) = 0$ and $f''(x) < 0$

(d) $f'(x) = 0$ and $f''(x) > 0$

(e) $f'(x) < 0$ and $f''(x) = 0$

SEE NEXT PAGE