

Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until Friday June 16th 2017.



MATHEMATICS SPECIALIST UNIT 3

Semester One

2017

SOLUTIONS

Calculator-free Solutions

$$\begin{aligned}
 1. \quad (a) \quad P(2i) &= 2(2i)^3 - 4(2i)^2 + 8(2i) - 16 && \checkmark \\
 &= 2(-8i) - 4(-4) + 16i - 16 && \checkmark \\
 &= -16i + 16 + 16i - 16 = 0 \quad \text{as required}
 \end{aligned}$$

$$\begin{aligned}
 (b) \quad z_2 &= \overline{z_1} = -2i && \checkmark \\
 \text{Using factorisation:} \\
 P(z) &= 2(z+2i)(z-2i)(z-k) && \checkmark \\
 &= 2(z^2+4)(z-k) = 2(z^3 - 2z^2 - 8z - 18) \\
 \therefore k &= 2 && \checkmark \\
 \therefore P(z) &= 2(z+2i)(z-2i)(z-2) && \checkmark
 \end{aligned}$$

OR using division of polynomials:

$$\begin{array}{r}
 \begin{array}{r} 2z \quad -4 \\ \hline z^2+4 \overline{) 2z^3 - 4z^2 + 8z - 16} \\ \underline{2z^3 \qquad \qquad + 8z} \\ 0 \quad -4z^2 \quad 0 \quad -16 \\ \underline{-4z^2 \qquad \qquad -16} \\ 0 \end{array} \\
 \therefore P(z) = (z^2+4)(2z-4) = 2(z+2i)(z-2i)(z-2)
 \end{array}$$

[6]

$$\begin{aligned}
 2. \quad (a) \quad f(z) &= z^5 - 32i = 0 \\
 z^5 &= 32i = 32 \operatorname{cis}\left(\frac{\pi}{2} + 2\pi k\right) \text{ with } k=0, \pm 1, \pm 2 && \checkmark \\
 \therefore z &= 2 \cos\left(\frac{\pi + 4\pi k}{10}\right) \text{ with } k=0, \pm 1, \pm 2 && \checkmark
 \end{aligned}$$

hence,

$$k=0 \rightarrow z_0 = 2 \cos\left(\frac{\pi}{10}\right)$$

$$k=1 \rightarrow z_1 = 2 \cos\left(\frac{\pi}{2}\right)$$


$$k=-1 \rightarrow z_2 = 2 \cos\left(\frac{-3\pi}{10}\right)$$

$$k=2 \rightarrow z_3 = 2 \cos\left(\frac{9\pi}{10}\right)$$

$$k = -2 \rightarrow z_0 = 2 \cos\left(\frac{-7\pi}{10}\right)$$

✓✓


2. (b)



✓ magnitude = 2

✓ $\frac{2\pi}{5}$ radians apart


3. (a)



✓ correct x position

✓ correct y position

(b) (i)




$$y = -1 - |x + 2|$$

$$\therefore a = -1 \text{ and } b = 2$$

✓✓

3. (b) (ii)



$$y = 3 - |x - 2|$$

$$\therefore a = 3 \text{ and } b = -2$$

✓✓

[2]

4. (a) $h(x) = f(f(x)) = \frac{1}{1+f(x)}$ with $x \neq -1$

$$\therefore 1 + f(x) \neq 0$$

✓

$$f(x) = \frac{1}{1+x} \neq -1$$

$$\therefore x \neq -2 \text{ and } x \neq -1$$

✓✓

$f(x) \neq 0$ on its natural domain, hence

✓

$f(0) = 1$ will not be generated by the second $f(x)$


$$\therefore \{y : y \neq 1 \wedge y \neq 0\}$$

✓

(b)
$$h(x) = \frac{1}{1 + \left(\frac{1}{1+x}\right)} = \frac{1}{\frac{1+x+1}{1+x}} = \frac{x+1}{x+2}$$

$$\therefore h(x) = \frac{x+2-1}{x+2} = \frac{x+2}{x+2} - \frac{1}{x+2} = \frac{-1}{x+2} + 1$$

✓



✓ asymptotes
x = -2 and y = 1

✓ discontinuity at
x = -1

✓ correct shape

4. (c) $x = 1 - \frac{1}{y+2}$

$$\therefore y = \frac{-1}{x+1} - 2 = \frac{1-2x}{x-1} \quad \checkmark$$

Domain $\{x : x \neq 1, x \neq 0\}$ \checkmark

Range $\{y : y \neq -2, y \neq -1\}$ \checkmark [12]

5. (a) Cartesian equations of the planes:

$$2x + y + z = 1$$

$$4y + 2z = 2$$

$$3x + y - z = 8$$

Using matrices:

$$\begin{bmatrix} 2 & 1 & 1 & 1 \\ 0 & 4 & 2 & 2 \\ 3 & 1 & -1 & 8 \end{bmatrix} \begin{matrix} R_1 \\ R_2 \\ R_3 \end{matrix}$$

$$\begin{bmatrix} 2 & 1 & 1 & 1 \\ 0 & 2 & 1 & 1 \\ 0 & -1 & -5 & 13 \end{bmatrix} \begin{matrix} R_1 \\ \frac{1}{2}R_2 \\ 2R_3 - 3R_1 \end{matrix} \quad \checkmark$$

$$\begin{bmatrix} 2 & 1 & 1 & 1 \\ 0 & 2 & 1 & 1 \\ 0 & 0 & -9 & 27 \end{bmatrix} \begin{matrix} R_1 \\ R_2 \\ R_2 + 2R_3 \end{matrix} \quad \checkmark$$

$$\therefore -9z = 27, \quad z = -3$$

$$2y - 3 = 1, \quad y = 2$$

$$2x + 2 - 3 = 1, \quad x = 1$$

Unique solution: $(1, 2, -3)$ \checkmark

(b) Fully simplified matrix in terms of a :

$$\begin{bmatrix} 2 & 1 & 1 & 1 \\ 0 & 2 & 1 & 1 \\ 0 & 0 & (4a-5) & 27 \end{bmatrix} \quad \checkmark$$

$$\therefore 4a - 5 = 0 \quad \checkmark$$

$$a \neq \frac{5}{4} \quad \checkmark \quad [6]$$

(Algebraic manipulation and substitution is also acceptable)

6. (a) Midpoint between $\begin{pmatrix} 0 \\ -4 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 4 \\ 0 \\ -4 \end{pmatrix}$ is $\begin{pmatrix} 2 \\ -2 \\ -2 \end{pmatrix}$ ✓✓

OR $\vec{AG} = \begin{pmatrix} 4 \\ 4 \\ -4 \end{pmatrix}$, centre = $\vec{OA} + \frac{1}{2}\vec{AG} = \begin{pmatrix} 0 \\ -4 \\ 0 \end{pmatrix} + \frac{1}{2}\begin{pmatrix} 4 \\ 4 \\ -4 \end{pmatrix} = \begin{pmatrix} 2 \\ -2 \\ -2 \end{pmatrix}$ ✓✓

radius = $\frac{1}{2}|\vec{AG}| = \frac{1}{2}\sqrt{4^2 + 4^2 + (-4)^2} = 2\sqrt{3}$ units ✓

$\left| r - \begin{pmatrix} 2 \\ -2 \\ -2 \end{pmatrix} \right| = 2\sqrt{3}$ ✓

(b) $\vec{AF} = \begin{pmatrix} 4 \\ 0 \\ -4 \end{pmatrix}$, $\vec{AB} = \begin{pmatrix} 4 \\ 4 \\ 0 \end{pmatrix}$

$\vec{AF} \times \vec{AB} = \begin{pmatrix} 0+16 \\ -16-0 \\ 16-0 \end{pmatrix} = \begin{pmatrix} 16 \\ -16 \\ 16 \end{pmatrix} = 16\begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$ ✓✓

$k = \begin{pmatrix} 4 \\ 0 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} = 4$ ✓

$\therefore r \cdot \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} = 4$ ✓

[8]

7. (a) $x = 2 \cos t \rightarrow \cos t = \frac{x}{2}$


$\sin^2 t + \cos^2 t = y^2 + \frac{x^2}{4} = 1$ ✓✓

(b) $x = 1 \rightarrow 2 \cos t = 1, \therefore t = \frac{\pi}{3}$ ✓

$\dot{r}(t) = -2 \sin t i + \cos t j$ ✓

$\dot{r} = -2 \sin\left(\frac{\pi}{3}\right)i + \cos\left(\frac{\pi}{3}\right)j$

$\dot{r} = -\sqrt{3}i + \frac{1}{2}j$ ✓



[6]

Calculator-Assumed Solutions

8. (a) $x=2$ and $y=\frac{-x}{2}+1$ ✓✓

(b) $f(x)=\frac{k}{x-2}+\left(\frac{-x}{2}+1\right)$

$f(3)=\frac{k}{1}+1-\frac{3}{2}=\frac{3}{2} \rightarrow k=2$ ✓

$\therefore f(x)=\frac{2}{x-2}-\frac{x}{2}+1=\frac{-x^2+4x}{2x-4}$ ✓

$\therefore a=-1, b=4, c=0, m=2, n=-4$ ✓✓

[6]

9. (a) $v(t)=\int [(6t)i+2j] dt=(3t^2+A)i+(2t+B)j$ ✓

$v(0)=\begin{pmatrix} 0+A \\ 0+B \end{pmatrix}=\begin{pmatrix} -2 \\ -3 \end{pmatrix}$

$\therefore v(t)=(3t^2-2)i+(2t-3)j$ ✓

$r(t)=\int [(3t^2-2)i+(2t-3)j] dt$

$i(t^3-2t+C)i+(t^2-3t+D)j$ ✓

$r(0)=\begin{pmatrix} 0+C \\ 0+D \end{pmatrix}=\begin{pmatrix} 0 \\ 0 \end{pmatrix}$

$\therefore r(t)=(t^3-2t)i+(t^2-3t)j$ ✓

(b) Condition to stop:

$|v(t)|=\begin{vmatrix} 3t^2-2 \\ 2t-3 \end{vmatrix}=0$ ✓

$\therefore t=\pm\sqrt{\frac{2}{3}}$ for x axis and $t=\frac{3}{2}$ for y axis ✓

Since there is no unique solution, the particle does not come to a stop. ✓


(c) Motion parallel to x axis = no y axis velocity

$\therefore 2t-3=0 \rightarrow t=\frac{3}{2}$ seconds ✓✓

$\therefore \left|v\left(\frac{3}{2}\right)\right|=\begin{vmatrix} 3\left(\frac{3}{2}\right)^2-2 \\ 0 \end{vmatrix}=\frac{19}{4}=4.75 \text{ ms}^{-1}$ ✓

$$r\left(\frac{3}{2}\right) = \left(\left(\frac{3}{2} \right)^3 - 2 \left(\frac{3}{2} \right) \right) = \left(\frac{3}{8} \right) = \frac{3}{8}i - \frac{9}{4}j \quad \checkmark \quad [11]$$

10. (a)




✓ Vertical asymptotes
 $x = -3, x = 1$

✓ Indicates $y \rightarrow 0^-$
as $|x| \rightarrow \infty$

✓ Shows local min
at $x = -1$

✓ Correct curvature at
 $x = -3$ and $x = 1$

(b)



✓ Correct mirror image
over x axis

✓ Correct mirror image
over y axis

✓ Indicates points
 $(0,3), (-1,0)$ and $(1,0)$

(c) Function can be inverted about its turning point.

i.e., for $x = -1 \rightarrow \therefore k = -1$

✓

Domain $\{x : x \leq 4\}$

✓

Range $\{y : y \geq -1\}$

✓

[10]

11. (a) At the xz plane $y = 0$.

✓

$$\therefore x^2 + z^2 - 2x = 7$$

$$(x-1)^2 + z^2 = 8$$

✓

circle centred at $(1, 0)$ and radius $2\sqrt{2}$ units

✓✓

- (b) Completing the squares of the original sphere:

$$(x-1)^2 + (y+1)^2 + z^2 = 9$$

$$\therefore \left| r - \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} \right| = 3$$

✓

centre of sphere is $C(1, -1, 0)$

$$\therefore n = \overrightarrow{CP} = \begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix} - \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix}$$

✓

$$k = n \cdot \overrightarrow{OP} = \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix} = 6 + 2 + 1 = 9$$

✓

$$\therefore r \cdot \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix} = 9 \text{ and hence } 2x + 2y + z = 9$$

✓

[8]

12. (a) $F(x) = [\cos(1-x)]^2 = f[\cos(1-x)] = f[g(1-x)] = f[g(h(x))]$

$$\therefore f(x) = x^2, g(x) = \cos x, h(x) = 1-x$$

✓✓✓


- (b) (i) $f(g(1)) = f(2) = 3$

✓

$$f(f(-3)) = f(0) = -3$$

✓

- (ii)




- ✓ correct sections
- ✓ correct y-intersect
- ✓ correct discontinuities

[8]

13. (a) $|z+2| \leq |z-2+2i|$

✓✓✓

(b) $r \leq \theta \therefore$ region under spiral



✓ spiral $r = \theta$

✓ region under spiral
and x axis

(b) $\frac{z+1}{z+i} = \frac{(x+1)+yi}{x+i(y+1)}$ such that $z \neq -i$

CAS: $i \left(\frac{x^2+y^2+x+y}{x^2+(y+1)^2} \right) + i \left(\frac{-x-y-1}{x^2(y+1)^2} \right)$

✓

$\text{Arg}\left(\frac{z+1}{z+i}\right) = \frac{\pi}{4} \rightarrow \Re\left(\frac{z+1}{z+i}\right) = \Im\left(\frac{z+1}{z+i}\right)$

$\therefore x^2+y^2+x+y = -x-y-1$

✓

$(x+1)^2 + (y+1)^2 = 1$


✓

circle centred at $(-1, -1)$ and radius $r=1$

✓

with discontinuity at $z = -i$

✓



[10]

14. (a) height = vertical velocity \times time = $90 \times 4 = 360$ m ✓

(b) Detonation at $\vec{OA} = \begin{pmatrix} 80 \\ -120 \\ 0 \end{pmatrix} + 4 \begin{pmatrix} 30 \\ -10 \\ 90 \end{pmatrix} = \begin{pmatrix} 200 \\ -160 \\ 360 \end{pmatrix}$ ✓

$$\vec{AP} = \vec{OP} - \vec{OA} = \begin{pmatrix} -200 \\ 350 \\ 20 \end{pmatrix} - \begin{pmatrix} 200 \\ -160 \\ 360 \end{pmatrix} = \begin{pmatrix} -400 \\ 510 \\ -340 \end{pmatrix}$$
 ✓

direct distance = $|\vec{AP}| = \sqrt{(-400)^2 + 510^2 + (-340)^2} = 731.92$ m ✓

time difference $\frac{731.92}{340} = 2.15$ seconds ✓

(c) speed $\sqrt{30^2 + (-10)^2 + 90^2} = 10\sqrt{91} = 95.39$ ms⁻¹ ✓

projection vector onto xy plane $\begin{pmatrix} 30 \\ -10 \\ 0 \end{pmatrix}$ ✓

$$\therefore \begin{pmatrix} 30 \\ -10 \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 30 \\ -10 \\ 90 \end{pmatrix} = \left| \begin{pmatrix} 30 \\ -10 \\ 0 \end{pmatrix} \right| \times \left| \begin{pmatrix} 30 \\ -10 \\ 90 \end{pmatrix} \right| \times \cos \theta$$
 ✓


$$\therefore \cos \theta = \frac{10}{\sqrt{910}} \rightarrow \theta = 70.64^\circ$$
 ✓

(d) distance vector $\begin{pmatrix} 200 \\ -160 \\ 360 \end{pmatrix} - \begin{pmatrix} 120 \\ -40 \\ 0 \end{pmatrix} = \begin{pmatrix} 80 \\ -120 \\ 360 \end{pmatrix}$ ✓

velocity $\frac{1}{4} \begin{pmatrix} 80 \\ -120 \\ 360 \end{pmatrix} = \begin{pmatrix} 20 \\ -30 \\ 90 \end{pmatrix}$ ms⁻¹ ✓

[11]

15. (a) (i)



15. (a) (ii) $|z+w|^2 = |z-w|^2$

$$(z+w)^2 = (z-w)^2 \quad \checkmark$$

$$z^2 + 2zw + w^2 = z^2 - 2zw + w^2 \quad \checkmark$$

$$4zw = 0 \text{ and hence } wz = 0 \quad \checkmark$$

if z, w are treated as vectors, then $z \perp w$ \checkmark

and hence $|\arg(z) - \arg(w)| = \frac{\pi}{2}$

(b) Let $u = x + yi$

$$|i+x+yi|^2 + |i-x-yi|^2$$


$$\checkmark (\sqrt{x^2+(y+1)^2})^2 + (\sqrt{(-x)^2+(1-y)^2})^2 \quad \checkmark$$

$$\checkmark x^2 + y^2 + 2y + 1 + x^2 + 1 - 2y + y^2 \quad \checkmark$$

$$\checkmark 2 + 2(x^2 + y^2) = 2 + 2(1) = 4 \quad \checkmark \checkmark$$

(c) (i) $iz^2 = \text{cis}\left(\frac{\pi}{2}\right) r^2 \text{cis}(2\theta) = r^2 \text{cis}\left(2\theta + \frac{\pi}{2}\right)$

$$\therefore |iz^2| = r^2 \text{ and } \arg(iz^2) = 2\theta + \frac{\pi}{2} \quad \checkmark \checkmark$$



$$\therefore |-2y-2xi| = |-2(y+xi)| = 2r \quad \checkmark$$

$$\arg(-2y-2xi) = -\left(\theta + \frac{\pi}{2}\right) \quad \checkmark$$

[14]

16. (a) CAS $\rightarrow Q(z) = z^2 + z - 1$ ✓✓
 (OR, using division of polynomials)
- (b) $(z^3 + 1)(z^2 + z - 1) = 0$
 $z^3 = -1$
 $\therefore z_1 = -1, z_2 = \frac{1}{2} + \frac{i\sqrt{3}}{2}, z_3 = \frac{1}{2} - \frac{i\sqrt{3}}{2}$ ✓✓✓
 $z = \frac{-1 \pm \sqrt{1+4}}{2}$
 $\therefore z_4 = \frac{-1 + \sqrt{5}}{2}, z_5 = \frac{-1 - \sqrt{5}}{2}$ ✓✓ [7]
17. (a) By De Moivre's theorem
 $(\cos \theta + i \sin \theta)^4 = \cos(4\theta) + i \sin(4\theta)$ ✓
 $i \cos^4 \theta + 4i \sin \theta \cos^3 \theta - 6 \sin^2 \theta \cos^2 \theta - 4i \cos \theta \sin^3 \theta + \sin^4 \theta$ ✓
 $\therefore \cos(4\theta) = \Re[(\cos \theta + i \sin \theta)^4]$
 $i \cos^4 \theta - 6 \sin^2 \theta \cos^2 \theta + \sin^4 \theta$ ✓
 $i \cos^4 \theta - 6 \cos^2 \theta (1 - \cos^2 \theta) + (1 - \cos^2 \theta)^2$ ✓
 $\therefore \cos(4\theta) = 8 \cos^4 \theta - 8 \cos^2 \theta + 1$ ✓
- (b) Let $x = \cos \theta$, then ✓
 $8x^4 - 8x^2 + 1 = 0$ when $\cos 4\theta = 0$ ✓
 $\therefore 4\theta = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \frac{7\pi}{2}$
 and $\therefore \theta = \frac{\pi}{8}, \frac{3\pi}{8}, \frac{5\pi}{8}, \frac{7\pi}{8}$
 $\therefore x = \cos \frac{\pi}{8}, \cos \frac{3\pi}{8}, \cos \frac{5\pi}{8}, \cos \frac{7\pi}{8}$ ✓ [8]
18. (a) If the dot product is zero, then a is perpendicular to the plane normal to b and c . ✓✓
 If the cross product is zero, then b and c are parallel ✓✓
- (b) LHS $i(a+b) \times (a-b)$
 $i a \times a - a \times b + b \times a - b \times b$ ✓
 $i 0 - (-(b \times a)) + b \times a - 0$ ✓
 $i b \times a + b \times a$ ✓

$$2(b \times a) = \text{RHS}$$

[7]