


Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until Wednesday 10th October 2018.



MATHEMATICS METHODS UNITS 3 & 4

Semester Two

2018

SOLUTIONS

Calculator-free Solutions

1. (a) $\log_x 9 + \log_x x^2$ ✓
 $= \frac{\log_3 9}{\log_3 x} + 2\log_x x$ ✓
 $= \frac{2}{p} + 2 \text{ or } \frac{2+2p}{p}$ ✓
- (b) $3^{2\log_3 3} = 3^2$ ✓
 $= 9$ ✓
- (c) $3^{2x^2}(2x) = 2x(3^{2x^2})$ ✓✓ [7]
2. (a) $\frac{dV}{dt} = 4\pi(12t - t^2)^2(12 - 2t)$ ✓
 $\frac{dV}{dt} = 0$ when $t(12 - t) = 0$ or $(12 - 2t) = 0$ ✓
 $\therefore t = 0$ or 6 or 12 ✓
Using the sign table yields Maximum occurs at $t = 6$ ✓
- (b) (i) $V = \frac{4}{3}\pi r^3 = \frac{4}{3}\pi(12t - t^2)^3$ ✓
 $\therefore r = 12t - t^2$ ✓
- (ii) When $t = 6$ then $r = 36$ ✓
 $r = 36$ is the maximum value of r , so max Volume. ✓ [8]
3. (a) $p = 0.5[(2) + (1.5) + (1.2)] = 0.5(4.7)$ ✓
 $= 2.35$ ✓
 $q = 0.5[(3) + (2) + (1.5)] = 0.5(6.5)$ ✓
 $= 3.25$ ✓
- (b) $\int_1^{2.5} \frac{3}{x} dx = [3 \ln x]_1^{2.5}$ ✓
 $= 3 \ln 2.5 - 3 \ln 1 = 3 \ln 2.5$ ✓ [6]
4. (a) 0.07 ✓
(b) 0.25 ✓
(c) $P(1) = (0.1)^2 = 0.01$ ✓
 $P(2) = (0.15)^2 = 0.0225$ ✓
 $\therefore P(1) + P(2) = 0.0325$ ✓
(d) Unbiased would be uniform ✓ [5]

5. (a) $\cos 5x - 5x \sin 5x$ ✓✓

(b) $\int \left(\frac{d}{dx}(x \cos 5x) \right) dx = \int (\cos 5x - 5x \sin 5x) dx$ ✓

$\therefore \int (\cos 5x) dx - \int (5x \sin 5x) dx = x \cos 5x + c$ ✓

$\therefore \int (5x \sin 5x) dx = \int (\cos 5x) dx - x \cos 5x + c$ ✓

$\therefore \int (x \sin 5x) dx = \frac{1}{25} \sin 5x - \frac{1}{5} x \cos 5x + c$ ✓ [6]

6. $4e^{2x} - 9e^x - 9 = 0$

Let $y = e^x$

$\therefore 4y^2 - 9y - 9 = 0$ ✓


$\therefore (4y + 3)(y - 3) = 0$ ✓

$\therefore y = -\frac{3}{4} \text{ or } y = 3$ ✓

$\therefore e^x = -\frac{3}{4} \rightarrow \text{no solution}$ ✓

and $e^x = 3 \rightarrow x = \ln 3$ ✓ [5]

7. (a)



$A = \int_{\frac{\pi}{2}}^{\pi} (\sin x - \cos x) dx$ ✓✓

(b) ✓

$= \left[-\cos x - \sin x \right]_{\frac{\pi}{2}}^{\pi}$ ✓

$= (0 + 1) - (-1 - 0) = 2$ ✓ [5]

8. $E_1 = 1.645 \sqrt{\frac{\hat{p}(1-\hat{p})}{n_1}}$

$E_2 = 1.645 \sqrt{\frac{\hat{p}(1-\hat{p})}{9n_1}}$ ✓

$\therefore \frac{E_2}{E_1} = \frac{\sqrt{n_1}}{\sqrt{9n_1}}$ ✓

$$= \frac{1}{3}$$

✓ [3]

9. (a) $\ln |\sin x| + c$ ✓✓

(b) $\left[\ln |\sin x| \right]_{\frac{\pi}{4}}^{\frac{\pi}{2}}$ ✓

$$= \ln(1) - \ln\left(\sin \frac{\pi}{4}\right) = 0 - \ln\left(\frac{1}{\sqrt{2}}\right) = \ln \sqrt{2}$$

✓

$$= \frac{1}{2} \ln 2$$

✓ [5]

Calculator-assumed Solutions

10. (a) $p + q = 0.4$ ✓

$$0.2 + 4p + 2.7 + 16q + 2.5 = 8.2$$

✓

$$\therefore \text{From CAS } p = 0.3 \text{ and } q = 0.1$$

✓

(b) $E(X) = 0.2 + 0.6 + 0.9 + 0.4 + 0.5 = 2.6$ ✓

(c) $E(Y) = 1 - 5.2 = -4.2$ ✓

[5]

11. (a) $\hat{p} = \frac{86}{200} = 0.43$ ✓

(b) $CI = 0.43 \pm 1.645 \sqrt{\frac{(0.43)(0.57)}{200}}$ ✓✓

$$= 0.43 \pm 0.0576$$

✓

$$\therefore 0.372 \leq p \leq 0.488$$

✓

(c) The mean suggested by the CI is $\frac{0.62 + 0.82}{2} = 0.72$ ✓

This does not lie within the 90% CI as calculated in (b) ✓

\therefore Evidence doesn't support the claim. ✓

(d) (i) Binomial requires independent events and a constant probability of success. Neither criteria apply here. ✓


(ii) $P(X = 1) = \frac{{}^7C_1 \times {}^3C_3}{{}^{10}C_4}$ ✓✓

$$= 0.1$$

✓

[12]

12. (a)



$$f(x) = \begin{cases} \frac{1}{4000} & 6000 < x < 10000 \\ 0 & \text{otherwise} \end{cases}$$

(b) (i) $P(8000 \leq X \leq 9000) = 0.25$ ✓✓

(ii) $P(X = 8500) = 0$ ✓

$$E(X) = \int_{6000}^{10000} \left(x \times \frac{1}{4000} \right) dx = 8000$$

(c) ✓✓

$$\text{VAR}(X) = \int_{6000}^{10000} \left((x - 8000)^2 \times \frac{1}{4000} \right) dx = 1\,333\,333.33$$

(d) (i) $\therefore \text{st dev} = 1154.7 = \1155 ✓✓

(ii) Standard deviation = \$1155 ✓

Not affected by change of origin ✓

[10]

13. (a) (i) $\ln V = \frac{7.22 - 6.91}{4} t + 6.91$ ✓

$\therefore \ln V = 0.0775t + 6.91$ ✓

(ii) $V = e^{0.0775t + 6.91}$ ✓

$\therefore V = 1002.25e^{0.0775t}$ ✓

(b) $2000 = 1002.25e^{0.0775t}$ ✓

$\therefore t = 8.91 \text{ years}$ ✓

(c) $V = V_0 e^{kt}$

$\therefore 2V_0 = V_0 e^{\frac{rt}{100}}$

$\therefore e^{\frac{rt}{100}} = 2$ ✓


$\therefore \frac{rt}{100} = \ln 2 \rightarrow t = \frac{100 \ln 2}{r}$ ✓

$\therefore t = \frac{69}{r}$ ✓

[9]

14. (a) $f'(x) = 2x \ln x + x^2 \left(\frac{1}{x} \right) = 2x \ln x + x$ ✓
 $\therefore 2x \ln x + x = 0$ when $x(2 \ln x + 1) = 0$
 $x = 0$ or $\ln x = -\frac{1}{2}$ ✓
 $\therefore x \neq 0$ or $x = e^{-\frac{1}{2}} = \frac{1}{\sqrt{e}} = 0.61$ ✓
 ie $f''(x) = 3 + 2 \ln x$ ✓
 $\therefore f''(0.61) > 0 \therefore \text{Min at } (0.61, -0.18)$ ✓
 (b) $f''(x) = 3 + 2 \ln x$
 $3 + \ln x = 0$ when $x = e^{-\frac{3}{2}}$ ✓
 $\therefore \text{Oblique POI at } (0.22, -0.08)$ ✓✓

(c)



✓✓✓✓ [12]

15. (a) (i) $v(t) = \int (3t + 5) dt = \frac{3t^2}{2} + 5t + 20$ ✓✓
 $\therefore v(3) = 48.5 \text{ m/sec}$ ✓
 (ii) $x(t) = \int \left(\frac{3t^2}{2} + 5t + 20 \right) dt = \frac{t^3}{2} + \frac{5t^2}{2} + 20t - 10$ ✓✓
 $\therefore x(3) = 86 \text{ m}$ ✓
 (b) $v = \frac{3t^2}{2} + 5t + 20 = 0$ when stopped ✓
 However $\frac{3t^2}{2} + 5t + 20 \neq 0$ so never stops ✓
 (c) Distance travelled = $x(3) - x(0)$ ✓
 $\therefore 86 - (-10) = 96 \text{ m}$ ✓

[10]

16. $X \sim \text{Bin}(36, 0.8)$

(a) $P(X \leq 30) = 0.7536$ ✓✓

(b) (i) $E(\hat{p}) = 0.8$ ✓

(ii) $\text{VAR}(\hat{p}) = \sqrt{\frac{(0.8)(0.2)}{36}} = \frac{1}{15}$ or 0.067 ✓✓

(c) $(0.9)^{10} = 0.349$ ✓✓

[7]

17. (a) Take proportions as follows:

$$20 - 29 : \frac{727}{3100} \times 100 = 23$$

$$30 - 39 : \frac{1050}{3100} \times 100 = 34$$

$$40 - 49 : \frac{800}{3100} \times 100 = 26$$

$$50 - 59 : \frac{523}{3100} \times 100 = 17$$

(b) (i) Normal distribution ✓✓✓

(ii) $\mu = 0.3387$ ✓

$$\text{Standard deviation} = \sqrt{\frac{(0.3387)(0.6613)}{100}} = 0.0473$$
 ✓

(c) $Y \sim N(0.3387, 0.0473^2)$ ✓

$\therefore P(X \geq 40) = P(Y \geq 0.4) = 0.0975$ ✓✓

[9]

18. (a) $P(7) = \frac{1}{6}$ ✓

(b) $P(<7) = \frac{15}{36}$ ✓

(c) $P(>7) = \frac{15}{36}$ ✓

(d) \$12 ✓

(e)

Game	7s	Unders	Overs
$P(X)$	$\frac{6}{36}$	$\frac{15}{36}$	$\frac{15}{36}$
Return	20	1	1

$$\therefore E(X) = \frac{6}{36} \times 20 + \frac{15}{36} \times 1 + \frac{15}{36} \times 1 = \frac{150}{36} = 4.16$$
 ✓✓

\therefore Club makes \$5 - \$4.16 = 84 cents per roll ✓

[7]

19. (a) $SA = (3x)(x) + (3x)(h)(2) + (x)(h)(2)$
 $= 3x^2 + 8xh$ ✓
 $V = 3x^2h = 18 \rightarrow h = \frac{6}{x^2}$ ✓
 $SA = 3x^2 + 8x\left(\frac{6}{x^2}\right) = 3x^2 + \frac{48}{x}$ ✓
 $\frac{d(SA)}{dx} = 6x - \frac{48}{x^2}$ ✓
 (b) $6x - \frac{48}{x^2} = 0 \rightarrow x = 2$ ✓
 \therefore Max occurs when $h = 1.5$ and $SA = 36 \text{ m}^2$ ✓
 $\frac{d(SA)}{dx} \approx \frac{\delta SA}{\delta x}$ ✓
 (c) $\delta SA = \left(\frac{d(SA)}{dx}\right)(\delta x) = \left(6x - \frac{48}{x^2}\right)(0.1)$ ✓
 When $x = 2$ then $\delta SA = 0 \times 0.1 = 0$ ✓
 (d) $\frac{dSA}{dt} = \frac{dSA}{dx} \times \frac{dx}{dh} \times \frac{dh}{dt}$
 $= \left(6x - \frac{48}{x^2}\right)\left(-\frac{1}{12x^{-3}}\right)(0.1)$ ✓✓
 When $x = 1$, $\frac{dSA}{dt} = 0.35 \text{ cm}^3/\text{sec}$ ✓ [11]
20. (a) $X \sim N(1.6, 0.4^2)$
 $P(X > 2) = 0.1587$ ✓✓
 (b) $P(X < k) = 0.9 \rightarrow k = 2.11$ ✓✓
 (c) (i) $X \sim \text{Bin}(4, 0.1587)$ ✓
 $\therefore P(X = 1) = 0.378$ ✓
 (ii) $0.1587 \times 0.5^3 \times ({}^4\text{C}_1)$ ✓
 $= 0.0794$ ✓ [8]