

Note: All part questions worth more than 2 marks require working to obtain full marks

Formula sheet provided: Yes

Task weighting: 10%

Marks available: 45 marks

Examinations

A4 paper, and up to three calculators approved for use in the WACE  
Drawing instruments, templates, notes on one unfolded sheet of

Standard items:  
Pens (blue/black preferred), pencils (including coloured), sharpener,  
correction fluid/tape, eraser, ruler, highlighters

Materials required:  
Calculator with CAS capability (to be provided by the student)

Number of questions: 6

Time allowed for this task: 40 mins

Task type:  
Response

Student name: \_\_\_\_\_ Teacher name: \_\_\_\_\_

## Course Methods test 4 Year 12

Perth Modern School  
Independent Public School  
Exceptional Schooling. Exceptional students.



## Q1 (3 marks)

Mary who goes to Perth Modern scored 67% in her exam which had a mean of 65% and a standard deviation of 18%. John who goes to Rossmoyne scored 62% in an exam with a mean of 58% and a standard deviation of 12%. Using standard scores, explain who had the better result.

**Solution**

**Edit Action Interactive**

The calculator screen shows two rows of calculations:

$\frac{67-65}{18}$  results in  $0.1111111111$

$\frac{62-58}{12}$  results in  $0.3333333333$

Below the calculator, the text "John has the higher z score hence better result." is displayed.

**Specific behaviours**

P shows calculation for z scores  
P states both z scores  
P states that John had the better result

## Q2 (3 marks)

Given that  $X \sim N(62, 196)$  and that  $P(k \leq x \leq 73) = 0.41$ , determine the value of  $k$  to two decimal places.

Perth Modern

Mathematics Department

(2 marks)

- a) Determine the proportion of calls that are between 11 and 17 mins.
- Calls that are longer than 17 mins are charged at double rate and are known as SUPER calls.
- The lengths of telephone calls,  $X$  in minutes, in Perth are Normally distributed with  $X \sim N(15, 3^2)$

Q3 (10 marks)

(Note - max 2 marks if working not shown)

P determines value of k to 2 dp

P subtracts to find area below k



P determines prob less than 73

**Specific behaviours**

k = 57.50

InvNormCDF(1, 0.37398, 14, 62) = 57.50

Accept below as working but not recommended in WACE!!!



Solution	
$\text{normCDF}(-\infty, 73, 196, 62)$ $0.783982554$ $0.783982554$ $\text{invNormCDF}("L", 0.37398, 14)$ $57.50137401$	
<input style="width: 40px; height: 25px; margin-right: 10px;" type="button" value="Edit"/> <input style="width: 40px; height: 25px; margin-right: 10px;" type="button" value="Action"/> <input style="width: 40px; height: 25px; margin-right: 10px;" type="button" value="Interactive"/>	
<input style="width: 25px; height: 25px; margin-right: 10px;" type="button" value="0.5"/> <input style="width: 25px; height: 25px; margin-right: 10px;" type="button" value="1"/> <input style="width: 25px; height: 25px; margin-right: 10px;" type="button" value="2"/>	
<input style="width: 40px; height: 25px; margin-right: 10px;" type="button" value="d&lt;sup&gt;-1&lt;/sup&gt;y/dx"/> <input style="width: 40px; height: 25px; margin-right: 10px;" type="button" value="Jdx"/> <input style="width: 40px; height: 25px; margin-right: 10px;" type="button" value="Simp"/>	
<input style="width: 40px; height: 25px; margin-right: 10px;" type="button" value="d&lt;sup&gt;2&lt;/sup&gt;y/dx&lt;sup&gt;2&lt;/sup&gt;"/> <input style="width: 40px; height: 25px; margin-right: 10px;" type="button" value="Jdx&lt;sup&gt;2&lt;/sup&gt;"/>	

**Solution**

**Edit Action Interactive**

$\frac{0.5}{2}$   $\frac{1}{2}$   $\frac{\partial}{\partial x}$   $\frac{\partial}{\partial x}$   $\int dx$   $\int dx$   $\text{Simp}$   $\frac{\partial}{\partial x}$   $\nabla$   $\nabla$   $\nabla$   $\nabla$

$\text{normCDF}(11, 17, 3, 15)$

0.6562962427

**Specific behaviours**

P uses correct parameters  
P states prob

**Extra working space**

- b) Given that a call is less than 13 mins, determine the probability that the call is greater than 9 mins. (3 marks)

**Solution**

$\frac{P(9 \leq X \leq 13)}{P(X \leq 13)}$

**Edit Action Interactive**

$\frac{0.5}{2}$   $\frac{1}{2}$   $\frac{\partial}{\partial x}$   $\frac{\partial}{\partial x}$   $\int dx$   $\int dx$   $\text{Simp}$   $\frac{\partial}{\partial x}$   $\nabla$   $\nabla$   $\nabla$   $\nabla$

$\text{normCDF}(9, 13, 3, 15)$


$\text{normCDF}(-\infty, 13, 3, 15)$

0.9098978046


**Specific behaviours**

P uses conditional prob formula  
P shows calculation for numerator and denominator  
P states prob

Q4 (15 marks)



- d) In Sydney the calls are also Normally distributed with a mean of 19 mins. If 25% of all calls are greater than 24 mins, determine the standard deviation to two decimal places. (3 marks)



- c) If the length of the call is in the lower 20% of calls, the call is not charged at all. Determine the longest length of the call in the lower 20% of calls. (2 marks)



- c) The consultant will stand outside a train station and will interview every fourth person who enters. Describe two sources of bias with this sampling scheme. (explain). (2 marks)

Mathematics Department


Perth Modern

In a particular city it is estimated that 30% of the population have blue eyes. A sample of 250 people is chosen and the number of blue eyed people are counted. The statistician will assume a Binomial distribution.

- a) State two reasons why the statistician believes that a Binomial distribution is appropriate.  
(2 marks)

Solution
The probability is independent of other people.
The probability is fixed.
Specific behaviours
P states one reason
P states two reasons

- b) Determine the probability that at least 75 people of the sample have blue eyes. (2 marks)

Solution

Specific behaviours
P uses Binomial with correct parameters
P states prob (Do not accept Normal-zero)


For each sample of 250 people the number of blue eyed people are counted and a sample proportion is calculated.

- c) State the **approximate** distribution of these sample proportions including all features.  
(2 marks)

Solution
Normal Distribution
Mean = 0.3
Standard deviation = 0.0290

Edit Action Interactive
$\text{invNormCDf("C", 0.82, 1, 0)}$ -1.340755034
$\text{solve}\left(0.015=1.340755034 \cdot \sqrt{\frac{0.5 \cdot 0.5}{n}}, n\right)$ $\{n=1997.360068\}$
n= 1998 Accept 1996-1999 due to various rounding of z score.
Specific behaviours
P sets up equation for n using p=0.5 P solves for decimal value P round up

- b) If the maximum sample size is 720 people, determine the maximum error in estimating a 95% confidence interval. (2 marks)

Solution

Specific behaviours
P determines z score P determines error (no need to round)

(4 marks)

- i) Another three samples of 250 people were taken and the number of blue eyed people were counted. See below. One of these samples is not from the city above. Explain which sample is and justify your answer using confidence intervals.

Solution	
<b>Specific behaviours</b>	
$\begin{array}{c} P \text{ states } Z \text{ score} \\ P \text{ states interval} \\ P \text{ rounds to 3 dp} \end{array}$	
$P \text{ divides } Z \text{ scores by root 3}$	
A traffic consultant has been hired to determine the proportion of people who drive to work.	

**Solution**

- a) A confidence interval of 82% will be used with an error within 0.015. Determine the minimum sample size that should be selected.

Q6 (7 marks) A traffic consultant has been hired to determine the proportion of people who drive to work.

Solution	
<b>Specific behaviours</b>	
$P \text{ determines all } Z \text{ scores}$	
$P \text{ divides } Z \text{ scores and multiplies by root 3}$	

- e) One sample of 250 people had 65 with blue eyes. Determine a 90% confidence interval for the population proportion to 3 decimal places.

Solution	
<b>Specific behaviours</b>	
$\text{normCDF}(0.35, \infty, 0.029, 0.3)$	

New width = double of  $2.285a = 4.57a$

Solution	
<b>Specific behaviours</b>	
$-1.644853627 \vee 3$	
$-2.170090378$	
$\text{invNormCDF}("C", 0.97, 1, 0)$	
$-1.644853627$	
$\text{invNormCDF}("C", 0.9, 1, 0)$	
$-2.170090378$	
$-1.644853627 \vee 3$	
$2.285131473$	

- d) Determine the approximate probability that a sample proportion will be greater than 0.35.

Solution	
<b>Specific behaviours</b>	
$P \text{ states Normal}$	
$P \text{ states mean and approx. st dev}$	

- c) Determine the new width of the confidence interval if a confidence of 97% is used and the sample size being one third of the original size. (2 decimal places)

Sample	Number of blue eyed people
A	55
B	72
C	41

Solution
<p>A: 0.177 - 0.263          B: 0.241 - 0.335          C: 0.126 - 0.203  <math>e</math>: 0.214 - 0.306</p> <p>Most probably Sample C as the confidence interval does not overlap with interval first calculated.          OR          We simply do not know whether any of the above intervals contain the true value of <math>p</math> as not all confidence intervals are expected to contain the true population <math>p</math>. Therefore no inference can be made.</p> <p><b>Specific behaviours</b></p> <p>P obtains at least one confidence interval of samples A,B&amp;C          P obtains confidence intervals for all samples          P gives an argument that mentions confidence intervals          P identifies sample C OR no inference possible</p>

Q5 (2, 2 &amp; 3 = 7 marks)


Consider a 90% confidence interval that has a margin of error of  $a$  units. Assume that the sample proportion does not change.

Answer in terms of  $a$ .

- a) Determine the new margin of error if the sample size is nine times the original.

Solution
$M \propto \frac{1}{\sqrt{n}} = \frac{1}{3}a$
Specific behaviours
<p>P uses inverse proportion with square root          P states in terms of <math>a</math></p>

- b) Determine the new margin of error if the sample size is unchanged but the confidence is decreased to 75%. (2 decimal places).

Solution
 <p>invNormCDF ("C", 0.9, 1, 0)  <math>-1.644853627</math></p> <p>invNormCDF ("C", 0.75, 1, 0)  <math>-1.15034938</math></p> $\frac{-1.15034938}{-1.644853627}a$ $0.6993627646 \cdot a$

Margin = 0.70a

**Specific behaviours**

P determines z scores for both confidences

P states in terms of  $a$  (no need to round)