

Question	Marks	Max	Question	Marks	Max
4	8	8			
3	8	8			
2	6	10			
1	5	8			
	6	13			

No other items may be taken into the examination room. It is your responsibility to ensure that you do not have any unauthorised material. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

**Important note to Candidates**

Special items: nil

Standard items: pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

**To be provided by the candidate**

Formula sheet

This Question/Answer booklet

Materials required/recommended for this section

Working time: fifty minutes  
Reading time before commencing work: five minutes

Your Teacher's Name:

Your Name:

Calculator-free

Section One:

UNIT 3 & 4

MATHEMATICS METHODS

Question/Answer booklet

Semester Two Examination, 2022



**Structure of this paper**

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	6	6	50	49	35
Section Two: Calculator-assumed	10	10	100	100	65
<b>Total</b>					<b>100</b>

**Instructions to candidates**


1. The rules for the conduct of the Western Australian Certificate of Education ATAR course examinations are detailed in the *Year 12 Information Handbook 2019*. Sitting this examination implies that you agree to abide by these rules.
2. Write your answers in this Question/Answer booklet.
3. You must be careful to confine your answers to the specific questions asked and to follow any instructions that are specific to a particular question.
4. Additional pages for the use of planning your answer to a question or continuing your answer to a question have been provided at the end of this Question/Answer booklet. If you use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number.
5. **Show all your working clearly.** Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
6. It is recommended that you **do not use pencil**, except in diagrams.
7. The Formula sheet is **not** to be handed in with your Question/Answer booklet.

$$k \left[ \frac{8}{(2x-4)^4} \right]_1^2 - k \left[ \frac{8}{(2x-4)^4} \right]_2^3 = 17$$

$$\int_2^3 k(2x-4)^3 dx - \int_1^2 k(2x-4)^3 dx = 17$$

**Solution**

If the area of the shaded region is 17 units<sup>2</sup>, determine the value of  $k$ .



The graph with equation  $y = k(2x-4)^3$  is shown below.

(5 marks)

**Question 1**

Working time: 50 minutes.

- Number of the question that you are continuing to answer at the top of the page. Fill in the original answer space where the answer is continued, i.e. give the page number.
- Continuing an answer: if you need to use the space to continue an answer, indicate this clearly at the top of the page.
  - Planning: if you use the spare pages for planning, indicate this clearly at the top of the page.
- Space pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.

This section has **six** questions. Answer all questions. Write your answers in the spaces provided.

**Section One: Calculator-free** (49 marks)

$$k([0-2] - [32-0]) = 17 - 34 \quad k = 17$$

$$k = \frac{-1}{2}$$

**Specific behaviours**

- ✓ Writes at least one definite integral to determine area.
- ✓ Recognises that the area from  $x=2$  to  $x=4$  is a signed area, and deals with this correctly.
- ✓ Correctly integrates using chain rule.
- ✓ Correctly substitutes boundaries and simplifies.
- ✓ Determines the value of  $k$ .

**Question 2 (10 marks)**

A random experiment can either result in a success with a probability of  $p$  or a failure.

In an event, this random experiment is conducted twice. Each experiment is independent of the other. Let the number of successes be represented by  $X$ .

- (a) State the distribution, and its parameters, that can be used to model the event described above. (2 marks)

**Solution**

$$X \text{ Bin}(2, p)$$

**Specific behaviours**

- ✓ Identifies the distribution as binomial.
- ✓ Writes down the correct parameters. .


The graph below shows the variance for the distribution in part (a) for various values of  $p$ .

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

Specific behaviours	
Solution	Mark
$Var(X) = 2 \times 0.5 \times 0.5 = 0.5$ $SD(X) = \sqrt{\frac{1}{2}}$	1 mark
In part (C)(i), Hence determine the exact value of the standard deviation, for the value of $p$	1 mark

P States value of $p$ .	
Specific behaviours	
Solution	Mark
$p = 0.5$	1 mark
(i) State the value of $p$ for which the standard deviation of $X$ is maximised.	1 mark

P Explains why the graph is continuous.	
Specific behaviours	
Solution	Mark
The probability is any value between 0 and 1, and hence the graph is continuous.	1 mark
(b) Explain why the graph is continuous, despite the distribution being discrete.	1 mark



End of questions

P Determines standard deviation

For the distribution in part (a), the probability that at least one of these experiments results in a success is 0.51.

- (d) (i) Show that  $(1-p)^2=0.49$  (2 marks)

**Solution**

$$\begin{aligned} P(X=0) &= 1 - 0.51 \\ (1-p)^2 &= 0.49 \end{aligned}$$

**Specific behaviours**

P Uses complement to  $P(X \geq 1)$ .  
P Shows that  $P(X=0)=(1-p)^2$ .

- (ii) Hence show that the value of  $p$  is 0.3. (1 mark)

**Solution**

$$\begin{aligned} 1-p &= 0.7 \\ p &= 0.3 \end{aligned}$$

**Specific behaviours**

P Correctly square roots 0.49 and determines  $p$ .

- (e) Determine  $E(X)$  for this distribution. (1 mark)

**Solution**

$$E[X]=0.6$$

ü smooth curve, concave up throughout

- (c) Given that  $\log_6(x+6)=\frac{\ln(x+6)}{\ln(6)}$ , determine the value of  $x$  where the slopes of  $y=f(x)$  and line  $L$  are the same. (2 marks)

**Solution**


$$\begin{aligned} f(x) &= -2 \log_6(x+6) - 1 = \frac{-2}{\ln(6)} \ln(x+6) - 1 \\ f'(x) &= \frac{-2}{\ln(6)} \times \frac{1}{x+6} = \frac{-2}{\ln(6)} \times \frac{1}{x+6} = \frac{-2}{x+6} \rightarrow x = \frac{5}{\ln(6)} - 6 \end{aligned}$$

**Specific behaviours**

P correctly differentiates  $f$   
ü solves for  $x$ -coordinate and fully simplified  
Note- no follow through if mistake makes solving too easy

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF






- b) A probability distribution of a random variable  $X$  is given by

$$P(X = x) = \frac{x}{15} \text{ where } x = 1, 2, 3, 4, 5$$

- (i) Graph the probability function on the axes below. (2 marks)



- (ii) Determine the probability of obtaining a value of  $X$  less than 3. (1 mark)

DO NOT WRITE IN THIS AREA AS IT WILL BE CUT OFF

- (b) The current,  $I$  amps, flowing through component B reaches a peak very quickly and then declines as time goes on, as modelled by  $I(t) = \frac{2+ln t}{4t}$ . Determine, in simplest form, the maximum current that flows through this component. (4 marks)

<b>Solution</b>
$I'(t) = \frac{\left(\frac{1}{t}\right)(4t) - (2+ln t)(4)}{(4t)^2} = \frac{4-4(2+ln t)}{4 \times 4t^2} = \frac{-1-4ln t}{4t^2}$
$I'(t) = 0 \Rightarrow ln t = -1t = e^{-1}$
$I(e^{-1}) = \frac{2-1}{4e^{-1}} = \frac{e}{4}$
Maximum current is $\frac{e}{4}$ amps.
<b>Specific behaviours</b>
P uses quotient rule ü obtains correct derivative and equates to zero ü obtains root of derivative ü calculates maximum current in simplified form

(p) 
$$\int_x^1 \frac{dx}{t} \cos(t+1) dt.$$

<b>Solution</b>	$\cos(x+1)$
<b>Specific behaviours</b>	$P(\cos(x+1))$
<b>Correct result</b>	

(1 mark)

(e) 
$$\int_3^0 \frac{dx}{t} \left[ e^{2x} \right] = 3e^6$$

<b>Solution</b>	$\left[ e^{2x} \right]_0^3 = 3e^6$
<b>Specific behaviours</b>	$P(e^{2x})$
<b>Correct result</b>	

(1 mark)

(iii) Determine  $P(X > 1 | X \leq 4)$ . (2 marks)

<b>Solution</b>	$\frac{9}{15} = \frac{3}{5}$
<b>Specific behaviours</b>	$P(\text{numerator})$
<b>Specific behaviours</b>	$P(\text{denominator})$
<b>Components A and B form part of an electronic circuit, and properties of these components are measured } t \text{ seconds after the circuit is turned on.}</b>	
<b>(a)</b> The rate of change of temperature, $T^\circ\text{C}$ , of component A is given by $\frac{dT}{dt} = \frac{18t}{3t+8}$ .	4 marks

Determine, in simplest form, the increase in temperature of this component during the first 4 seconds.

<b>Solution</b>	$\Delta T = \int_0^4 \frac{18t}{3t+8} dt = 3 \int_0^4 \frac{6t}{3t+8} dt = 3 \left[ \ln(3t+8) \right]_0^4 - \ln(56) + \ln(8) = 3 \ln(7) + C$
<b>Specific behaviours</b>	$P(\text{determines at least two correct probs})$
<b>Specific behaviours</b>	$P(\text{all probs correct})$
<b>(iv)</b> State the cumulative probability distribution for $X$ . (2 marks)	

<b>Solution</b>	$P(X \leq 1) = \frac{1}{15}, P(X = 2) = \frac{3}{15}, P(X = 3) = \frac{6}{15}, P(X = 4) = \frac{10}{15}, P(X = 5) = \frac{15}{15}$
<b>Specific behaviours</b>	$P(\text{determines at least two correct probs})$
<b>Specific behaviours</b>	$P(\text{all probs correct})$
<b>(v)</b> State the cumulative probability distribution for $X$ . (2 marks)	

<b>Solution</b>	$\frac{15}{15} = \frac{5}{1}$
<b>Specific behaviours</b>	$P(\text{determines prob})$
<b>Specific behaviours</b>	$P(\text{determines prob})$

(p) 
$$\int_x^1 \frac{dx}{t} \cos(t+1) dt.$$

**Question 4**

(8 marks)

Determine the following:

(a)  $\int 6 e^{2x-3} dx.$

(1 mark)

<b>Solution</b>
$3e^{2x-3} + c$
<b>Specific behaviours</b>
P correct antiderivative, with constant of integration

(b)  $\int_0^{\frac{\pi}{8}} \sin(4x) dx.$

(2 marks)

<b>Solution</b>
$\left[ -\frac{1}{4} \cos(4x) \right]_0^{\pi/8} = 0 - \left( -\frac{1}{4} \right) = \frac{1}{4}$
<b>Specific behaviours</b>
P correct antiderivative ü correct value

(c)  $f\left(\frac{\pi}{6}\right)$  when  $f(x) = \frac{\cos(3x)}{2+\sin(x)}.$

(3 marks)

<b>Solution</b>
$f'(x) = \frac{-3\sin(3x)(2+\sin(x)) - \cos(3x)\cos(x)}{(2+\sin(x))^2} f\left(\frac{\pi}{6}\right) = \frac{-3(2+0.5)-0}{(2+0.5)^2} \textcolor{red}{i} - \frac{3}{2.5} = \frac{-6}{5}$
<b>Specific behaviours</b>
P correctly uses quotient rule ü correctly differentiates all trig terms ü correctly evaluates