



## Hale School Semester Two Examination Sample

Write your name below:

---

### CHEMISTRY Stage 3

For Examiners only	
Part 1	
Part 2	
Part 3	
Total	

#### Time allowed for this paper

Reading time before commencing work: ten minutes

Working time for paper: three hours

#### Materials required/recommended for this paper

##### *To be provided by the supervisor*

This Question/Answer Booklet

Multiple-choice Answer Sheet

Chemistry Data Sheet

##### *To be provided by the candidate*

Standard items: pens, pencils, eraser, correction fluid/tape, ruler, highlighters

Special items: non-programmable calculators satisfying the conditions set out by the Curriculum Council for this course

#### Important note to candidates

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised notes or other items of a non-personal nature in the examination room. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

## Structure of this paper

Section	Number of questions available	Number of questions to be answered	Suggested working time (minutes)	Marks available	Percentage of exam
Section One: Multiple-choice	25	25	50	50	25
Section Two: Short answer	12	12	60	70	35
Section Three: Extended answer	5	5	70	80	40
					100

## Instructions to candidates

- The rules for the conduct of Curriculum Council examinations are detailed in the *Student Information Handbook*. Sitting this examination implies that you agree to abide by these rules.
- Answer the questions according to the following instructions:

### Section One

Answer **all** questions, using a 2B, B or HB pencil, on the separate Multiple Choice Answer Sheet provided. Do not use a ball point or ink pen.

### Section Two

Answer in the spaces provided in this Question/Answer Booklet.

### Section Three

Write your answers in the Section 3 Question/Answer Booklet which is provided.

- A blue or black ball point or ink pen should be used. Marks for answers in pencil (besides section 1) can't be contested.
- For full marks, chemical equations should refer only to those species consumed in the reaction and the new species produced. These species may be **ions** [for example  $\text{Ag}_{(\text{aq})}^+$ ], **molecules** [for example  $\text{NH}_{3(\text{g})}$ ,  $\text{NH}_{3(\text{aq})}$ ,  $\text{CH}_3\text{COOH}_{(\text{l})}$ ,  $\text{CH}_3\text{COOH}_{(\text{aq})}$ ] or **solids** [for example  $\text{BaSO}_{4(\text{s})}$ ,  $\text{Cu}_{(\text{s})}$ ,  $\text{Na}_2\text{SO}_{4(\text{s})}$ ]

See next page

**Section One: Multiple-choice****25% (25 Marks)**

This section has **25** questions. Answer **all** questions on the separate Multiple-choice Answer Sheet provided. For each question shade the box to indicate your answer. Use only a blue or black pen to shade the boxes. If you make a mistake, place a cross through that square, do not erase or use correction fluid, and shade your new answer. Marks will not be deducted for incorrect answers. No marks will be given if more than one answer is completed for any question.

Suggested working time for this section is 50 minutes.

---

1. Which of the following best describes the molecular shape and molecular polarity of a chloroform molecule whose formula is  $\text{CHBr}_3$ ?

- (a) pyramidal, non polar
- (b) tetrahedral, non polar
- (c) pyramidal, polar
- (d) tetrahedral, polar

2. An element X has the following five successive ionisation energies (in  $\text{kJmol}^{-1}$ )

680	1600	8000	11600	14500
-----	------	------	-------	-------

What would be the formula of the compound formed when “X” reacts with oxygen ?

- (a)  $\text{X}_2\text{O}$
- (b)  $\text{XO}$
- (c)  $\text{X}_2\text{O}_3$
- (d)  $\text{XO}_2$

3. When  $1.0 \text{ mol L}^{-1}$  solutions of the following are mixed, which combinations will result in the formation of precipitates?

- i)  $\text{Ba}(\text{NO}_3)_2$  and  $\text{HCl}$
- ii)  $\text{Ca}(\text{NO}_3)_2$  and  $\text{Na}_2\text{CO}_3$
- iii)  $\text{Cu}(\text{NO}_3)_2$  and  $\text{KOH}$
- iv)  $\text{Pb}(\text{NO}_3)_2$  and  $\text{H}_2\text{SO}_4$

- (a) i), ii) and iii) only
- (b) ii) and iii) only
- (c) i), ii), iii) and iv)
- (d) ii), iii) and iv) only


See next page

4. The conjugate base of the acid  $\text{HCrO}_4^-$  is:
- (a)  $\text{H}_2\text{CrO}_4$
  - (b)  $\text{H}_2\text{CrO}_4^-$
  - (c)  $\text{CrO}_4^{2-}$
  - (d)  $\text{CrO}_4^-$
5. Which of the following physical properties **decrease** with increasing atomic number for both the alkali metals and the halogens?
- I. Atomic radius
  - II. Ionization energy
  - III. Melting point
- (a) I only
  - (b) II only
  - (c) III only
  - (d) I and III only
6. Which of the following equations represents a redox equation? (States not included)
- (a)  $\text{NaOH} + \text{HNO}_3 \rightarrow \text{NaNO}_3 + \text{H}_2\text{O}$
  - (b)  $2\text{AgNO}_3 + \text{Cu} \rightarrow 2\text{Ag} + \text{Cu}(\text{NO}_3)_2$
  - (c)  $\text{H}_2\text{SO}_4 + 2\text{KOH} \rightarrow \text{K}_2\text{SO}_4 + 2\text{H}_2\text{O}$
  - (d)  $\text{CaCl}_2 + \text{Ba}(\text{OH})_2 \rightarrow \text{Ca}(\text{OH})_2 + \text{BaCl}_2$
7. Which one of the following solids contains covalent bonds only?
- (a)  $\text{SiO}_2$
  - (b)  $\text{MgO}$
  - (c)  $\text{NH}_4\text{Br}$
  - (d)  $\text{Ne}$
8. If the pH of a solution changes from 2 to 4, then the hydronium ion concentration
- (a) is doubled.
  - (b) is halved.
  - (c) increases by a factor of 100.
  - (d) decreases by a factor of 100.

9. A crystal of iodine,  $I_2$ , produces a purple vapour when gently heated. Which pair of statements correctly describes this process?

	Type of bond broken	Formula of purple species
(a)	covalent	I
(b)	covalent	$I_2$
(c)	dispersion forces	$I_2$
(d)	dipole-dipole	$I_2$

10. Household bleach contains sodium hypochlorite, NaClO, as the active ingredient. The concentration of NaClO in the bleach can be determined by reacting a known amount with aqueous hydrogen peroxide,  $H_2O_2$ , according to the equation:




When 25.0 mL of bleach is treated with an excess of aqueous  $H_2O_2$ , 0.0350 mol of oxygen gas is given off.

What is the concentration of NaClO in the bleach?

- (a) 1.40 mol  $L^{-1}$   
 (b) 0.700 mol  $L^{-1}$   
 (c) 0.875 mol  $L^{-1}$   
 (d)  $8.75 \times 10^{-4}$  mol  $L^{-1}$

11. In the contact process reaction:



If the equilibrium system **temperature** is increased, what effect will this have on the equilibrium constant, K, and the yield?

	Equilibrium constant, K	Yield increase
(a)	decrease	products
(b)	decrease	reactants
(c)	increase	products
(d)	increase	reactants


12. Deposits of ammonium compounds, including ammonium sulfate, have been discovered in areas of high atmospheric pollution. A chemical reaction believed to occur is:




What does **not** occur in this reaction?

- (a) acid/base neutralisation
  - (b) coordinate (dative) bond formation
  - (c) oxidation/reduction
  - (d) ionic bond formation
13. Galvanic cells are used as portable sources of electrical energy. One common cell is the rechargeable nickel-cadmium cell.

#### Nickel–Cadmium Cell



The net equation representing the discharge of the nickel-cadmium cell is:




The reaction at the **anode** during the discharge of the cell is:

- (a)  $\text{Cd}(\text{s}) + 2\text{OH}^-(\text{aq}) \rightarrow \text{Cd}(\text{OH})_2(\text{s}) + 2\text{e}^-$
- (b)  $\text{Cd}(\text{s}) + 2\text{OH}^-(\text{aq}) + 2\text{e}^- \rightarrow \text{Cd}(\text{OH})_2(\text{s})$
- (c)  $\text{NiO}_2(\text{s}) + 2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightarrow \text{Ni}(\text{OH})_2(\text{s}) + 2\text{OH}^-(\text{aq})$
- (d)  $\text{NiO}_2(\text{s}) + 2\text{H}_2\text{O}(\text{l}) \rightarrow \text{Ni}(\text{OH})_2(\text{s}) + 2\text{OH}^-(\text{aq}) + 2\text{e}^-$


See next page

14. Which one of the following has the same electronic arrangement as  $\text{Li}^+$ ?
- $\text{Na}^+$
  - $\text{Be}^{2+}$
  - $\text{F}^-$
  - Ne
15. The largest mass of silver chloride is precipitated when an excess of silver nitrate solution is added to:
- 25.0 mL of a  $0.800 \text{ mol L}^{-1}$  solution of hydrochloric acid.
  - 30.0 mL of a  $0.300 \text{ mol L}^{-1}$  solution of iron(III) chloride.
  - 50.0 mL of a  $0.200 \text{ mol L}^{-1}$  solution of magnesium chloride.
  - 50.0 mL of a  $0.500 \text{ mol L}^{-1}$  solution of sodium chloride.
16. The IUPAC name for the structure below is:



- 2,2,5-trimethylheptane
- 3,6,6-trimethylheptane
- 2-ethyl-5,5-dimethylhexane
- 5-ethyl-2,2-dimethylhexane





17. Which one of the following species does **not** have eight valence electrons surrounding the central atom?
- (a)  $\text{CHCl}_3$  molecule
  - (b)  $\text{NO}_2$  molecule
  - (c)  $\text{NH}_4^+$  ion
  - (d)  $\text{OF}_2$  molecule
18. Select, from the list below, the compound that can be polymerised to give:



- (a) 2-methylbut-1-ene
- (b) 2-methylbut-2-ene
- (c) pent-2-ene
- (d) pent-1-ene




19. Which of the following rows identifies the structural diagram and the corresponding IUPAC name of the compound with the chemical formula,  $C_8H_{16}$ ?

	Structural Diagram	IUPAC Name
(a)		ethylbenzene
(b)		ethylcyclohexane
(c)		cyclopentylpropane
(d)		propylcyclopentene


20. When the compounds HF,  $H_2O$ ,  $NH_3$ , and  $CH_4$  are listed in order of increasing boiling point, which order is correct?
- (a)  $CH_4 < NH_3 < H_2O < HF$   
 (b)  $NH_3 < CH_4 < H_2O < HF$   
 (c)  $CH_4 < NH_3 < HF < H_2O$   
 (d)  $HF < CH_4 < H_2O < NH_3$
21. The reductant that can convert  $1.0\text{ M Fe}^{3+}(\text{aq})$  to  $\text{Fe}^{2+}(\text{aq})$  but not  $1.0\text{ M Sn}^{2+}(\text{aq})$  to  $\text{Sn}(\text{aq})$ , at STP is:
- (a)  $\text{Cu}(\text{s})$   
 (b)  $\text{Au}(\text{s})$   
 (c)  $\text{Ni}(\text{s})$   
 (d)  $\text{HOOC}(\text{COOH})$

22. A cell was incorrectly connected, as shown below. Which statement is **incorrect**?



- (a) The anode is the zinc electrode.
- (b) There would be no electron current flow from one half cell to the other.
- (c) If electrodes are interchanged the cell emf (potential difference) would be  $-1.1\text{ V}$  (at  $25^\circ\text{C}$ ).
- (d) The concentration of  $\text{Cu}^{2+}$  ions will decrease.
23. Which of the following statements is **correct**?
- (a) Covalent network solids include diamond, graphite and sulfur.
- (b) Metal solids and ionic solids exhibit non-directional interparticle bonding.
- (c) Ionic solids conduct electricity very well in the aqueous and solid states.
- (d) Heated covalent molecular solids tend to decompose before melting.
24. Which of the following statements about the third row of the Periodic Table is correct?
- (a) Elements on the right side of the row form acidic oxides, whilst those on the left side form basic oxides.
- (b) Elements on the left side of the row have a greater range of oxidation states than elements on the right side.
- (c) Elements on the right side of the row are stronger reducing agents than elements on the left side.
- (d) Electronegativity decreases across a row from left to right of the period.

25. A common painkiller has the structure:



Which of the options below best represents its characteristics?

	<b>Type</b>	<b>Functional groups</b>
(a)	aromatic	carboxyl, hydroxyl
(b)	aliphatic	hydroxyl, alkene
(c)	aromatic	hydroxyl, ester
(d)	aliphatic	carbonyl, hydroxyl

**END OF SECTION 1**

**Section Two: Short answer****35% (70 Marks)**

This section has **12** questions. Answer **all** questions. Write your answers in the space provided.

Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.

- Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
- Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question(s) that you are continuing to answer at the top of the page.

**Suggested working time for this section is 60 minutes.**

---

**Question 26****(5 marks)**

A buffer solution is needed for preserving “Tango” fruit juice. A chemist at the fruit juice company prepared a benzoic acid/sodium benzoate buffer with concentrations of  $0.105 \text{ mol L}^{-1}$   $\text{C}_6\text{H}_5\text{COOH}$  and  $0.125 \text{ mol L}^{-1}$   $\text{C}_6\text{H}_5\text{COONa}$ .

- (a) What is a “buffer solution”, and what is its purpose? (2 mark)

---

---

---

- (b) Explain, using Le Chatelier’s principle, how this solution acts as a buffer solution. (use equations in your answer). (3 marks)

---

---

---


---

---

**Question 27****(2 marks)**

See next page

Toluene (methyl benzene),  $\text{C}_7\text{H}_8$  (g) is an important solvent and precursor to many other organic compounds such as trinitrotoluene (TNT). It can be produced according to the following equilibrium:



When 3.00 mol of  $\text{C}_7\text{H}_{14}$  (g) was introduced into a 1.00 L container, 1.20 mol of  $\text{H}_2$ (g) was produced at equilibrium.

What were the equilibrium concentrations, in  $\text{mol L}^{-1}$ , of  $\text{C}_7\text{H}_8$  and  $\text{C}_7\text{H}_{14}$ ?

---



---

### Question 28

(4 marks)

Write **observations** for any reactions that occur in the following procedures. In each case describe in full what you would observe, including any:

- colours
- odours
- precipitates (give the colour)
- gases evolved (give the colour or describe as colourless).

If no change is observed, you should state this.

- (a) Potassium sulfide solution is added to lead (II) nitrate solution. (2 marks)

Observation: \_\_\_\_\_

---



---



---

- (b) Sodium metal is added to pentanol. (2 marks)

Observation: \_\_\_\_\_

---



---




---

### Question 29

(3 marks)

- (a) Consider the reaction half equations and then balance the following redox equation:

See next page




---



---



---



---

- (b) Re-write the redox equation, for alkaline (basic) conditions. (1 mark)

---



---

### Question 30

(4 marks)

Write the equation for the reaction that occurs in each of the following procedures. If no reaction occurs, write 'no reaction'. For full marks, chemical equations should refer only to those species consumed in the reaction and the new species produced. These species may be ions [for example  $\text{Ag}^+(\text{aq})$ ], molecules [for example  $\text{NH}_3(\text{g})$ ,  $\text{NH}_3(\text{aq})$ ,  $\text{CH}_3\text{COOH}(\ell)$ ] or solids [for example  $\text{BaSO}_4(\text{s})$ ,  $\text{Cu}(\text{s})$ ,  $\text{Na}_2\text{CO}_3(\text{s})$ ].

- (a) Potassium phosphate solution is added to copper (II) sulfate solution. (2 marks)

Equation: \_\_\_\_\_

- (b) Propene gas is shaken with hydrogen bromide gas in the presence of uV light. (2 marks)


Equation: \_\_\_\_\_

### Question 31

(7 marks)

Commercial buses, and cars, can operate using a proton exchange membrane fuel cell (PEM) to provide the required energy. They use a solid polymer sandwiched between two sheets of carbon fibre paper as an electrolyte, and porous carbon

See next page



electrodes containing a platinum catalyst. They need only hydrogen, oxygen from the air, and water to operate. They are typically fuelled with pure hydrogen supplied from storage tanks or onboard reformers. The cell operates at a temperature of around 80°C.

A Picture of the Hydrogen Fuel Cell Transit Bus



- (a) What is **one** advantage of a solid polymer membrane electrolyte over the liquid chemical electrolyte, KOH (used in other fuel cells)? (1 mark)

---



---

- (b) State **one** environmental advantage, and **one** sustainability advantage; of this PEM fuel cell over the use of conventional fuels such as diesel? (2 marks)

---



---



---

- (c) What are **two** disadvantages of the  $O_2/H_2$  fuel cell, as energy sources, over conventional fuels such as diesel? (2 marks)

---



---



---

- (d) Some fuel cells use methanol as a source of hydrogen (as  $H^+$ ), which combines with oxygen to produce carbon dioxide and water. Write an overall balanced redox equation given the following skeleton equations;  $CH_3OH \rightarrow CO_2$  and  $O_2 \rightarrow H_2O$  (2 marks)

---




---



---

**Question 32****(8 marks)**

On heating, a mixture of potassium and bromine react to form potassium bromide, according to the equation:



- (a) The melting points of potassium, bromine and potassium iodide are 63.3 °C, -7.2 °C and 734 °C respectively. For each of the substances state the type of **inter-particle** bonding present and the nature of the attractive forces holding each substance together. (6 marks)

Substance	Type of bonding	Nature (strength) of bonding
Potassium		
Bromine		
Potassium bromide		

- (b) Briefly explain why the melting point of bromine is much lower than that of sodium bromide. (2 marks)

---



---



---



---



---

**Question 33****(15 marks)**

- (a) For each species listed in the table below, draw the structure, representing all valence shell electron pairs either as : or as — **and** state or draw the shape of the molecule or ion. (9 marks)

(for example, water  $\text{H}:\ddot{\text{O}}:\text{H}$  or  $\text{H}-\ddot{\text{O}}-\text{H}$  or  $\text{H}-\overline{\text{O}}-\text{H}$  bent)

Compound	Electron-dot structure (showing all valence shell electrons)	Shape (sketch or name)
----------	---	---------------------------

See next page



Carbon disulfide $\text{CS}_2$		
Strontium nitrate $\text{Sr}(\text{NO}_3)_2$		
Urea $\text{H}_2\text{NCONH}_2$		

- (b) Compare, and explain, the molecular polarity of carbon disulfide and urea. (6 marks)

Compound	Polar or non-polar	Explanation
$\text{CS}_2$		

See next page

H <sub>2</sub> NCONH <sub>2</sub>		

**Question 34****(5 marks)**

- (a) **Draw and label** the geometric isomeric forms of 2-pentene (pent-2-ene). (3 marks)

Form:	Form:
-------	-------

- (b) What chemical test could be used to distinguish between pent-2-ene and pentane?  
(chemical equation required) (2 marks)

---



---



---



---




---

**Question 35****(10 marks)**

Industrially, calcium carbonate can be thermally decomposed (roasted) to form calcium oxide ('quicklime'). This product is used to change pH, in calcium silicate brick manufacture, in aluminium and gold production, and in the building industry (plaster).

The chemical reaction for its production is:

See next page



- (a) What is the  $K_{\text{eq}}$  expression for the reaction? (1 mark)

- (b) In practice, conditions can be changed to alter the rate of reaction and the yield of product. Indicate any effects of imposed change on the system in the table below. (9 marks)

Imposed change	Effect on forward reaction rate	Effect on reverse reaction rate	Effect on the value of K
Increase the partial pressure of the carbon dioxide			
The temperature is decreased			
Increase the surface area of the $\text{CaCO}_3$			

### Question 36

(3 marks)

Older second hand cars can have bubbling of paint on the panels due to iron corrosion (“rusting”) under the sealed paint (i.e. an anodic region). This is often noticed close to the wheel hubs or the base of doors or windows. The paint under normal circumstances provides a physical barrier to corrosion.

See next page

- (a) What is the likely cause of the corrosion? (2 marks)

---

---

---

- (b) Suggest a practical chemical solution to the problem. (1 mark)

---

---

---

**Question 37**

**(4 marks)**

Vinegar is about 4% by mass acetic acid and is safe to consume in foods. The same concentration sulfuric acid is not safe to consume. Explain why. Include equations.

---

---

---

---

---

---

---

---

---

---

---

**END OF SECTION 2**

**EXTRA WORKING SPACE**

---

---

---

See next page





**Write your name below:**

---

# CHEMISTRY

## Stage 3

### Section Three: Extended answers

**40% (80 Marks)**

This section contains **five (5)** questions. You must answer **all** questions. Write your answers in the space provided in this question and answer booklet. Answer should be in blue or black pen.

Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.

- Planning: If you use the spare pages for planning, indicate this clearly at the top of the page.
- Continuing an answer: If you need to use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number. Fill in the number of the question(s) that you are continuing to answer at the top of the page.

Suggested working time for this section is 70 minutes.

---


### Question 38

**(14 marks)**

A student carried out two acid-base practical investigations, at 25°C:

See next page

A titration was performed where 20.0 mL of an unknown solution was pipetted into a conical flask and titrated with another unknown solution from a burette. The pH was monitored with a pH meter, recorded with a data logger and the results displayed on a computer screen throughout the experiment. The changes in pH are shown below:



- (a) What is meant by the term “equivalence point”? (1 mark)

---

---

---

- (b) What was the pH of the mixture at the equivalence point? (1 mark)

---

- (c) (i) A pH meter had to be used in this investigation. Explain, why there was no suitable acid-base indicator that could be used for this titration? (1 mark)

---

---

- (ii) What does this tell us about the acid and base added together? (1 mark)

---



---

---

(iii) Give an example of the acid and base that could have been used to give these results? (1 mark)

Acid\_\_\_\_\_ Base\_\_\_\_\_

(iv) Which species was in the conical flask? (1 mark)

---

(e) What was the concentration of  $\text{OH}^-(\text{aq})$  ( $\text{mol L}^{-1}$ ) in the mixture after 15.0 mL of titrant had been added? (2 marks)

---

---

---

---

A commercial brand of antacid, "Easiflux", was tested to see how much acid it could neutralize. A 5.00 mL sample of the mixture was used where the active ingredient was stated to be magnesium hydroxide. The mixture was reacted with 12.90 mL of  $72.9 \text{ g L}^{-1}$  hydrochloric acid, HCl (stomach acid).

(f) Write a balanced equation for this reaction. (1 mark)

---

(g) How many moles of HCl reacted? (2 marks)

---

---

---

(h) What mass (g) of magnesium hydroxide would react with this amount of HCl? (2 marks)

---

See next page


- 
- 
- (i) The suggested dose for bad indigestion is 800 mg of  $\text{Mg}(\text{OH})_2(\text{s})$  in a 5.00 mL dose. How does the sample tested compare with this? (1 mark)
- 
- 

**Question 39****(17 marks)**

There are many ways of observing and measuring the rates of chemical reactions. A student investigated one reaction involving colour change; an “iodine clock” reaction.

At room temperature, a reaction occurs when potassium iodate solution is mixed with sodium hydrogen sulfate solution that contains a small amount of starch. In a laboratory, 12.00 mL of a 0.0160 M  $\text{NaHSO}_3(\text{aq})$  solution containing starch were placed in each of six test tubes. Different volumes of 0.0240 M  $\text{KIO}_3(\text{aq})$  and enough distilled water to maintain a constant volume were added to each test tube and the time taken for the dark-blue colour to appear was measured. The data were recorded in the table over the page.

The chemical reaction is:



- (a) Is this a redox reaction? Justify your answer using half equations and oxidation numbers (3 marks)
- 
- 
- 
- 

- (b) Describe a procedure for diluting the stock 0.0240 mol  $\text{L}^{-1}$   $\text{KIO}_3$  solution to give a 0.00800 mol  $\text{L}^{-1}$  reaction mixture test solution. (2 marks)
-

---

---

---

(c) What is the independent variable for the investigation? (1 mark)


---

(d) What variables need to be controlled? (1 mark)

---

(e) Complete the table below, and graph the data of reaction rate ( $1/t$ ) versus concentration of potassium iodate ( $\text{mol L}^{-1}$ ). (5 marks)

Concentration of $\text{IO}_3^-$ ( $\text{mol L}^{-1}$ ) in reaction mixture.	Reaction time (s).	Reaction rate, $1/t$ ( $\text{s}^{-1}$ ).
0.00200	210	
0.00400	88	
0.00600	49	
0.00800	39	
0.0100	33	
0.0120	27	



See next page

- (f) What conclusion can be drawn from the graph about the relationship between the rate of the reaction and the concentration of the potassium iodate? (1 mark)

---

---

- (e) Use your graph to predict the time taken for a  $0.00500 \text{ mol L}^{-1}$  potassium iodate solution, at standard room temperature, to react. (1 mark)

---

- (f) The procedure described above was repeated at a number of different temperatures between  $10^\circ\text{C}$  and  $70^\circ\text{C}$  and the reaction times were measured as before. Would you expect the reaction times to increase, decrease, or stay the same, as the temperature was increased? Use Collision Theory to justify your answer. (3 marks)

---

---


---

---

---

**Question 40****(15 marks)**

Iron is one of the most abundant metals on Earth, is essential to most forms of life and to normal human physiology. Sometimes people take iron supplements. The iron content of a particular brand of iron tablets was determined by titration with a freshly standardised solution of potassium permanganate,  $\text{KMnO}_4$ . The equation for the titration reaction is:



- (a) Why are iron tablets sometimes medically prescribed? (1 mark)

---

---

- (b) What is the oxidation number of manganese in the permanganate ion? (1 mark)

---

- (c) (i) Why must potassium permanganate solutions be standardised? (2 mark)

---

---

- (ii) What reagent is used for this purpose? (1 mark)

---

**250 mL of  $\text{Fe}^{2+}$  solution was prepared from ten (10) iron tablets, each of mass 0.328 g.**

- (e) Explain why additional dilute sulfuric acid must be added to the titration flask before each titration is carried out. (1 mark)

---

---

- (f) How was the end-point detected? (1 mark)

---

**A standardised 0.0100 M potassium permanganate was used to react with 25.0 mL portions of the iron solution prepared from the ten tablets.**

- (g) A number of titrations were performed and the following titre values obtained.

See next page

	Trial	1	2	3
Titre (mL)	21.00	18.79	18.76	18.70

What is the average titre used? (1 mark)

---

(g) What is the **concentration** ( $\text{mol L}^{-1}$ ) of the  $\text{Fe}^{2+}$  solution? (3 marks)

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

(h) What is the **total mass** (mg) of iron in one tablet? (3 marks)

---

---

---

---

---

---

---

---

- (i) What is the **percentage, by mass**, of iron in each tablet? (1 mark)

---

---

---

---


---

---

**Question 41**

**(22 marks)**

Many drugs are produced by chemically combining several molecules. A newly developed amino acid, in combination with another substance, has the potential to be a new anti-inflammatory drug. The structure of the amino acid ( $M_r = 179.214$ ) is:



- (a) (i) Explain, why it is considered to be an amino acid? (2 marks)

---

See next page


(ii) Would you expect this molecule to be reasonably soluble in water? Explain.

(2 marks)


The amino acid above exists as a zwitterion in aqueous solution. Zwitterion is the general name given these compounds when they form an ion which has a positive charge on one atom and a negative charge on another.

(b) Draw the structure of this **zwitterion**.

(2 marks)



The anti-inflammatory drug is made by combining the amino acid, with the molecule shown below, called compound X. R represents a small side chain.



(c) (i) Combine both structures to show the structure of the **drug**.

(1 mark)



See next page



--

(ii) What type of chemical linkage forms? (1 mark)

---

(iii) What type of reaction occurs, and what small molecule is released? (2 marks)

---

---

(d) Spectroscopy analysis showed compound **X** contained the elements C, H, O, and chlorine (Cl).

Experimentally, the identity of **R** was found by combusting 0.425 g of compound **X** in excess air. It was found that 0.844 g of carbon dioxide and 0.171 g of water was produced.

Further testing of a further 0.396 g sample of compound **X**, produced 25.7 mLs of chlorine gas ( $\text{Cl}_2$ ), under conditions of 40 °C and 100 kPa.

(i) Determine the **percentage composition**, by mass, of each element in compound **X**. (7 marks)

---

---

---

(3 marks)

---

---

---

---

---

(2 mark)

---

---

---

**Question 42****(12 marks)**

Butanoic acid,  $\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH}$ , is a ubiquitous, oily, colourless substance with a diverse range of origins and uses. It occurs naturally but can be manufactured through the fermentation of sugar and starch, and then the addition of putrefying cheese, with calcium carbonate added to neutralize the acid.

- (a) **Draw** and **name** two structural isomers of  $\text{C}_4\text{H}_8\text{O}_2$  (other than butanoic acid). (4 marks)

Name: \_\_\_\_\_

See next page

Name:

Butanoic acid is a rancid smelling substance that gives parmesan cheese its characteristic odour. It has been used as a nausea inducing repellent by anti-whaling protesters against whalers. Butanoic acid can undergo **esterification** with ethanol,  $\text{CH}_3\text{CH}_2\text{OH}$ , to form a much more pleasant smelling pineapple flavoured ester.

- (b) What is the common **catalyst** used for this reaction? (1 mark)

---

- (c) Draw, and name the **ester** formed. (2 marks)

Name:

Fats and oils can be removed from clothing by the action of soaps. Soaps are typically sodium or potassium salts of long chain fatty acids e.g.  $\text{CH}_3(\text{CH}_2)_{16}\text{COONa}$ .

- (5 marks)

[illegible]

[illegible]

**END OF SECTION 3**

[illegible]

