


Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until Friday June 15th 2018.



MATHEMATICS SPECIALIST UNIT 3

Semester One

2018


SOLUTIONS

Calculator-free Solutions

1. Since $z = \pm 2$ are roots, then $z^2 - 4$ is a factor. ✓
 \therefore dividing $P(z)$ by $z^2 - 4$ gives the quadratic factor $2z^2 - 4z + 4$ ✓
 and using the quadratic formula gives $z = 1 \pm i$ ✓✓
 $\therefore P(z) = 2(z+2)(z-2)(z-1-i)(z-1+i)$ ✓ [5]


2. (a) $w = z^3 = \left[3 \operatorname{cis} \left(\frac{-\pi}{2} \right) \right]^3 = 27 \operatorname{cis} \left(\frac{-3\pi}{2} \right) = 27 \operatorname{cis} \frac{\pi}{2}$ ✓✓

(b)



- ✓ magnitude = 3
 ✓ $\frac{2\pi}{3}$ radians apart

(c)



$x^2 = 9 + 9 - 2(9) \cos 120^\circ$ ✓
 $\therefore x = 3\sqrt{3}$ ✓
 \therefore perimeter is $9\sqrt{3}$ units, ✓ [7]


3. (a) $3a + 7 = 0$ ✓
 $\therefore a = -\frac{7}{3}$ ✓
 (b) $a \neq -\frac{7}{3}$ ✓

$$\therefore z = \frac{25}{3a+7}, y = \frac{23a+87}{3a+7}, x = \frac{5a+20}{3a+7}$$

✓✓✓

[6]

4. (a)



✓ intersects $f(x)$
along the line $y=x$

✓ correct shape and
location

(b) $f^{-1}(x) = -(x+1)^2 + 4 = -x^2 - 2x + 3$ ✓✓

Domain $\{x \in R : x \geq -1\}$ ✓

(c) $f(g) = \sqrt{4-g} - 1 = \sqrt{x^2} - 1 = |x| - 1$ ✓

(d) Condition for composition to exist: $4 - g(x) \geq 0$ ✓

$$4 - 4 + x^2 \geq 0$$

$$x^2 \geq 0$$

$$\therefore x \in R$$
 ✓

$$\therefore \text{no changes needed for the domain of } g(x)$$
 ✓

Range $\{y \in R : y \geq -1\}$ ✓

[10]

5. (a) $\vec{FD} = \begin{pmatrix} -3 \\ 3 \\ 2 \end{pmatrix}$ ✓

$$\therefore r = \begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix} \pm \lambda \begin{pmatrix} -3 \\ 3 \\ 2 \end{pmatrix} \text{ OR } \begin{pmatrix} 0 \\ 3 \\ 2 \end{pmatrix} \pm \lambda \begin{pmatrix} -3 \\ 3 \\ 2 \end{pmatrix}$$
 ✓

(b) Use \vec{FD} as the normal vector of the plane ✓

$$\therefore k = \begin{pmatrix} -3 \\ 3 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 3 \\ 2 \end{pmatrix} = -9 + 9 + 4 = 4$$
 ✓

$$\therefore \begin{pmatrix} x \\ y \\ z \end{pmatrix} \cdot \begin{pmatrix} -3 \\ 3 \\ 2 \end{pmatrix} = 4 \rightarrow -3x + 3y + 2z = 4 \quad \checkmark$$

5. (c) $\begin{pmatrix} 3-3\lambda \\ 3\lambda \\ 2\lambda \end{pmatrix} \cdot \begin{pmatrix} -3 \\ 3 \\ 2 \end{pmatrix} = 4 \rightarrow \therefore \lambda = \frac{13}{22}$ ✓✓

$$\therefore \begin{pmatrix} 3 - \frac{39}{22} \\ \frac{39}{22} \\ \frac{26}{22} \end{pmatrix} = \frac{1}{22} \begin{pmatrix} 27 \\ 39 \\ 26 \end{pmatrix} = \frac{27}{22}i + \frac{39}{22}j + \frac{13}{11}k$$
 ✓


(d) Radius $\therefore p = \frac{1}{2}|FD| = \frac{1}{2} \left| \begin{pmatrix} -3 \\ 3 \\ 2 \end{pmatrix} \right| = \frac{\sqrt{22}}{2}$ ✓

Centre $\therefore \vec{OF} + \frac{1}{2}\vec{FD}$ ✓

$$\therefore \begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} -3 \\ 3 \\ 2 \end{pmatrix} = \begin{pmatrix} \frac{3}{2} \\ \frac{3}{2} \\ 1 \end{pmatrix}$$
 ✓

$\therefore a = b = \frac{3}{2}c = 1$ ✓ [12]

6. $y = x + \frac{2}{x+1}$ (after long division or otherwise) ✓




- ✓ Vertical asymptote $x = -1$
- ✓ Oblique asymptote $y = x$
- ✓ y intercept at $(0, 2)$
- ✓ Shape and accuracy

[5]

7. (a) $|z| = \sqrt{3k^2 + k^2} = 2|k| \quad (k \neq 0)$

✓

(b)



$$\arg(z) = \frac{\pi}{6} \quad \text{for } k > 0$$

✓

$$\arg(z) = \frac{-5\pi}{6} \quad \text{for } k < 0$$

✓

(c) $\arg(z) = \frac{-\pi}{3} \quad \text{for } k > 0$

✓

$$\arg(z) = \frac{2\pi}{3} \quad \text{for } k < 0$$

✓

[5]

Calculator-Assumed Solutions

8. For ABC to be collinear,
- $AB \parallel AC \parallel BC$
- (any two)

$$\overrightarrow{AB} = \begin{pmatrix} 4 \\ \alpha - 1 \\ -2 \end{pmatrix} \quad \overrightarrow{AC} = \begin{pmatrix} 8 \\ 2 \\ \beta - 1 \end{pmatrix} \quad \checkmark$$

$$\therefore \begin{pmatrix} 8 \\ 2 \\ \beta - 1 \end{pmatrix} = k \begin{pmatrix} 4 \\ \alpha - 1 \\ -2 \end{pmatrix} \quad \checkmark$$

$$\therefore k = 2 \rightarrow \alpha = 2\beta = -3 \quad \checkmark\checkmark \quad [4]$$

9. (a) $\omega = \frac{1}{8} \rightarrow T = \frac{2\pi}{\omega} = \frac{2\pi}{\frac{1}{8}} = 16\pi = 50.27 \text{ seconds}$ $\checkmark\checkmark$

(b) $\frac{d}{dt} r(t) = \dot{r}(t) = \frac{5}{4} \cos\left(\frac{t}{8}\right) i + \frac{3}{4} \sin\left(\frac{t}{8}\right) j$ $\checkmark\checkmark$

(c) speed $|\dot{r}(t)| = \sqrt{\frac{25}{16} \cos^2\left(\frac{t}{8}\right) + \frac{9}{16} \sin^2\left(\frac{t}{8}\right)}$ $\checkmark\checkmark$


$$|\dot{r}| = \sqrt{\left[\frac{9}{16} \cos^2\left(\frac{t}{8}\right) + \frac{9}{16} \sin^2\left(\frac{t}{8}\right) \right] + \cos^2\left(\frac{t}{8}\right)} = \sqrt{\frac{9}{16} + \cos^2\left(\frac{t}{8}\right)}$$

max speed at $\cos^2\left(\frac{t}{8}\right) = 1$ or $\cos\left(\frac{t}{8}\right) = \pm 1$

$\therefore \text{max speed} = \frac{5}{4}$ for $t = 8n\pi, n = 0, 1, 2, \dots$ \checkmark

$\therefore r(8n\pi) = \pm 6j$ $\checkmark\checkmark$

- (d) Elliptical path with intercepts $(\pm 10, 0)$ and $(0, \pm 6)$ $\checkmark\checkmark$



$$r(0) = -6j \quad \checkmark$$


$$\dot{r}(0) = \frac{5}{4}i \quad \checkmark$$

$$r(6) = 6.82i - 4.39j \quad \checkmark$$

$$\dot{r}(6) = 0.91i + 0.51j \quad \checkmark$$


[15]

10. (a)



- ✓ Mirror image curve over the y axis
- ✓ Location and accuracy

(b)




- ✓ Vertical asymptote $x = -3$
- ✓ Indicates $y \rightarrow |\infty|$ as $|x| \rightarrow -3$
- ✓ Correct domain.
- ✓ crosses $f(x)$ when $y=1$ or -1

(c) from the graph below $x=\pm 5$

✓✓

[8]



11. (a) (i) $\Im(z) \geq -2$ ✓
 $\Im(z) \leq 2\Re(z) + 1$ ✓✓


(i) $2 \leq |z| \leq 4$ ✓
 $\frac{-\pi}{2} \leq \arg(z) \leq \frac{\pi}{10}$ ✓✓

(b) Let $z = x + yi$ and $z \neq 0$

$$\frac{1}{x+yi} - \frac{1}{x-yi} = \frac{-2yi}{x^2+y^2} = i$$
 ✓

$$-2y = x^2 + y^2$$

$$\therefore x^2 + (y+1)^2 = 1$$
 ✓



- ✓ circle centred at (0, -1) with radius = 1
- ✓ discontinuity at the origin

[10]

12. (a) $45 \text{ min} \rightarrow t = \frac{3}{4}$

$$\begin{pmatrix} -6 \\ 12 \\ -0.6 \end{pmatrix} + \frac{3}{4}v = 0$$
 ✓

$$\therefore v = \frac{4}{3} \begin{pmatrix} 6 \\ -12 \\ 0.6 \end{pmatrix} = \begin{pmatrix} 8 \\ -16 \\ 0.8 \end{pmatrix} = 8i - 16j + 0.8k \text{ km/h}$$
 ✓

12. (b) $20 \text{ min} \rightarrow t = \frac{1}{3}$

$$\vec{OA} = O + \frac{1}{3} \begin{pmatrix} 12 \\ -12 \\ 0 \end{pmatrix} = \begin{pmatrix} 4 \\ -4 \\ 0 \end{pmatrix} \quad \checkmark$$

$$\vec{OS} = \begin{pmatrix} -6 \\ 9 \\ -0.6 \end{pmatrix} + \frac{1}{3} \begin{pmatrix} 6 \\ -9 \\ 0.6 \end{pmatrix} = \begin{pmatrix} -4 \\ 6 \\ -0.4 \end{pmatrix} \quad \checkmark$$

$$\therefore \vec{AS} = \begin{pmatrix} -4 \\ 6 \\ -0.4 \end{pmatrix} - \begin{pmatrix} 4 \\ -4 \\ 0 \end{pmatrix} = \begin{pmatrix} -8 \\ 10 \\ -0.4 \end{pmatrix} \quad \checkmark$$

$$|\vec{AS}| = \sqrt{\begin{pmatrix} -8 \\ 10 \\ -0.4 \end{pmatrix}} = 12.81 \text{ km} \quad \checkmark$$

(c) From $t_0 = 1500$:

$$\vec{OT} = \begin{pmatrix} 8 \\ -8 \\ 0 \end{pmatrix} + t \begin{pmatrix} x \\ y \\ z \end{pmatrix} \text{ where } \sqrt{\begin{pmatrix} x \\ y \\ z \end{pmatrix}} = 500$$

Condition for collision:

$$\vec{OT} = \begin{pmatrix} 8 \\ -8 \\ 0 \end{pmatrix} + t \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -2 \\ 3 \\ -0.2 \end{pmatrix} + t \begin{pmatrix} 6 \\ -9 \\ 0.6 \end{pmatrix} = \vec{OS} \quad \checkmark$$

$$\therefore x = 6 - \frac{10}{t} \quad y = \frac{11}{t} - 9 \quad z = 0.6 - \frac{0.2}{t} \quad \checkmark$$

$$\text{and } \left(6 - \frac{10}{t}\right)^2 + \left(\frac{11}{t} - 9\right)^2 + \left(0.6 - \frac{0.2}{t}\right)^2 = 500^2 \quad \checkmark$$

$$\text{CAS: } t = 0.029112 \text{ or } 1.75 \sim 2 \text{ min (1502hrs)} \quad \checkmark$$

$$\text{and } \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -337.50 \\ 368.85 \\ -6.27 \end{pmatrix} \text{ km/h} \quad \checkmark$$

[11]

13. (a) $y = \frac{-1}{2}(x-2)^2 + 8 = \frac{-x^2}{2} + 2x + 6$

$$\therefore a = \frac{-1}{2} \quad b = 2 \quad c = 6 \quad \checkmark \checkmark$$

OR

$$y = \frac{1}{2}(x-2)^2 - 8 = \frac{x^2}{2} - 2x - 6$$


$$\therefore a = \frac{1}{2}b = -2c = -6$$

✓


(BOTH solutions must be given)

13. (b)

✓✓



(c)



From the diagram above:

(i) $n \geq m - 6$

✓✓

OR $n \geq -m - 2$

✓

(both solutions must be given)

(ii) $n < m - 6$

✓


AND $n \leq m - 2$

✓

[10]

14. (a) $(x-1)^2 + (y-1)^2 + z^2 = 2$ ✓
 \therefore centre at $(1,1,0)$ and radius $\sqrt{2}$ ✓✓
- (b) $(x-1)^2 + (y-1)^2 + 1^2 = 2$
 $\therefore (x-1)^2 + (y-1)^2 = 1$
 centre at $(1,1,1)$ and radius 1 ✓✓
- (c) $d = \overrightarrow{CP} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$ ✓
 $r = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$ ✓
 $y=1$ and $x-1=z$ ✓
- (d) $n = d = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$
 $k = n \cdot \overrightarrow{DP} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix} = 2+0+1=3$ ✓
 $\therefore r \cdot \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} = 3$ ✓ [10]

15. (a) See diagram below.



(i) $|z|_{\max} = |\overrightarrow{OC}| + r = 2\sqrt{2} \times \sqrt{2} + 2 = 6$ units

✓✓

(ii) $\sin \phi = \frac{2}{4} \rightarrow \phi = \frac{\pi}{6}$

✓

$$\tan(\phi + \theta) = \frac{2\sqrt{2}}{2\sqrt{2}} = 1 \rightarrow \phi + \theta = \frac{\pi}{4}$$

✓

$$\therefore \theta = \frac{\pi}{4} - \frac{\pi}{6} = \frac{\pi}{12}$$

✓

(b) (i) $\cos(n\theta) = \frac{z^n + z^{-n}}{2}$

✓

$$\sin(n\theta) = \frac{z^n - z^{-n}}{2i}$$

✓

(ii) $\cos^3 \theta = \left(\frac{z^1 + z^{-1}}{2} \right)^3$

✓

$$\therefore \frac{1}{8} [z^3 + 3z^2 z^{-1} + 3z^1 z^{-2} + z^{-3}]$$

✓

$$\therefore \frac{1}{4} \left[\frac{z^3 + z^{-3}}{2} \right] + \frac{3}{4} \left[\frac{z^1 + z^{-1}}{2} \right]$$

✓✓

$$\therefore \frac{1}{4} \cos(3\theta) + \frac{3}{4} \cos \theta \text{ as required}$$

[11]

$$16. \quad (a) \quad f(f) = \frac{f+2}{f-1} = \frac{\frac{x+2}{x-1}+2}{\frac{x+2}{x-1}-1} = x \quad \checkmark \checkmark$$

$\therefore f(x)$ is its own inverse \checkmark

$$(b) \quad f(x) = \frac{x+2}{x-1} = 1 + \frac{3}{x-1}$$

Domain $\{x \in \mathbb{R} : x \neq 1\}$ \checkmark

Range $\{y \in \mathbb{R} : y \neq 1\}$ \checkmark

(c) Since $y=x$ and $y=-x$ are perpendicular, a reflection of $y=x$ does not affect the symmetry of $g(x)$ over $y=-x$ \checkmark
therefore, the reflected function continues to be its own inverse \checkmark

$$(d) \quad f^{-1}(x) = f(x) \rightarrow h(x) = f^{-1}(f(h)) \quad \checkmark$$

$$\therefore \frac{f(h)+2}{f(h)-1} = \frac{\frac{1}{x}+2}{\frac{1}{x}} \quad \checkmark$$

$$\therefore h(x) = 3x+1 \quad \checkmark$$

[10]

$$17. \quad P\left(\frac{1}{w}\right) = \left(\frac{1}{w}\right)^{8n} - \left(\frac{1}{w}\right)^{4n} + 1 = \frac{1}{w^{8n}} - \frac{1}{w^{4n}} + 1 \quad \checkmark$$

$$\therefore \frac{1}{w^{8n}}(1 - w^{4n} + w^{8n}) = \frac{1}{w^{8n}} \times P(w) \quad \checkmark$$

$$\frac{1}{w^{8n}} \times 0 = 0 \quad \therefore \frac{1}{w} \text{ is also a root of } P(w) \quad \checkmark$$

$$P(iw) = (iw)^{8n} - (iw)^{4n} + 1 = i^{8n} w^{8n} - i^{4n} w^{4n} + 1 \quad \checkmark$$

$$\therefore 1 \times w^{8n} - 1 \times w^{4n} + 1 = P(w) = 0 \quad \checkmark$$

$\therefore iw$ is also a root of $P(w)$ \checkmark [6]

$$18. \quad \cos \theta = \frac{x}{2} \quad \checkmark$$

$$y = \cos 2\theta = 2 \cos^2 \theta - 1 \quad \checkmark$$

$$\therefore y = 2 \left(\frac{x}{2}\right)^2 - 1 = \frac{x^2}{2} - 1 \quad \checkmark$$

$$x = 2 \cos \theta, \therefore -2 \leq x \leq 2$$

$$\text{Domain } \{x \in \mathbb{R} : -2 \leq x \leq 2\}$$

✓

$$y = \cos(2\theta) \text{ Range } \{y \in \mathbb{R} : -1 \leq y \leq 1\}$$

✓

[5]