

No other items may be taken into the examination room. It is **your responsibility** to ensure that you do not have any unauthorised material if you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

© 2018 WA Exam Papers. Kennedy Baptist College has a non-exclusive licence to copy and communicate this document for non-commercial educational use within the school. No other copying, communication or use is permitted without the express written permission of WA Exam Papers. NS245-125-4.

Important note to candidates

Special items: drawing instruments, templates, notes on two unfolded sheets of A4 paper, and up to three calculators approved for use in this examination

Standard items: pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

Formula sheet (referred from Section One)

This Question/Answer booklet

Materials required/recommended for this section

Time allowed for this section
Reading time before commencing work: ten minutes
Working time: one hundred minutes

Your name _____
In words _____

--	--	--	--	--	--	--	--

Student number: In figures _____

Calculator-assumed
Section Two:
UNITS 3 AND 4
METHODS
MATHEMATICS

Question/Answer booklet

Semester Two Examination, 2018

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	8	8	50	52	35
Section Two: Calculator-assumed	13	13	100	98	65
Total					100

Supplementary page

Question number: _____

Instructions to candidates

1. The rules for the conduct of examinations are detailed in the school handbook. Sitting this examination implies that you agree to abide by these rules.
2. Write your answers in this Question/Answer booklet.
3. You must be careful to confine your response to the specific question asked and to follow any instructions that are specified to a particular question.
4. Supplementary pages for the use of planning/continuing your answer to a question have been provided at the end of this Question/Answer booklet. If you use these pages to continue an answer, indicate at the original answer where the answer is continued, i.e. give the page number.
5. Show all your working clearly. Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
6. It is recommended that you do not use pencil, except in diagrams.
7. The Formula sheet is not to be handed in with your Question/Answer booklet.

Supplementary page

Section Two: Calculator-assumed

Methods Units 3 and 4

Calculator-Assumed

Working time: 100 minutes.

Question 9

(a) Assuming no further contamination occurred, determine the level of Strontium-90 in a contaminated soil sample at the start of 1995 was 0.55 mg/kg.

The level of Strontium-90 has a half-life of 28.8 years and decays continuously such that $S = S_0 e^{-kt}$, where S is the level of Strontium-90, t is the time in years since the level was S_0 and k is a constant.

(i) the level of Strontium-90 in the sample at the start of 2018.

(3 marks)

<input checked="" type="checkbox"/> Assumptions	Assume the sample contained 0.55 mg/kg of Strontium-90 at the start of 1995.
<input checked="" type="checkbox"/> Specific behaviours	Strontium-90 has a half-life of 28.8 years and decays continuously.
<input checked="" type="checkbox"/> Solution	$S(23) = 0.55 e^{-0.02407 \times 23} = 0.316 \text{ mg/kg}$
<input checked="" type="checkbox"/> Working	Value for S that rounds to 0.32

(ii) the rate of change of the level of Strontium-90 in the sample at the start of 2018.

(1 mark)

<input checked="" type="checkbox"/> Assumptions	Assume the sample contained 0.55 mg/kg of Strontium-90 at the start of 1995.
<input checked="" type="checkbox"/> Specific behaviours	Strontium-90 has a half-life of 28.8 years and decays continuously.
<input checked="" type="checkbox"/> Solution	$-0.02407 \times 0.316 = -0.0076 \text{ mg/kg/year}$
<input checked="" type="checkbox"/> Working	Answer (i) multiplied by k

(b) Strontium-90 decays into yttrium-90. The mass of yttrium-90 decays continuously such that $Y = Y_0 e^{-0.010t}$, where Y is the mass of Yttrium-90 and t is the time in hours since the level was Y_0 . Determine the time taken for a mass of Yttrium-90 to decrease by 90%.

(2 marks)

<input checked="" type="checkbox"/> Assumptions	Assume the sample contained 0.10 mg of Strontium-90 at the start of 1995.
<input checked="" type="checkbox"/> Specific behaviours	Strontium-90 decays into Yttrium-90 with a half-life of 28.8 years.
<input checked="" type="checkbox"/> Solution	$e^{-0.010t} = 0.10$ $t = 228 \text{ hours}$
<input checked="" type="checkbox"/> Working	Solves for t Writes equation for t

Question number: _____
 Supplementary page _____

Question 10

(6 marks)

A local council wants to know what proportion of its ratepayers support a recent decision to start charging for parking at its 15 car parks.

- (a) Comment, with reasons, on whether the following sampling methods are likely to introduce bias.

- (i) Send a council worker to one randomly selected council car park at 10 am on a Monday morning and get them to record the responses of the first 20 drivers who arrive. (2 marks)

Solution	
Biased, as	
- small sample size	
- only ask users of car park chosen	
- car parkers may not be ratepayers, etc, etc	
Specific behaviours	
✓ indicates bias, with reason	
❑ second reason	

- (ii) In a council newsletter sent to all ratepayers, include a link to a public page on the council website where users can click a 'yes' or 'no' button to register their support. (2 marks)

Solution	
Biased, as	
- volunteer sampling	
- ratepayers may not have internet access	
- web site visitors may not be ratepayers, etc, etc	
Specific behaviours	
✓ indicates bias, with reason	
❑ second reason	

- (b) Following the analysis of a large random sample, a student calculated the 95% confidence interval for ratepayer statements below as **true** or **false**, where the statement was taken logically from the council's report.

- (i) There is a 95 % chance that the true proportion of ratepayers lies between 0.1 and 0.3.

Solution	
False. (See notes)	
Specific behaviours	
✓ correct response	

- (ii) If the random sampling was repeated many times, 95% of the time the true proportion of supportive ratepayers would lie between 0.1 and 0.3.

Solution	
False. (See notes)	
Specific behaviours	
✓ correct response	

See next page

SN245-125-4

Question 21

(6 marks)

A game is played at a carnival where two fair 4-sided dice with faces numbered 1, 2, 3 and 4 are tossed at the same time. Patrons pay \$3 for each play of the game, winning a major prize if both dice show a four or a minor prize if just one of the dice shows a four. The operator of the game buys major prizes for \$22 each, minor prizes for \$2.50 and must pay overhead costs of \$95 per day.

Determine how many times the game must be played per day so that the operator can expect to make a daily profit of at least \$150.

Solution			
$X = \$\text{profit per game for operator}$			
x	-19.00	0.50	3.00
$P(X=x)$	$\frac{1}{16}$	$\frac{6}{16}$	$\frac{9}{16}$

$$E(X) = \frac{-19 + 3 + 27}{16} = \frac{11}{16} = \$0.6875$$

$$\frac{11}{16}n \geq 95 + 150 \\ n \geq 356.4$$

Require at least 357 patrons to play per day.

Specific behaviours	
✓ defines random variable	
❑ table with row showing values RV can take	
❑ correct probabilities for all outcomes	
❑ calculates $E(X)$	
❑ forms inequality $n \times E(X) \geq \text{overheads+profit}$	

Examiners note	
Interpretation of Confidence Intervals	
Suppose that a 95% confidence interval is calculated as $[a, b]$.	
A common misconception is to think this means there is a 95% chance that the true population proportion falls between a and b . This is incorrect. Like any population parameter, the population proportion is a constant, not a random variable. It does not change. The probability that a constant falls within any given range is either 0 or 1.	
The confidence level describes the uncertainty associated with a sampling method. Suppose we used the same sampling method to select different samples and to compute a different interval estimate for each sample. Some interval estimates would include the true population proportion, and some would not. A 95% confidence level means that we would expect 95% of the interval estimates to include the true population proportion.	

rted that
ch of the
not follow

lies
(1 mark)

ie the true
(1 mark)

End of questions

SN245-125-4

Question 11

The discrete random variable X has $E(X) = 3.2$ and probability function


$$P(X=x) = \begin{cases} a+bx & x=2,3,4 \\ 0 & \text{elsewhere.} \end{cases}$$

(8 marks)

(2 marks)

Solution					
Sum of probabilities: $a+2b+3b+a+4b=1$					
Expected value: $2(a+2b)+3(a+3b)+4(a+4b)=3.2$					
Sum of probabilities: $a+2b+3b+a+4b=1$					
Another function A is defined on the interval $-3 \leq x \leq 5$ by					
$A(x) = \int_x^{-3} f(t) dt.$					

It is known that $A(-1) = A(5) = -9$ and $A(3) = 0$. Sketch the graph of $y = A(x)$ on the axes below, clearly indicating the location of all x -intercepts, turning points, points of inflection and other key features.




The graph of $y = f(x)$ is shown below.

Another function A is defined on the interval $-3 \leq x \leq 5$ by

$$A(x) = \int_x^{-3} f(t) dt.$$

Solution	
Sketched over defined interval	
x-intercepts	
local minimum	
labelled endpoint	
approx-position pt. of inflection	



(c) A second random variable Y is a linear transformation of X such that $Y = kX + 4$, where k is a constant and $E(Y) = 20$. Determine $\text{Var}(Y)$.

(2 marks)

Solution	
Correct Variance	
indicates probabilities	
specific behaviours	
$\text{Var}(X) = \frac{75}{47} = 0.626$	
P($X=x$)	
x	2
	3
	4
	13
	30
	30
	10
	30
	7
Solution	

(2 marks)

Solution	
Correct Variance	
indicates probabilities	
specific behaviours	
$\text{Var}(Y) = 5^2 \times \frac{75}{47} = 15.6$	
3.2k + 4 = 20 $\Rightarrow k = 5$	
Solution	

(2 marks)


Solution	
Correct Variance	
indicates probabilities	
specific behaviours	
$\text{Var}(X) = \frac{75}{47} = 0.626$	
P($X=x$)	
x	2
	3
	4
	13
	30
	30
	10
	30
	7
Solution	

(2 marks)

(8 marks)

Question 12

- (a) The graphs of the probability density functions of three normally distributed random variables W , X and Y are shown below.



State, with justification, which of the three random variables has

- (i) the largest mean?

Solution	
Y - maximum furthest to right	
Specific behaviours	
✓ correct variable	

(1 mark)

- (ii) the smallest standard deviation?

Solution	
X - highest f_{max} , so least spread.	
Specific behaviours	
✓ correct variable	

(1 mark)

- (b) Empty bottles are filled with A mL of water, where A is a normally distributed random variable with mean of 380 mL and standard deviation of 4.5 mL.

- (i) Determine the probability that a bottle is filled with less than 373 mL. (1 mark)

Solution	
$P[X < 373] = 0.0599$	
Specific behaviours	
✓ correct probability	

- (ii) Determine the probability that a bottle is filled with more than 375 mL, given that it is filled with less than 380 mL. (2 marks)

Solution	
$P[375 < A < 380] = 0.3667$	
$p = \frac{0.3667}{0.5} = 0.7335$	
Specific behaviours	
✓ numerator	
✗ correct probability	


- (iii) The mean of A is to be increased by k mL so that 99% of all bottles are filled with at least 375 mL. Determine the value of k . (3 marks)

Solution	
$\frac{375 - \bar{x}}{4.5} = -2.326 \Rightarrow \bar{x} = 385.47$	
$k = 385.47 - 380 = 5.47$ mL	
Specific behaviours	
✓ equation showing correct z-score	
✗ solves for mean	
✗ correct value of k	

See next page

SN245-125-4

- (b) The graph of $x=2y^2-12$ and the line $4y=x+6$ are shown below.



Determine the area bounded by the line and the curve. (4 marks)

Solution	
Line-curve intersect when $x=-10, 6$ (CAS)	
When $y=0, x=-12$.	
Curve: $y = \pm\sqrt{0.5x+6}$	
Line: $y = 0.25x + 1.5$	

$$A_1 = \int_{-10}^6 \left((\sqrt{0.5x+6}) - (0.25x+1.5) \right) dx \quad \text{using } \frac{56}{3}$$

$$A_2 = \int_{-12}^{-10} \left((\sqrt{0.5x+6}) - (-\sqrt{0.5x+6}) \right) dx \quad \text{using } \frac{8}{3}$$

$$A = A_1 + A_2 \quad \text{using } \frac{64}{3}$$

Specific behaviours	
✗ points of intersection	
✗ correct integrand	
✗ correct bounds	
✗ correct area	

Alternative Solution	
Line and curve intersect when $y=-1, 3$ (CAS)	
Line: $x = 4y - 6$	
$A = \int_{-1}^3 ((4y - 6) - (2y^2 - 12)) dy$	
$\text{using } \frac{64}{3}$ sq units	

Specific behaviours	
✗ points of intersection	
✗ correct integral A_1	
✗ correct integral A_2	
✗ correct area	

See next page

SN245-125-4

Question 13

(a) If there were 36 000 people living in the city, estimate the actual number of these who had visited a doctor in the last year. (2 marks)

Solution	
Specific behaviours	
✓ indicates proportion	
✗ estimate to nearest 100	
0.19165 × 36000 ≈ 6 900 people	
Specific behaviours	
✓ indicates margin of error	
$E = 0.011487 \times 2.576 = 0.0296$	
Solution	
Determine the approximate margin of error for the 99 % confidence interval for the proportion of people who had visited a doctor in the last year.	

Solution	
Specific behaviours	
✓ indicates standard deviation	
✗ correct margin of error	
$E = 0.011487 \times 2.576 = 0.0296$	
Solution	
Determine the approximate margin of error for the 99 % confidence interval for the proportion of people who had visited a doctor in the last year. (2 marks)	


Solution	
Specific behaviours	
✓ indicates standard deviation	
✗ correct margin of error	
$sd = \sqrt{0.19165(1-0.19165)} = 0.011487$	
Solution	
Determine an approximate 99 % confidence interval for the true proportion of people who had visited a doctor in the last year. (2 marks)	

Solution	
Specific behaviours	
✓ indicates $p \pm E$	
✗ correct interval	
$[0.1621, 0.2212]$	
Solution	
Determine an approximate 99 % confidence interval for the true proportion of people who had visited a doctor in the last year. (2 marks)	

Solution	
Specific behaviours	
✓ indicates correct method	
✗ indicates size	
$n = \frac{2.576^2 \cdot 0.19165(1-0.19165)}{0.055^2} \approx 340$	
Solution	
In order to confirm the sample proportion obtained from the random sample, another sample is to be taken. Estimate, to the nearest 10 people, the sample size required to obtain an approximate margin of error for a 99 % confidence interval that is close to 0.055. (2 marks)	

Solution	
Specific behaviours	
✓ indicates correct method	
✗ indicates size	
$n = \frac{2.576^2 \cdot 0.19165(1-0.19165)}{0.055^2} \approx 340$	
Solution	
In order to confirm the sample proportion obtained from the random sample, another sample is to be taken. Estimate, to the nearest 10 people, the sample size required to obtain an approximate margin of error for a 99 % confidence interval that is close to 0.055. (2 marks)	

(a) The graph of $y = f(x)$ is shown together with some values of $f(x)$.
225 out of a random sample of 1174 people in a city had visited a doctor in the last year.
Estimate the actual number of these who had visited a doctor in the last year. (8 marks)



(a) The graph of $y = f(x)$ is shown together with some values of $f(x)$.

Solution	
Specific behaviours	
✓ over estimate of area	
✗ under estimate of area	
✗ averages for best estimate of area	
✗ correct sign for integral	
Solution	
$\therefore \int_1^3 f(x) dx \approx 8.4$	
Area estimate = $(8.85 + 7.95) \div 2 = 8.4$	
Under estimate = $0.25(9.5 + 8.5 + 7.2 + 6.6) \approx 7.95$	
Over estimate = $0.25(10.2 + 9.5 + 8.5 + 7.2) \approx 8.85$	
Solution	
Calculate a numerical approximation for $\int_1^3 f(x) dx$.	
By considering the areas of the rectangles shown and using values of $f(x)$ from the table,	
calculate a numerical approximation for $\int_1^3 f(x) dx$. (4 marks)	

x	$f(x)$	x	$f(x)$	x	$f(x)$	x	$f(x)$
0.75	-9.3	-10.2	-9.5	-1.25	-8.5	1.5	-7.2
1	-1.5	2	-6.6	2.25	-5.9		
1.25	1.75	1.75	2	2.25			
1.5	2	2.25					

(7 marks)

Question 14

The table below shows the sign of the polynomial $f(x)$ and some of its derivatives at various values of x . There are no other zeroes of $f(x)$, $f'(x)$ or $f''(x)$ apart from those shown in the table.

x	-2	-1	0	1	2	3	4
$f(x)$	+ <i>i</i>	0	- <i>i</i>	- <i>i</i>	- <i>i</i>	0	+ <i>i</i>
$f'(x)$	- <i>i</i>	- <i>i</i>	0	+ <i>i</i>	+ <i>i</i>	0	+ <i>i</i>
$f''(x)$	+ <i>i</i>	+ <i>i</i>	+ <i>i</i>	0	- <i>i</i>	0	+ <i>i</i>

- (a) For what value(s) of x is the graph of the function concave up? (1 mark)

Solution
$x < 1$ and $x > 3$


Specific behaviours
✓ correct inequalities and domain

- (b) At what location does the graph of f have a turning point? Explain your answer. (2 marks)

Solution
At $x=0$. The gradient is zero and f is concave up on either side.

Specific behaviours
✓ location ☒ explanation

- (c) Sketch a possible graph of $y=f(x)$ on the axes below. (4 marks)



Solution
See graph

Specific behaviours
✓ correct location of roots ☒ correct location of stationary points ☒ point of inflection at $x=1$ ☒ concave up everywhere except $1 < x < 3$

- (c) Determine the mean time to process an order in minutes and seconds. (4 marks)

Solution
$g(t) = \frac{-1}{48}(t-12), 4 < t \leq 12$ $E(T) = \int_0^4 t \left(\frac{t}{24}\right) dt + \int_4^{12} t \left(\frac{-1}{48}(t-12)\right) dt$ $\frac{16}{3} = 5\frac{1}{3}$ Mean is 5 min 20 sec.

Specific behaviours

- ✓ indicates $g(t)$ for second interval
- ☒ indicates both integrals
- ☒ evaluates mean
- ☒ writes mean as required

The variance of T is 6 minutes 13 seconds.

- (d) Two new procedures will affect the processing time of an order. The first will decrease the time by 15% and the second will then add one-and-a-half minutes. Determine the new mean and standard deviation of the time to process an order. (3 marks)

Solution
$E(0.85T + 1.5) = 0.85 \times 5\frac{1}{3} + 1.5$ $\textcolor{brown}{i} 6.03 \text{ min (6 m 2 s)}$ $\sigma_{old} = \sqrt{6\frac{13}{60}} = 2.493$ $\sigma_{new} = 0.85 \times 2.493 \textcolor{brown}{i} 2.12 \text{ min (2 m 7 s)}$

Specific behaviours

- ☒ new mean
- ✓ indicates original sd
- ☒ new sd

Every day a scientific researcher randomly catches 8 fish from an inland lake containing a large number of fish, 63% of which are thought to be trout.

(a) The random variable X is the number of trout in the daily catch.

(i) Describe the distribution of X .

$X \sim B(8, 0.63)$	Solution
✓ indicates binomial	✓ indicates parameters
✓ correct value	✓ correct number of days
Days.	

(ii) Over a period of 14 days, how many times would you expect the daily catch to contain more trout than fish of other species?

$P(X \geq 5) = 0.6626$	$P(X \leq 5) = 0.6626$
$n = 0.6626 \times 14 \approx 9$ days	$P(Y = 15) = 0.0058$
✓ indicates method	✓ correct probability
Solution	

(iii) Determine the probability that a total of 15 trout are caught over two consecutive days.

$p = P(X=7) \times P(X=8) \times 2$	$P(Y=15) = 0.0058$
$? 0.1166 \times 0.0248 \times 2 = 0.0058$	$P(Y=15) = 0.0058$
✓ correct probability	✓ indicates method
Solution	

(b) Determine the probability that a total of 15 trout are caught over two consecutive days.

(i) Calculate an approximate 90% confidence interval for the proportion of trout in the lake given that over a 10-day period, a total of 48 trout were caught.

$x = 48, n = 80, p = 0.6$	Solution
✓ states interval	✓ indicates x and n
Solution	

(ii) Use the confidence interval to comment on the researcher's suspicion. (2 marks)


No evidence that proportion is lower, as 63% is within the CI, but there is evidence that proportion is more than 50%, as 50% is below the lower bound of CI.	Solution
✓ comment on lower than 63%	✓ comment on more than 50%
Solution	

(c) Calculate an approximate 90% confidence interval for the proportion of trout in the lake given that over a 10-day period, a total of 48 trout were caught.

(i) Use the confidence interval to comment on the researcher's suspicion. (2 marks)

No evidence that proportion is lower, as 63% is within the CI, but there is evidence that proportion is more than 50%, as 50% is below the lower bound of CI.	Solution
✓ comment on lower than 63%	✓ comment on more than 50%
Solution	

The time T to process orders at a warehouse is random variable which can take any value in the interval 0 to 12 minutes. The graph of the triangular probability density function of T is shown below.



The time T to process orders at a warehouse is random variable which can take any value in the interval 0 to 12 minutes. The graph of the triangular probability density function of T is shown below.

(a) Determine the value of k .

$f(t) = \frac{t}{24}, 0 \leq t \leq 4$	Solution
$\int_0^4 \frac{t}{24} dt = \frac{3}{16}$	✓ indicates integral
$P(T < 3) = \int_0^3 \frac{t}{24} dt = \frac{3}{16} = 0.1875$	✓ correct probability
Solution	

(b) Determine the probability that the time to process an order takes less than 3 minutes.

Question 16

(7 marks)

At time $t=0$, a small body P is at the origin O and is moving with a velocity of 18 ms^{-1} . The acceleration of P for $t \geq 0$ is given by

$$a = \frac{-3}{\sqrt{t+4}} \text{ ms}^{-2}.$$

- (a) Determine the velocity of P when $t=5$.

(4 marks)

Solution	
$v = \int a dt \quad -6\sqrt{t+4} + c$	$v(5) = 18 + \int_0^5 \left(\frac{-3}{\sqrt{t+4}} \right) dt$
$c = 18 + 6\sqrt{4} = 30$	$\cancel{18} - 12 = 6$
$v = 30 - 6\sqrt{t+4}$	
$v(5) = 30 - 6\sqrt{9} = 12 \text{ ms}^{-1}$	NB Using net change is quicker in (a), but since an expression for $v(t)$ is needed in (b) best to determine it here.
Specific behaviours	
✓ indicates v is integral of a	
✗ correct integral	
✗ evaluates c	
✗ correct velocity	

- (b) Determine the distance of P from O at the instant P is stationary.

(3 marks)

Solution	
$v = 0 \Rightarrow 30 - 6\sqrt{t+4} = 0 \Rightarrow t = 21$	
$OP = \int_0^{21} (30 - 6\sqrt{t+4}) dt \approx 162 \text{ m}$	
Specific behaviours	
✓ determines value of t	
✗ writes integral for change in displacement	
✗ correct distance	

Question 17

(8 marks)

A student repeatedly took random samples of size 150 from a large population in which it was known that 38% of people were classified as overweight. For each sample, the proportion of overweight people was calculated and recorded as the sample proportion.

- (a) Use an appropriate binomial distribution to determine the probability that the sample proportion is no more than 0.34 in a randomly chosen sample. (3 marks)

Solution	
$X \sim B(150, 0.38)$	
$0.34 \times 150 = 51$	
$P(X \leq 51) = 0.1777$	
Specific behaviours	
✓ states parameters	
✗ indicates most number of successes	
✗ correct probability	

- (b) After recording a large number of sample proportions, the student used them to create a histogram from which the approximate normality of their distribution was evident.

- (i) Determine the expected mean and standard deviation of the observed normal distribution. (2 marks)

Solution	
mean = 0.38	
$sd = \sqrt{\frac{0.38(1-0.38)}{150}} \approx 0.0396$	
Specific behaviours	
✓ correct mean	
✗ correct sd	

- (ii) Use this normal distribution to determine the probability that the sample proportion is no more than 0.34 in a randomly chosen sample. (1 mark)

Solution	
$P(X < 0.34) = 0.1564$	
Specific behaviours	
✓ correct probability	

- (iii) Describe how the parameters calculated in (i) would change if the student took smaller random samples. (2 marks)

Solution	
Mean would stay the same.	
SD would increase.	
Specific behaviours	
✓ states no change in mean	
✗ states increase in sd	