


Copyright for test papers and marking guides remains with *West Australian Test Papers*.  
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.  
Test papers should be withdrawn after use and stored securely in the school until 14<sup>th</sup> October 2021.



# **SEMESTER TWO**

## **MATHEMATICS METHODS UNITS 3 & 4**

**2021**

## **SOLUTIONS**

**Calculator-free Solutions**

1. (a)  $(2\pi, 0)$

$$\frac{dy}{dx} = \frac{2\tan(x)}{x} + \frac{2\ln x}{\cos^2(x)}$$

✓

When  $x = 2\pi$ ,  $m = 0 + 2\ln 2\pi$

✓

$$0 = 2\ln 2\pi(2\pi) + c \therefore c = -4\pi \ln 2\pi$$

$$y = (2\ln 2\pi)x - 4\pi \ln 2\pi$$

✓

(b)  $2t \ln(t^4) \tan(t^2)$

✓✓

[5]

$$\frac{1+2+3+4+5}{k} = 1$$

2. (a)

$$k = 15$$

✓

$$\frac{12}{15} = \frac{4}{5}$$

(b)

✓

$$\frac{1+4+9+16+25}{15} = \frac{55}{15} = \frac{11}{3}$$

(c)

✓

(d) (i) Expected value  $\left(\frac{1}{10}\right)\left(\frac{11}{3}\right) + 2 = \frac{71}{30}$

✓

$$\frac{1+8+27+64+125}{15} - \frac{121}{9} = \frac{14}{9}$$

Variance of Y:

$$\frac{\sqrt{14}}{3}$$

Standard deviation of Y =

✓

$$\frac{\sqrt{14}}{30}$$

Standard deviation of X =

✓

$$P\left(X \leq \frac{11}{5}\right) = P(Y \leq 2) = \frac{3}{15}$$

(ii)

✓✓

[8]


3. (a)  $g''(x) = -2x + 1$  when  $x > \frac{1}{2}$   $g(x)$  is concave down

✓✓

(b) Maximum

✓

(c)



	Correct point of inflection	✓	
	Shape showing local minimum and maximum	✓	[5]
4.	(a) $-1 = \log_c(2)$		
	$c^{-1} = 2 \therefore c = \frac{1}{2}$	✓	
	$b = -1$	✓	
	$0 = \log_c x \therefore x = 1 \rightarrow C(1, 0)$		
	$p(x) = 2a^x - 1$ given $(1, 0)$		
	$2a - 1 = 0 \therefore a = \frac{1}{2}$	✓	
	$= 3\log_5 5 + \log_5 2 - \frac{1}{3}(3)\log_5 2$	✓	
	$= 3$	✓	
	(c) $\log_4(x^2 - 6x) = 2$	✓	
	$x^2 - 6x - 16 = 0$		
	$(x - 8)(x + 2) = 0$		
	$\therefore x = 8$	✓	[7]
5.	(a) 0.16	✓	
	(b) 0.34	✓	
	$\frac{0.16}{0.5} = 0.32$		
	(c)	✓✓	[4]
6.	$\log_2 y = \frac{1}{4}\log_2 x + 3$		
	$\log_2 y = \frac{1}{4}\log_2 x + 3\log_2 2$	✓	
	$\log_2 y = \log_2 8x^{0.25}$		
	$y = 8x^{\frac{1}{4}}$	✓	
	$k = 8 \quad n = \frac{1}{4}$	✓	[3]

$$7. \quad m'(x) = \frac{(x^2 - 1)(2xe^{x^2-1}) - (e^{x^2-1})(2x)}{(x^2 - 1)^2}$$

$$= \frac{2xe^{x^2-1}(x^2 - 2)}{(x^2 - 1)^2}$$

Stationary points occur when  $2xe^{x^2-1}(x^2 - 2) = 0$

$$\therefore x = 0, \pm\sqrt{2}$$

$$\therefore \left(0, -\frac{1}{e}\right) \quad (\sqrt{2}, e) \quad (-\sqrt{2}, e)$$

[4]

$$8. \quad (a) \quad \rho(x) = \begin{cases} \frac{1}{20} & 0 \leq x \leq 20 \\ 0 & x > 20 \end{cases}$$

$$\frac{\frac{2}{20}}{\frac{17}{20}} = \frac{2}{17}$$

(b)

(c) 10 minutes

[4]

$$9. \quad \int_{-1}^1 k(1 - x^2) - 2k(x^2 - 1) \, dx = 3k \int_{-1}^1 (1 - x^2) \, dx$$

$$3k \left[ x - \frac{x^3}{3} \right]_{-1}^1 = 8$$

$$4k = 8$$

$$k = 2$$


[3]

$$10. \quad (a) \quad 0 < x \leq 1$$

$$(b) \quad \left(\frac{4}{3}\right)^p = \frac{16}{9}$$

$$p = 2$$

(c) (i)



✓

$$\log_{\frac{4}{3}} x = \frac{\ln x}{\ln \frac{4}{3}}$$

10. (c) (ii)

$$f'(x) = \frac{1}{x \ln \frac{4}{3}} = \frac{1}{x(\ln 4 - \ln 3)}$$

✓✓

[7]

**Calculator-Assumed Solutions**

11. (a)  $M_P(t) = M_R(t) - M_c(t) = -0.4t + 8 - 0.3t - 2$

✓

Maximum profit occurs when  $M_P(t) = -0.7t + 6 = 0$   
therefore  $t = 8.57$  years

✓

(b)  $\int_0^{8.57} -0.7t + 6 \, dt = 25.714285$

✓

Maximum profit = \$25714285 - \$12000000 = \$13 714 285

✓

[4]

12. (a) Discrete data; Independent events

✓

Only two outcomes: under the limit or over the limit.

✓

(b) (i)  $X \sim \text{Bin}(10, 0.05)$   $Y \sim \text{Bin}(10, 0.013)$

✓

$$P(X = 3) + P(X = 2)P(Y = 1) + P(X = 1)P(Y = 2) + P(Y = 3) \\ (0.01048) + (0.07463)(0.11556) + (0.31512)(0.00685) + (0.00024) \\ = 0.02150$$

✓

(ii)  $P(1 \leq Y \leq 10) = 0.12265$

✓

(iii)  $X \sim \text{Bin}(275, 0.05)$   $E(X) = 13.75$   
 $\therefore 13$  drivers.

✓

Variance =  $275 \times 0.05 \times 0.95 = 13.0625$

Standard deviation = 3.6142

✓

(iv)  $Y \sim \text{Bin}(n, 0.013)$   $P(Y \geq 1) \leq 0.6$

$$1 - P(X = 0) \leq 0.6$$

✓

$$P(Y = 0) \geq 0.4$$

$$0.987^n \geq 0.4 \quad \therefore n \leq 70.025$$

✓

Therefore the maximum value of  $n = 70$

✓

[11]

13. Point of intersection:

$$x^3 - 4x^2 + 3x + 1 = x^2 - 3x + 1$$

$$x = 0 \text{ or } 2 \text{ or } 3$$

Area of shaded part:

$$\int_0^3 |(x^3 - 4x^2 + 3x + 1) - (x^2 - 3x + 1)| \, dx = \frac{37}{12} \text{ units}^2 \quad \checkmark \checkmark \text{ Alternate}$$

Shaded area under line:

$$\int_0^2 1 - x - (x^2 - 3x + 1) \, dx = \frac{4}{3} \text{ units}^2 \quad \checkmark$$

Fraction:  $\frac{16}{37}$   $\checkmark$  [4]

14. (a)  $AC = t - 30 \quad \therefore p^2 = 30^2 - (t - 30)^2$   $\checkmark$   
 $p^2 = -t^2 + 60t$   $\checkmark$

Volume of cone =  $\frac{1}{3} \pi r^2 h$

$$V(t) = \frac{1}{3} \pi (-t^2 + 60t)(t) = -\frac{\pi t^3}{3} + 20\pi t^2 \quad \checkmark$$

(b)  $V'(t) = 40\pi t - \pi t^2 = 0$   $\checkmark$   
 $t = 0 \text{ or } 40$   $\checkmark$   
 $V''(t) = 40\pi - 2\pi t$   $\checkmark$   
 $V''(40) = -40\pi \therefore \text{maximum at } t = 40 \text{ cm}$   $\checkmark$

(c) Volume of cone at  $t = 40 \text{ cm}$  is  $\frac{32000\pi}{3} = 33510.322 \text{ cm}^3$   $\checkmark$   
 Volume of sphere =  $113\,097.336 \text{ cm}^3$   $\checkmark$   
 Percentage = 29.63%  $\checkmark$

(d)  $\frac{\delta r}{r} = -0.015 \quad \frac{dV}{dr} = 4\pi r^2 \quad \frac{\delta V}{\delta r} \approx \frac{dV}{dr}$   $\checkmark$   
 $\frac{\delta V}{V} \approx \frac{4\pi r^2}{\frac{4}{3}\pi r^3} \times \delta r$   $\checkmark$

$$\frac{\delta V}{V} \approx \frac{3\delta r}{r}$$

$$\frac{\delta V}{V} \approx 3(-0.015)$$

Approximately 4.5% decrease in volume.  $\checkmark$  [11]

15. (a)  $50 = 100(1 - e^{3k})$   
 $k = -0.23105$  ✓
- (b) Less than 5 minutes:  $P = 100(1 - e^{5 \times -0.23105}) = 68.5\%$  ✓  
 Percentage 5 minutes or longer = 31.5% ✓
- (c) (i)  $t$  is a continuous random variable where  $0 \leq t \leq 30$ .  
 The function in the domain  $0 \leq t \leq 30$  is positive.  
 The probability at  $t = 0$  is 0 and  $t = 30$  is 1,  
 therefore is cumulative. ✓✓
- (ii)  $P(t) = 1 - e^{-0.23105(15)} = 0.96896$  ✓
- (iii)  $\frac{0.68502}{0.90079} = 0.7605$  ✓✓ [8]

16.  $T(t) = 21e^{-kt} + 4$
- $9.8 = 21e^{-kt} + 4$  ✓
- $6.5 = 21e^{-k(t+15)} + 4$  ✓
- $k = 0.056104 \quad t = 22.9334$  ✓
- The liquid was placed in the fridge at 11:37 am ✓ [4]

17. (a) 6 year old tree is growing at a rate of 17.69 cm/year ✓  
 50 year old tree is growing at a rate of 2.277 cm/year ✓  
 7.8 times faster ✓
- (b) (i) Convenience sample: this sample may not be  
 representative of all the six year old trees in the plantation.  
 The sample is not large enough. ✓✓
- (ii) Stratified sample where a number of trees from each  
 area according to the size of the area are chosen at random.  
 Or  
 Systematic and Array sample: Every  $n$ th tree starting  
 at a randomly assigned tree as one walks down each row. ✓✓
- (c) (i)  $P(Z > z) = 0.0094 \quad \therefore z = 2.3495$  ✓
- $\frac{326 - \mu}{12} = 2.3495$  ✓
- $\mu = 297.8 \text{ cm}$  ✓
- (ii) According to the model, the height of a six year old tree  
 is 297.8 cm and the mean of the sample is the same,  
 therefore the model is suitable. ✓
- (iii)  $X \sim (5, 0.0094) \quad P(X = 0) = 0.95388$  ✓
- (d)  $\frac{132}{400} \pm 2.58 \sqrt{\frac{\left(\frac{132}{400}\right)\left(\frac{268}{400}\right)}{400}} = (0.269, 0.391)$  ✓✓

With a 99% confidence level, between 26.9% and 39.1% of the

- mature trees can be used for luxury furniture. ✓ [14]
18. Over - estimate =  $0.2(0.127+0.172+0.184+0.184+0.181) = 0.1696$  ✓  
 Under - estimate =  $0.2(0+0.127+0.172+0.181+0.173) = 0.1306$  ✓  
 Average =  $0.1501 \text{ units}^2$  ✓ [3]
19. (a) (i)  $n > 30$   
 $np = 300 \times 0.12 = 36$   $npq = 300 \times 0.12 \times 0.88 = 31.68$   
 $np > 10$   $npq > 10$  therefore  
 a normal distribution can apply. ✓✓
- (ii)  $\hat{p} \sim N(0.12, 0.01876^2)$  ✓✓
- (iii)  $P(\hat{p} > 0.125) = 0.3949$  ✓
- (iv)  $P(\hat{p} < k) = 0.25$   $k = 0.1073$  ✓  
 10.7% ✓
- (b) (i)  $\int_5^{18} \frac{1}{15} dx = \frac{13}{15}$  ✓
- (ii) Expected waiting time = 12.5 days ✓  
 Normal distribution  $\hat{p} \sim N(0.5, 0.07906^2)$  ✓✓
- (iii)  $P(\hat{p} < 0.7) = 0.9943$  ✓ [12]
20. (a)  $\ln\left(\sqrt{\frac{x+1}{x^3}}\right) = \frac{1}{2} \ln(x+1) - \frac{3}{2} \ln x$  ✓
- $\frac{d}{dy} \left( \frac{1}{2} \ln(x+1) - \frac{3}{2} \ln x \right) = \frac{1}{2} \left( \frac{1}{x+1} \right) - \frac{3}{2x}$  ✓✓
- $= \frac{1}{2x+2} - \frac{3}{2x}$
- (b) (i)  $pH = -\log_{10} [H_3O^+]$   
 Hydronium ions range from  
 $[H_3O^+] = 10^{-4.5} = 0.00003162$  moles per litre ✓  
 to  
 $[H_3O^+] = 10^{-4.7} = 0.000019953$  moles per litre ✓
- (ii)  $pH = -\log_{10} [H_3O^+] = -\log_{10} 4.8 \times 10^{-8}$  ✓  
 $pH = 7.32$  therefore it is not acidic. ✓ [7]



21. (a)  $p = \frac{28}{80}$  ✓
- (b) (i) (0.2455, 0.4545) ✓✓
- (ii)  $ME = \frac{1.960 \sqrt{\frac{\frac{28}{80} \left( \frac{52}{80} \right)}}{80} = 0.1045$  ✓
- (iii) 95% the width is 0.209 ✓
- 99% the width is 0.2747 ✓
- $0.209x = 0.2747$  ✓
- $x = 1.3144$  ✓
- Therefore an increase of 31.45%
- (c)  $1.960 \sqrt{\frac{\frac{28}{80} \left( \frac{52}{80} \right)}{n}} = 0.02$  ✓
- $n = 2184.9$  ✓
- Therefore sample size of 2185 ✓
- (d)  $0.95 \times 40 = 38$  ✓
- ∴ Approximately 38 would contain the true proportion ✓ [10]
22. (a)  $-192\cos(4t) - 80\sin(4t) = 0$
- $t = 1.27679$  (max) ✓
- $v(t) = 20\cos(4t) - 48\sin(4t) + c$
- $v(t) = 20$  when  $t = 0$  therefore  $c = 0$  ✓
- $v(1.27679) = 52$  units/sec ✓
- (b)  $v(2) = -50.4$  units/sec ✓
- $a(2) = -51.2$  units/sec<sup>2</sup> ✓
- The velocity and the acceleration are both negative, therefore the particle is moving faster to the left.
- ∴ Increasing speed ✓
- (c)  $x(t) = 5\sin 4t + 12\cos 4t + c$
- $x(0) = 12 \therefore c = 0$  ✓
- $x(2) = 3.2008$  units ✓
- (d)  $\int_4^5 20\cos(4t) - 48\sin(4t) dt = 22.393$  units ✓✓
- (e)  $\int_0^5 |20\cos(4t) - 48\sin(4t)| dt = 160.538$  units ✓✓ [12]

End of Questions