

Question	Marks	Max	Question	Marks	Max
12			12		
11			10		
10			17		
8			8		
9			16		
9			15		
8			11		
7			10		
7			13		
9			14		
9			15		
8			11		
8			10		

No other items may be taken into the examination room. It is your responsibility to ensure that you do not have any unauthorised material if you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

Important note to candidates
and up to three calculators approved for use in this examination
drawing instruments, templates, notes on two unruled sheets of A4 paper,
Special items:

To be provided by the candidate
Standard items: pens (blue/black preferred), pencils (including coloured), sharpener,
correction fluid/tape, eraser, ruler, highlighters

To be provided by the supervisor
Materials required/recommended for this section
Formula sheet (retained from Section One)
This Question/Answer booklet

Time allowed for this section
Materials required/recommended for this section
Working time:
Reading time before commencing work: ten minutes
one hundred minutes

Your Teacher's Name:

Your Name:

Calculator-assumed
Section Two:

UNIT 3 & 4
MATHEMATICS METHODS

Question/Answer booklet

Semester Two Examination, 2021



INDEPENDENT PUBLIC SCHOOL
Exceptional school living. Exceptional students.

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	6	6	50	50	33
Section Two: Calculator-assumed	11	11	100	100	67
Total					100

Instructions to candidates

1. The rules for the conduct of the Western Australian Certificate of Education ATAR course examinations are detailed in the *Year 12 Information Handbook 2019*. Sitting this examination implies that you agree to abide by these rules.
2. Write your answers in this Question/Answer booklet.
3. You must be careful to confine your answers to the specific questions asked and to follow any instructions that are specific to a particular question.
4. Additional pages for the use of planning your answer to a question or continuing your answer to a question have been provided at the end of this Question/Answer booklet. If you use the space to continue an answer, indicate in the original answer space where the answer is continued, i.e. give the page number.
5. **Show all your working clearly.** Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
6. It is recommended that you **do not use pencil**, except in diagrams.
7. The Formula sheet is **not** to be handed in with your Question/Answer booklet.

\checkmark Determines the distance between 2 and 3 seconds
Specific Behaviours
$19.6 + 4.9 = 24.5 \text{ m}$
$\int_0^2 -9.8t + 19.6 = -4.9m$
OR
$\int_0^3 -9.8t + 19.6 \checkmark = -24.5m \checkmark \quad (3 \text{ marks})$
Solution

(c) Determine the **total distance** travelled over the time interval $0 \leq t \leq 3$. (3 marks)

\checkmark Sets up the correct integral.
Specific Behaviours
$15.6 + \int_0^2 -9.8t + 19.6 = 34.9 \text{ m}$
Solution

(b) Determine this maximum height of the projectile. (2 marks)

\checkmark States the expression for $v(t)$.
Specific Behaviours
$v(t) = 0, t = 2 \text{ s}$
$v(t) = -9.8t + 19.6$
Solution

(a) Given that $a(t) = -9.8 \text{ m/s}^2$, determine an expression for $v(t)$ and use it to find the time at which the projectile reaches its maximum height. (2 marks)

A projectile is fired upward from a cliff at a speed of 19.6 m/s and falls into a valley below. The acceleration g due to Earth's gravity is about 9.8 m/s^2 downwards.

Working time: 100 minutes.

Question 7

- Planning: if you use the spare pages for planning, indicate this clearly at the top of the page.
 - Continuing an answer: if you need to use the space to continue an answer, indicate this clearly at the top of the page.
 - Original answer space where the answer is continued: i.e. give the page number. Fill in the number of the question that you are continuing to answer at the top of the page.
- Spare pages are included at the end of this booklet. They can be used for planning your responses and/or as additional space if required to continue an answer.
- This section has **eleven** questions. Answer all questions. Write your answers in the spaces provided.
- Section One: Calculator-assessed
(100 Marks)

Question number: _____

Additional working space

- ✓ Adds the distance to 19.6m
- ✓ Determines the total distance

Question 8 (9 marks)

The table below summarises census information about the number of children in the households of an Australian town.

Number of children	0	1	2	3	4 or more
Percentage of households	23	32	35	7	3

A random sample of 20 households is selected from this town.

- a. State the distribution and determine the probability that the sample will contain:

- i. 3 households only with no children. (2 marks)

Solution

X the number of household with no children

$$X \text{ Bin}(20, 0.23)$$

$$P(X=3) = \text{binomialPDF}(3, 20, 0.23) = 0.163091$$

Specific Behaviours

- ✓ States binomial distribution with correct parameters
- ✓ Determines the correct probability

- ii. more than half the households, with at least 2 children. (3 marks)

Solution

Y at least 2 children household

$$Y \text{ B}(20, 0.45)$$

$$P(Y > 10) = 1 - P(Y \leq 10)$$

$$1 - \text{binomialCDF}(0, 10, 20, 0.45)$$

$$0.2492893598$$

Specific Behaviours

- ✓ States binomial distribution with correct parameters
- ✓ Recognises the correct probability expression $P(Y > 10)$
- ✓ Determines the correct probability

Additional working space

Question number: _____

Solution	
Specific Behaviours	
	<ul style="list-style-type: none"> ✓ States binomial distribution with correct parameters ✓ Checks probability for $n=4$ ✓ Checks probability for $n=3$ ✓ Concludes $n=4$
	$\therefore n = 4.$
	$\text{When } n = 5, \quad P(A \geq 1) > 0.1$ $1 - P(A = 0) > 0.1$ $P(A = 0) < 0.9$ $P(A = 0) < 0.03^5 \cdot 0.9^5 = 0.858$ $n = 5, \quad P(A = 0) = 0.03^5 \cdot 0.9^5 = 0.885$ $n = 4, \quad P(A = 0) = 0.03^4 \cdot 0.9^4 = 0.913$ $n = 3, \quad P(A = 0) = 0.03^3 \cdot 0.9^3 = 0.973$
<p style="text-align: right;">Question number: _____</p>	

A new random sample of households is selected from this town. The probability that this new sample contains a household with 4 or more children is more than 10%.

b. Determine the smallest value of n . (4 marks)

Additional working space	
	<ul style="list-style-type: none"> ✓ States the optimal route is to run around the pool. ✓ Calculates $T(0)$ and $T\left(\frac{\pi}{L}\right)$.

Question 9**(9 marks)**

When refuelling the car, the rate of flow of petrol into the tank is given by

$$\frac{dV}{dt} = 9e^{-(t+2)}(8-t) \text{ for } 0 \leq t \leq 8,$$

where V is the litres of petrol in the tank at time t in minutes. Initially the tank has two litres of petrol.

(a)

- i. Determine the **exact** initial rate of flow of petrol into the tank. (1 mark)

Solution	
Define $f(t)=9 \cdot e^{-(t+2)} \cdot (8-t)$	done
$f(0)$	$72 \cdot e^{-2}$
Specific behaviours	

P States correct value

- ii. Determine the value of t for which $\frac{dV}{dt}=0$. (1 mark)

Solution	
solve($f(t)=0, t$) $0 \leq t \leq 8$	{ $t=8$ }
Specific behaviours	

P States correct value


- iii. Determine the time, to the nearest second, when the rate is 1 litre per minute. (2 marks)

Solution	
solve($f(t)=1, t$) $0 \leq t \leq 8$	
{ $t=1.990556694$ }	
60×0.990556694	
59.43340164	
$t=1$ minutes 59 seconds	
Specific behaviours	

PP states to nearest second (do not accept decimal minutes)

Question 17**(7 marks)**

Consider a lifeguard at a circular pool, at position A, with diameter of 60 m. He must reach someone who is drowning on the exact opposite side of the pool, at position C. The lifeguard is considering taking a two-stage route, first swimming at angle θ with a speed 1 m/s to position B and then leaving the pool and running around the pool at speed 3 m/s.



- (a) Determine the function $T(\theta)$ as the total amount of time it takes to reach the drowning person, in terms of the swim angle θ (radians). Hint: Angle BOC = 2 θ . (3 marks)

Solutions	
AB = $60 \cos \theta$,	
Arclength BC = $30 2\theta = 60\theta$	
$T(\theta) = \frac{60\theta}{3} + \frac{60 \cos \theta}{1} = 20\theta + 60 \cos \theta$	
Behaviour	

✓ Determines the correct expression for AB.
✓ Determines the correct expression for arc length BC.
✓ Determines the correct expression for $T(\theta)$. No need to simplify.

- (b) Using Calculus, justify whether this two-stage route will **minimise** the time the lifeguard takes to reach the drowning person. If not, determine the route that will. (4 marks)

Solutions	
$T'(\theta) = 20 - 60 \sin \theta = 0, \theta \approx 0.34 \text{ radians}$	
$T''(\theta) = -60 \cos(0.34) < 0, \text{local maximum}$	
Hence, the two stage route will not give minimum time.	
$T(0) = 60 \text{ s}, T\left(\frac{\pi}{2}\right) = 31.41593 \text{ s}$	
Therefore, the fastest route is to run around the pool directly.	
Behaviour	

✓ Equates $T'(\theta)$ to 0 and solve for θ in radians.
✓ Uses 2nd derivative to show it is a local max and hence not the optimal route.

Solution	
Specific behaviours	
P states to nearest second (do not accept decimal minutes)	P states the correct definite integral
2 minutes 22 seconds	
22.29878094	
60×0.371646349	
$\{x=2, 371646349\}$	
$\int_0^x f(t) dt = 8, x 0 \leq x \leq 8$	
solve(

(c) Determine the time, to the nearest second, when there are 10 litres of petrol in the tank. (2 marks)

Solution	
Specific behaviours	
$V = 10.5 \text{ L}$	
10.52653144	
8.526531443 + 2	
8.526531443	
$\int_8^0 f(t) dt$	

(b) Determine the amount of petrol in the tank, to one decimal place, when $t = 8$. (3 marks)

P justifies answer by saying that only 90% of intervals are expected to contain the true proportion OR compares the overlap with a new 90% confidence interval using 0.2528

Question 10**(8 marks)**

The volume, $V \text{ cm}^3$, of a plastic bottle is given by $V = \pi r^2 h + \frac{2}{3} \pi r^3$, where $h \text{ cm}$ is the height and $r \text{ cm}$ is the radius of its cap.

- (a) Given that $h=3 \text{ cm}$, what is the approximate increase in volume when the radius expands by $p \text{ cm}$ from 2 cm ? Use a calculus method to find your answer. (4 marks)

Solution	
$\delta V \approx \frac{dV}{dr} \times \delta r \approx (2\pi r^2 + 6\pi r) \times \delta r \text{ At } r=2 \text{ cm, } \delta r=p \text{ cm}$ $\delta V \approx [2\pi \times 2^2 + 6\pi \times 2] \times p = 20\pi p \text{ cm}^3$	
Specific behaviours	
P Determines $\frac{dV}{dr}$ P Determines an expression for δV in terms of r P Substitutes the correct δr P Determines the correct δV	

Question 16**(8 marks)**

5000 boxes of a certain kind of cereal are stored in a warehouse. The cereal manufacturer wishes to estimate the proportion of boxes in the warehouse that weigh less than the 300g stated on the box. The manufacturer asks a warehouse employee to sample and weigh 200 cereal boxes. The boxes are stored in the warehouse on 50 numbered shelves (with 100 boxes arranged in a line on each shelf).

- a) Briefly describe a suitable method for selecting the sample of 200 boxes. (2 marks)

Solution	
The employee could use a random number generator. E.g. to select each box they could first generate a random number from 1-50 to select the shelf, and then generate a random number from 1-100 to select the box from that shelf.	
Specific behaviours	
P indicates some random mechanism P indicates that the boxes are selected accordingly	

The warehouse employee finds that 38 of the 200 selected boxes weigh less than 300g.

- b) Determine the sample proportion of underweight cereal boxes, and thus show that a 90% confidence interval for the proportion of underweight boxes (to 3 decimal places) is $(0.144, 0.236)$. (4 marks)

Solution	
$\hat{p} = \frac{38}{200} = 0.19$ $90\% \text{ confidence interval} = \left(0.19 - 1.645 \times \sqrt{\frac{0.19 \times 0.81}{200}}, 0.19 + 1.645 \times \sqrt{\frac{0.19 \times 0.81}{200}} \right)$ $(0.19 - 0.046, 0.19 + 0.046) \rightarrow (0.144, 0.236)$	
Specific behaviours	
P states the correct sample proportion P uses the correct value for the standard error P uses the correct z-value interval P shows correct method for calculating the confidence interval	

- c) The manufacturer later consults factory records to conduct a census of the weights of all 5000 boxes in the warehouse, and finds that the proportion of underweight boxes is 0.2528 (assume that this is the correct proportion). The manufacturer accuses the employee of being careless in their measurements or calculations. Is this accusation justified? Briefly explain your answer. (2 marks)

Solution	
No. Only 90% of 90% confidence intervals are expected to contain the true proportion. It is possible that the survey and calculation by the warehouse employee was performed appropriately, but happened to yield one of the 10% of confidence intervals that do not contain the true proportion.	
Specific behaviours	
P answers 'No' with a reference to part (b) - MUST make reference to the confidence interval calculated in part (b)	

$$-\frac{dx}{\sqrt{\frac{1}{3} \cdot a^3 \cdot (\sin(x))^2 \cdot (1+\cos(x))}} = \frac{540}{-(a^3 \cdot (\sin(x))^3 \cdot \sqrt{2-a^2} \cdot (\cos(x))^2 \cdot \sin(x) \cdot \sqrt{2-a^2} \cdot \cos(x) \cdot \sin(x) \cdot \sqrt{2})}$$

If used CAS

$$\theta = 70.5^\circ \text{ (Reject } 180^\circ \text{ and } 289.5^\circ)$$

$$\cos \theta = \frac{3}{1} \text{ or } \cos \theta = -1$$

$$(3\cos \theta - 1)(\cos \theta + 1) = 0$$

$$3\cos^2 \theta + 2\cos \theta - 1 = 0$$

$$2\cos \theta + 2\cos^2 \theta = 1 - \cos^2 \theta$$

$$2\cos \theta (1 + \cos \theta) = 1 - \cos^2 \theta$$

$$\text{Since } \sin \theta \neq 0, 2\cos \theta (1 + \cos \theta) = \sin^2 \theta$$

$$2\sin \theta \cos \theta (1 + \cos \theta) = \sin^3 \theta$$

$$-\sin^3 \theta + 2\sin \theta \cos \theta (1 + \cos \theta) = 0$$

$$\frac{1}{3} \pi a^3 (-\sin^3 \theta + 2\sin \theta \cos \theta (1 + \cos \theta)) = 0$$

$$\frac{d\theta}{dV} = 0 \text{ for maximum or minimum volume}$$

$$\frac{d\theta}{dV} = \frac{1}{3} \pi a^3 (-\sin^3 \theta + 2\sin \theta \cos \theta (1 + \cos \theta))$$

$$\frac{d\theta}{dV} = \frac{1}{3} \pi a^3 (\sin^2 \theta (-\sin \theta) + (1 + \cos \theta) 2\sin \theta \cos \theta)$$

Solution

<p>P Uses incremental formula</p> <p>P Determines an expression for $\frac{dV}{dr}$ in terms of r</p> <p>P Substitutes and states correct expression for $\frac{dV}{dr}$</p> <p>P Simplifies and states correct percentage change</p>	
<p>Specific behaviours</p>	
$\frac{dV}{dr} \approx \frac{V}{r} \times \frac{6r}{3(r+6)} \times \frac{6\pi r^2 + \frac{3}{2}\pi r^3}{2\pi r^2 + 12\pi r}$ $A/r = 4\text{cm}, \frac{dV}{dr} = a%$ $V \approx \frac{A}{3} \times \frac{4+9}{4+6} \times a = 2.3\% \text{ a}$	
$\frac{dV}{dr} = \frac{V}{r} \times \frac{6r}{3(r+6)} \times \frac{(4+9)}{3 \cdot (4+6)} \times a$ $2.307692308 \cdot a$	
$\frac{d}{dr}(f(t))$ done	
$\text{Define } f(r) = 6\pi r^2 + \frac{2\pi r^3}{3}$ $\text{Simplify } (2\pi r^2 + 12\pi r)$ $2\pi r^2 + 12\pi r$ $6\pi r^2 + 2\pi r^3$ $\frac{d}{dr}(f(r))$ $\frac{r \cdot (r+9)}{3 \cdot (r+6)}$ $\frac{3 \cdot (r+9)}{3 \cdot (r+6)} \times a$	

- (b) If $h=6\text{cm}$ and $r=4\text{cm}$, use calculus to find the approximate percentage change, to one decimal place, in the volume when the radius increases by $y\%$.

- d) Given that a is a constant, find $\frac{dV}{d\theta}$ and hence find the value(s) of θ for which the volume is maximised.

volumne is maximised.

volume is maximised.


- ✓ Substitutes results from (a) and (b) into the volume formula
- ✓ Factors out $\frac{1}{3} \pi (a^3 \sin^2 \theta)$

Question 11**(10 marks)**

An aircraft designer is trailing a new wing shape with a cross section at a certain point enclosed by the function

$$y = -k(x^2 - 0.01) \ln(x)$$

where k is a positive constant, together with an interval of the x -axis from $x=a$ to $x=b$.



- (a) Determine the values of a and b (given that $a < b$). (2 marks)


Solution	
$-k(x^2 - 0.01) \ln(x) = 0$	
$x^2 - 0.01 = 0$ or $\ln(x) = 0$	
$x = 0.1$ or $x = 1$	
$a = 0.1$ and $b = 1$	
Specific behaviours	
P equates function to 0	
P states both correct values	

- (b) Determine the value of c (to 3 decimal places) above which the highest point of the top edge of the cross section occurs. (3 marks)

Solution	
$\frac{dy}{dx} = -k(2x \ln(x) - k(x^2 - 0.01) \times \frac{1}{x}) = 0$	
$x = 0.61461$	
$c = 0.615$	
Specific behaviours	
P writes correct derivative	
P equates derivative to 0	
P states correct value of c to 3 decimal places	

Question 15**(9 marks)**

An engineering project requires an original part PBQ that is conical in shape. At a later stage of the project, a smaller cone PAQ will be extracted from the original such that $AP = AB = AQ$. We will use the letter a to represent the three equal lengths AQ , AB and AP . We will use the letter r to represent the radii OP and OQ of the circular section of the cone. We will use the letter h to represent the altitude OB of the original cone. Both angles $\angle OAP$ and $\angle OAQ$ have a magnitude θ . It is advised that you label the diagram accordingly.



- a) Find h , the altitude of the original cone, in terms of a and θ . (1 mark)

Solution	
$h = a + a \cos\theta$	
Specific behaviours	
✓ States h in terms of a and θ	

- b) Find r , the radius of the circular section of the cone, in terms of a and θ . (1 mark)

Solution	
$r = a \sin\theta$	
Specific behaviours	
✓ States r in terms of a and θ	

The volume of the cone is given by $V = \frac{1}{3}\pi r^2 h$.

- c) Use the results from (a) and (b) to show that (2 marks)

$$V = \frac{1}{3}\pi a^3 \sin^2 \theta (1 + \cos\theta)$$

Solution	
$V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi (a^2 \sin^2 \theta)(a + a \cos\theta)$	
$= \frac{1}{3}\pi (a^3 \sin^2 \theta) + \frac{1}{3}\pi (a^3 \sin^2 \theta)(\cos\theta)$	
$= \frac{1}{3}\pi a^3 \sin^2 \theta (1 + \cos\theta)$	
Specific behaviours	

	Solution
	<p>P substitutes value of c from part (b) into function</p> <p>P equates to 0.15</p> <p>P states correct value of k to 3 decimal places</p> <p>$k = 0.838$</p>
(c)	<p>Find, to 3 decimal places, the value of k such that the maximum height of the cross section is 0.15 units.</p> <p>$0.15 = -k(0.615^2 - 0.01) \ln(0.615)$</p>
(d)	<p>Determine, to 3 decimal places, the area of the cross section for the value of k determined in part (c).</p> <p>$\int_1^{0.1} -0.838(x^2 - 0.01) \ln(x) dx = 0.087$</p>

- (c) Find, to 3 decimal places, the value of k such that the maximum height of the cross section is 0.15 units.
- (d) Determine, to 3 decimal places, the area of the cross section for the value of k determined in part (c).

Question 12**(12 marks)**

Two electrical engineering companies produce light globes – one company produces type A and the other produces type B. Both types are normally distributed. A light globe is considered premium if it has a lifespan longer than 850 hours.

- (a) The company producing type A globes claims that their product has a mean lifespan of 818 hours and a standard deviation of 112 hours.

- i. What is the probability of a type A globe being premium? (2 marks)

Solution

$$A \sim N(818, 112^2)$$

$$P(A > 850) = 0.3875$$

Specific Behaviours

- ✓ Uses the correct probability statement
- ✓ Determines the correct probability

- ii. What is the probability of a type A globe being premium, given that its lifespan exceeds 800 hours? (2 marks)

Solution

$$P(A > 850 | A > 800) = \frac{P(A > 850)}{P(A > 800)} = \frac{0.3875}{0.5638} = 0.6873$$

Specific Behaviours

- ✓ Uses the conditional probability formula
- ✓ Determines the correct probability

- iii. What lifespan is exceeded by 90% of all type A globes produced? (2 marks)

Solution

$$P(A > k) = 0.9$$

$$z = -1.2816$$

$$-1.2816 = \frac{k - 818}{112}$$

$$\therefore k = 674.47$$

Specific Behaviours

- ✓ Determines the z-score
- ✓ Uses the z-score to determine the value of k

- iv. If the company selects 50 type A batteries one at a time, find the probability that it takes a selection of eight batteries before six premium batteries are selected. (3 marks)

Solution

$$X \sim Bin(7, 0.3875)$$

$$P(X=5) \times 0.3875$$

$$0.06883 \times 0.3875$$

$$0.02667$$

Specific Behaviours

- ✓ Sets up a binomial distribution with correct parameters

- (ii) What proportion of accelerometers will operate at least at 100 °C (2 marks)

Solution

$$Pr(T < 100) = Pr\left(Z \leq \frac{100 - 95.6692}{5.6727}\right)$$

CAS Solve

$$\frac{100 - 95.6692}{5.6727} = 0.672716695$$

$$0.7634417005$$

$$normCDF(-\infty, 0.7634417005, 1, 0)$$

$$0.7773999907$$

$$Pr(T \leq 100) = 0.7774$$

$$\therefore Pr(T \geq 100) = 1 - 0.7774 = 0.2226$$

22.26% of accelerometers will operate at 100 °C

Specific behaviours

- ✓ States the proportion that will operate at or below 100 °C

- b) During the process of thorough testing, the manufacturer took repeated samples and one sample of 300 accelerometers revealed that 51 continued to function at 100 °C

- (i) State the approximate distribution of the sample proportions and justify your choice. (3 marks)

Solution

$$\hat{p} \sim N\left(0.17, \frac{0.17 \times 0.83}{300}\right)$$

$$\hat{p} \sim N(0.17, 0.0216871^2)$$

A normal approximation model is appropriate as the value of n is large enough for the distribution to approach normality.

Specific behaviours

- ✓ Indicates Normal distribution
- ✓ Calculates the correct mean and standard deviation (or variance)
- ✓ Correctly justifies why the normal distribution can be used

- (ii) Provide a 99% confidence interval for the proportion of accelerometers that can function at a temperature of at least 100 °C (2 marks)

Solution

$$\text{Lower } 0.1141376$$

$$\text{Upper } 0.2258624$$

$$\hat{p} 0.17$$

$$n 300$$

A 99% confidence interval is [0.1141, 0.2259]

Question 14	
Solution	(11 mark)
<p>A company that manufactures professional photogrammetry drones uses a particular model of accelerometer that fails when it becomes too hot. The temperature $T^\circ\text{C}$ at which a random sample of accelerometers fails normally distributed.</p> <p>Accelerometer fails at least 105°C.</p> <p>In preliminary laboratory tests, 97% of a random sample of accelerometers continues working at least 105°C.</p> <p>(i) Calculate the mean and standard deviation of the distribution</p>	$\Pr\left(Z \leq \frac{85 - \mu}{\sigma}\right) = 0.97$ $\Pr\left(Z \leq \frac{105 - \mu}{\sigma}\right) = 0.05$ $\frac{105 - \mu}{\sigma} = -1.880794$ $\frac{\mu - 85}{\sigma} = 1.6448536$ $\sigma = 95.66921152, \mu = 5.672716695$
<p>(b) The company produces type B light globes that 31% of their product has a lifespan over 860 hours, and 19% have a lifespan below 700 hours. Determine the mean and standard deviation for the type B light globes.</p> <p>(3 marks)</p>	$P(B > 860) = 0.312 = 0.4959$ $P(B < 700) = 0.192 = 0.8779$ $0.4959 = \frac{860 - \mu}{\sigma}$ $-0.8779 = \frac{700 - \mu}{\sigma}$ $(CAS Simultaneous): \mu = 802.25 \text{ and } \sigma = 116.47$
<p>Specific Behaviours</p> <ul style="list-style-type: none"> ✓ Determines $P(X=5)$ ✓ Determines the correct probability 	<p>Specific Behaviours</p> <ul style="list-style-type: none"> ✓ Sets up simultaneous equations to solve for the parameters ✓ Determines the z-scores for both probabilities statements ✓ Determines the mean and standard deviation ✓ Solves simultaneously

Solution	
<p>The mean and standard deviation are approximately 95.7°C and 5.6°C.</p> <p>The mean and standard deviation for the standard score of 105°C respectively.</p>	<p>Specific behaviours</p> <ul style="list-style-type: none"> ✓ Forms an equation for the standard score of 85°C ✓ Forms an equation for the standard score of 105°C
<p>(ii) Calculate the mean and standard deviation of the distribution</p>	<p>Specific Behaviours</p> <ul style="list-style-type: none"> ✓ An equation for the standard score of 105°C

Question 13**(10 marks)**

A discrete random variable Y is defined by $P(Y=y) = a \log(x-1)$ for $x=6, 11$ and 21

(a) Determine the value of a .

(3 marks)

Solution

$$a \log(5) + a \log(10) + a \log(20) = 1$$

$$a \log(5 \times 10 \times 20) = 1$$

$$a \log 1000 = 1$$

$$a = \frac{1}{\log 1000} = \frac{1}{3}$$

Specific Behaviours

- ✓ Substitutes and sums the probabilities to 1
- ✓ Uses log laws to add the logs together
- ✓ Determines the value of a

(b) Determine $P(Y=21|Y>6)$ in exact form.

(3 marks)

Solution

$$P(Y=21|Y>6) = \frac{P(Y=21)}{P(Y>6)}$$

$$\frac{\frac{1}{3} \log 20}{\frac{1}{3} \log(10 \times 20)}$$

$$\frac{\log 20}{\log 200}$$

Specific Behaviours

- ✓ Uses the conditional probability formula
- ✓ Substitutes the correct numerator
- ✓ Substitutes the correct denominator and leaves answer in exact form

(c) The expected value $E(Y) = \frac{38}{3} + m \log n$ where the constants m and n are prime numbers. Determine the values of m and n . (4 marks)

Solution

$$E(Y) = 6\left(\frac{1}{3} \log 5\right) + 11\left(\frac{1}{3} \log 10\right) + 21\left(\frac{1}{3} \log 20\right)$$

$$\textcolor{red}{i} 2 \log 5 + \frac{11}{3} \log 10 + 7 \log 20$$

$$\textcolor{red}{i} 2 \log 5 + \frac{11}{3} + 7 \log 20$$

$$\textcolor{red}{i} 2 \log 5 + \frac{11}{3} + 2 \log 20 + 5 \log 20$$

$$\textcolor{red}{i} 2 \log 100 + \frac{11}{3} + 5 \log 2 + 5 \log 10$$

$$\textcolor{red}{i} 4 + 5 + \frac{11}{3} + 5 \log 2$$

$$\textcolor{red}{i} \frac{38}{3} + 5 \log 2$$

$$\therefore m=5 \text{ and } n=2$$

Specific Behaviours

- ✓ Expresses $E(Y)$ as the sum of logs
- ✓ Simplifies and splits the $7 \log 20$ term
- ✓ Combines logs and splits $5 \log 20$ term
- ✓ Determines the values of m and n