

The 90% confidence interval of the sample proportion \hat{p} , from the initial survey is $0.649 \leq \hat{p} \leq 0.725$.

(d) Use the 90% confidence interval of the initial sample to compare the following samples:

- (i) A random sample of 365 people at a shopping centre found that 258 had a preference for the phablet style smart phone. (2 marks)

Solution
$\hat{p} = \frac{258}{365} = 0.71$ and $0.668 \leq \hat{p} \leq 0.746$ ✓
The confidence interval for this second survey overlaps, significantly, the 90% confidence interval of the initial survey so this indicates we are sampling from the same population.
Specific behaviours
✓ calculates 90% confidence interval for \hat{p} correctly ✓ states the similarity of results

8 d (ii)

$$\hat{p} = \frac{52}{75} = 0.693$$

and $0.5789 \leq \hat{p} \leq 0.7545$ ✓

Again the \hat{p} falls within the C.I. and is similar to initial survey results so sampling from the same population.

(No need to talk about Brian: Maths Teacher inside Retirement Village) $\overline{Ch S}$ 3 apps

Any reasonable comment ✓



PERTH MODERN SCHOOL
exceptional schooling, exceptional students.

Continuous Random Variables
The Normal Distribution
Sample Proportions

Test 5

Semester Two 2018
Year 12 Mathematics Methods
Calculator Assumed

Name: Sol 4 Trends

Date: Fri 17th Aug. 7:45am

You may have a formula sheet for this section of the test.

Classpad Calculators

1 page of Notes

Total _____/46

50 minutes

- Teacher: _____
- Mr McClelland _____
- Mrs. Berry _____
- Mr Gannon _____
- Mrs Cheng _____
- Mr Staffe _____
- Mr Strain _____

Question 1

(5 marks)

The life of an electronic component is given by the probability density function:

$$f(x) = \begin{cases} \frac{100}{x^2} & x > 100 \\ 0 & \text{otherwise} \end{cases}$$

Find:

- (a) the probability that a component lasts for more than 250 hours.

(2 marks)

$$1 - \int_{100}^{250} \frac{100}{x^2} dx = 0.4$$

- (b) the median life of a component.

(2 marks)

$$\int_{100}^{\infty} \frac{100}{x^2} dx = 0.5 \Rightarrow \left[-\frac{100}{x} \right]_{100}^{\infty} = 100 \left[0 - \left(-\frac{1}{100} \right) \right] = \frac{100}{k} = 0.5$$

- (c) the lifetime for 95% of components.

(1 mark)

$$\int_k^{\infty} \frac{100}{x^2} dx = 0.05; k = 2000 \text{ hrs} \quad \left\{ \begin{array}{l} P(100 < X \leq k) = 0.95 \\ \therefore \text{The Lifetime is } 100 < X \leq 2000 \end{array} \right.$$

Question 2

(4 marks)

- (a) $\Pr(Z < -0.376)$, where Z is a standard normal variable is:

(1 mark)

$$X \sim N(0, 1) \Rightarrow 0.3535 \checkmark$$

- (b) If Z is a standard normal random variable, and $\Pr(Z > c) = 0.75$, then the value of c is?

(1 mark)

$$c = -0.6745 \checkmark$$

- (c) If X is a normally distributed random variable with mean $\mu = 4$ and standard deviation, $\sigma = \sqrt{2}$, then the transformation that maps the curve of the density function of X , $f(x)$, to the curve of the standard normal distribution is:

(2 marks)

$$Z = \frac{x - \mu}{\sigma} = \frac{x - 4}{\sqrt{2}}$$

$$\therefore (x, y) \rightarrow \left(\frac{x - 4}{\sqrt{2}}, \sqrt{2}y \right)$$

Question 8

(10 marks)

A random survey was conducted to estimate the proportion of mobile phone users who favoured standard smart phones over the new *phablet* style smart phones. It was found that 283 out of 412 people surveyed preferred the new *phablet* style smart phones.

- (a) Determine the sample proportion \hat{p} of those in the survey who preferred a phablet style smart phone.

(1 mark)

Solution
$\hat{p} = \frac{283}{412} = 0.6869$
Specific behaviours
✓ calculates \hat{p} correctly

- (b) Use the survey results to estimate the standard deviation of \hat{p} , for the sample proportions.

(2 marks)

Solution
Standard deviation = $\sqrt{\frac{283}{412} \left(1 - \frac{283}{412} \right)} = 0.0228$
Specific behaviours
✓ substitutes correctly into standard deviation formula ✓ calculates standard deviation correctly

- (c) A follow-up survey is to be conducted to confirm the results of the initial survey. Working with a confidence interval of 95%, estimate the sample size necessary to ensure margin of error of at most 4%.

(3 marks)

$$0.6869 \quad 0.3/31 \quad n = 517 \checkmark$$

Specific behaviours
✓ writes an equation to evaluate n from the margin of error ✓ solves correctly for n ✓ rounds n up to the nearest whole number

(c) Determine the probability that in a random sample of 120 people, the number who had taken a plane flight in the last year was greater than 26. (3 marks)

Solution

The distribution is binomial with $p = 0.19$ and $n = 120$.
 $P(X > 26) = P(X \geq 27)$, since n is discrete

If $n \neq 26$
 $\text{prob} = 0.2602$ [2 marks]

BinomialCD

Lower	27
Upper	120
Normal	120
pos	0.19

prob 0.9998

BinomialCD

Lower	27
Upper	120
Normal	120
pos	0.19

prob 0.9998

Hence the required probability is 0.9998 (to four decimal places)

Specific behaviours

- identifies the distribution as binomial - bin(120, 0.19)
- uses 27 as the lower bound in the binomial cumulative distribution
- states the correct probability

(d) If seven surveys were taken and for each a 95% confidence interval for p was calculated, determine the probability that at least four of the intervals included the true value of p . (2 marks)

Solution
$\text{bin}(7, 0.95) \Rightarrow P(4 \leq x \leq 7) = 0.9998$
Specific behaviours
identifies the distribution as binomial - bin(7, 0.95)
calculates the probability correctly


Accept: Normal Dist

$$\sigma = \sqrt{0.19 \times 0.81} = 0.0358$$

$$P(X > 26) \sim N(0.19, 0.0358^2)$$


$$P(X > 26) = 0.2282$$

Classical normal $\left(\frac{26}{120}, \infty, 0.0358, 0.19\right)$



Question 3

The weight of a population of teenage females is normally distributed with a mean of 55 kg and a standard deviation of 8 kg. If the lowest 5% of teenage females is classified as underweight, what is the cut-off weight for this group?



Question 4

A probability density function is given by

$$f(x) = 4x(6 - x)^2 \quad 0 < x < 6$$

Find the value of A and hence the mean and the standard deviation of this distribution.

$$A \int_0^6 x(6-x)^2 dx = 1$$

$$\therefore A = \frac{1}{108} = 0.009259$$

$$E(X) = \frac{1}{108} \int_0^6 x^2 \times x(6-x)^2 dx$$

$$= \frac{1}{108} \int_0^6 x^3(6-x)^2 dx$$

$$= \frac{1}{108} \int_0^6 (x^3(36 - 12x + x^2)) dx$$

$$= \frac{1}{108} \int_0^6 (36x^3 - 12x^4 + x^5) dx$$

$$= \frac{1}{108} \left[9x^4 - \frac{12}{5}x^5 + \frac{1}{6}x^6 \right]_0^6$$

$$= \frac{1}{108} \left[9(6^4) - \frac{12}{5}(6^5) + \frac{1}{6}(6^6) \right]$$

$$= \frac{1}{108} \left[9(1296) - \frac{12}{5}(7776) + \frac{1}{6}(46656) \right]$$

$$= \frac{1}{108} \left[11664 - 18662.4 + 7776 \right]$$

$$= \frac{1}{108} \left[7776 \right]$$

$$= 72$$

$$= 2.4$$

$$\text{Var}(X) = E(X^2) - \mu^2$$

$$= 7.2 - 2.4^2$$

$$= 1.44$$

$$\therefore \sigma_x = 1.2$$

$$E(X^2) = \int_0^6 x^2 \times f(x) dx$$

$$= \frac{1}{108} \int_0^6 x^2 \times x(6-x)^2 dx$$

$$= \frac{1}{108} \int_0^6 x^3(6-x)^2 dx$$

$$= \frac{1}{108} \int_0^6 (x^3(36 - 12x + x^2)) dx$$

$$= \frac{1}{108} \int_0^6 (36x^3 - 12x^4 + x^5) dx$$

$$= \frac{1}{108} \left[9x^4 - \frac{12}{5}x^5 + \frac{1}{6}x^6 \right]_0^6$$

$$= \frac{1}{108} \left[9(6^4) - \frac{12}{5}(6^5) + \frac{1}{6}(6^6) \right]$$

$$= \frac{1}{108} \left[11664 - 18662.4 + 7776 \right]$$

$$= \frac{1}{108} \left[7776 \right]$$

$$= 7.2$$

8

