


Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until Friday 12th June 2020.



SEMESTER ONE

MATHEMATICS METHODS UNIT 3

2020

SOLUTIONS

Calculator-free Solutions

1. (a) $\frac{d}{dx} \left[\left(\sin \left(\frac{x}{2} \right) \right)^3 \right] = 3 \sin^2 \left(\frac{x}{2} \right) \times \cos \left(\frac{x}{2} \right) \times \frac{1}{2}$ ✓✓
- (b) $2t(\tan t) + \frac{t^2}{\cos^2 t}$ ✓✓
- (c) $f(y) = \cos \left\{ (\sin y)^{\frac{1}{2}} \right\}$
- $f'(y) = -\sin \sqrt{\sin y} \times \frac{1}{2} \left(\sin^{-\frac{1}{2}} y \right) \times \cos y$ ✓✓✓ [7]
2. (a) $v(t) = 2e^{2t} - 2e^2 t + c$ ✓
- $x(t) = e^{2t} - e^2 t^2 + ct + k$ ✓
- When $t = 0, x = 0 \rightarrow k = -1$ ✓
- When $t = 1, x = 0 \rightarrow c = 1$ ✓
- $x(t) = e^{2t} - e^2 t^2 + t - 1$
- ∴
- (b) 0 ✓
- (c) $v(0) = 3$ ✓ [6]
3. (a) $2 \sin \frac{x}{2} - e^{\cos x} + c$ ✓✓
- (b) $f'(y) = (1 - 2y)^{-\frac{1}{2}}$
- $f(y) = \frac{(1 - 2y)^{\frac{1}{2}}}{\left(\frac{1}{2} \right) \times (-2)} + c$ ✓✓
- ∴
- $(-4, 3) \rightarrow 3 = -\sqrt{9} + c \rightarrow c = 6$ ✓
- ∴ $f(y) = -\sqrt{1 - 2y} + 6$ ✓
- ∴
- (c) $g(x) = - \int_{-1}^{2x} \frac{2}{\tan^2 t} dt$ ✓
- $g'(x) = - \frac{2}{\tan^2(2x)} \times 2 = - \frac{4}{\tan^2 2x}$ ✓✓ [9]
- ∴

4. (a) 6 represents $\frac{1}{8}$ of sample then there were 48 responses. ✓
 $\therefore g = 10$ ✓
 $\therefore f = 12$ ✓
 $h = \frac{1}{4}$ ✓
 and ✓
- (b) $E(X) = 1 \times \frac{1}{8} + 2 \times \frac{1}{6} + 3 \times \frac{1}{4} + 4 \times \frac{1}{4} + 5 \times \frac{5}{24}$ ✓
 $\therefore E(X) = 3\frac{1}{4}$ ✓ [6]
5. (a) (i) $\int_{-1}^0 f(x) \, dx - \int_0^2 f(x) \, dx$ ✓✓
 (ii) $\int_u^p [f(x) - g(x)] \, dx$ ✓✓
- (b) $\frac{d}{dx}(f(x)) = \frac{d}{dx}(x^3 - x^2 - 2x) = 3x^2 - 2x - 2$ ✓
 $\therefore 3x^2 - 2x - 2 = 3$ ✓
 $\therefore 3x^2 - 2x - 5 = 0$
 $\therefore (3x - 5)(x + 1) = 0$ ✓
 $\therefore p = \frac{5}{3}$ ✓ [8]
6. (a) (i) $\int_{-4}^5 h(t) \, dt = \int_{-4}^0 h(t) \, dt + \int_0^5 h(t) \, dt$
 $\therefore \int_0^5 h(t) \, dt = 14$ ✓
- (ii) $\int_{-4}^0 [2 - h(t)] \, dt = \int_{-4}^0 2 \, dt - \int_{-4}^0 h(t) \, dt$
 $= 8 - (-4) = 12$ ✓✓
- (iii) $2 + 3 \int_{-4}^5 h(t) \, dt = 2 + 30 = 32$ ✓✓
- (b) We would need to know where $h(t)$ lies below the x-axis. ✓ [6]

7. $(0, 2) \rightarrow e = 2$

$$y' = 4ax^3 + 3bx^2 + cx + d$$

$$\therefore y' = 0 \text{ when } x = 0 \rightarrow d = 0$$

$$y'' = 12ax^2 + 6bx + 2c$$

$$\therefore y'' = 0 \text{ when } x = 0 \rightarrow c = 0$$

$$\text{and } y'' = 0 \text{ when } x = -2 \rightarrow 48a - 12b = 0$$

$$(-2, 0) \rightarrow 0 = 16a - 8b + 2 \rightarrow 16a - 8b = -2$$

$$\text{Hence } b = \frac{1}{2} \text{ and } a = \frac{1}{8}$$

[8]

Calculator-Assumed Solutions

8. (a) $a + b = 26$

$$\frac{30 + 80 + 150 + 20a + 25b}{50} = 16.6$$

$$\therefore a = 16 \text{ and } b = 10$$


(b) Standard deviation = 6.44

(c) (i) $16.6 \left(1 - \frac{d}{100} \right)$

(ii) $41.47 \left(1 - \frac{d}{100} \right)^2$

[9]

9. (a)



(b) (i) $\int_{-2}^2 f'(x) dx = f(2) - f(-2) = 2 - (-2) = 4$

(ii) $\int_{-2}^2 f''(x) dx = f'(2) - f'(-2) = 0 - 0 = 0$

(c)
$$\text{Area} = \int_{-2}^2 |f'(x)| dx = \int_{-2}^2 f'(x) dx \text{ since positive}$$
$$= f(2) - f(-2)$$

$$= 4$$

✓

[10]

10. (a) $2.5 = 2e^{k \times 5}$

✓

$$\therefore k = 0.04463$$

✓

$$\therefore M(t) = 2\,000\,000e^{0.04463t}$$

✓

(b) $M(15) = 2\,000\,000e^{0.04463 \times 15}$

✓

$$= 3\,906\,325 = 3\,906\,500$$

✓

(c) $M'(t) = 89\,260e^{0.04463t}$

✓

$$\therefore M'(15) = 174\,339 \text{ microbes/min}$$

✓

(d) $2\,000\,000 = 3\,906\,500e^{-0.05t}$

✓

$$\therefore t = 13.39 \text{ min}$$

✓

$$\therefore 12:58 \text{ pm}$$

✓

[10]

11. (a) (i) $\text{Length}(x) = \sqrt{x^2 + 16} + \sqrt{(30 - x)^2 + 100}$

✓

$$L'(x) = \frac{1}{2}(16 + x^2)^{-\frac{1}{2}}(2x) + \frac{1}{2}(x^2 - 60x + 1000)^{-\frac{1}{2}}(2x - 60)$$

✓

$$L'(x) = \frac{x}{\sqrt{x^2 + 16}} + \frac{x - 30}{\sqrt{x^2 - 60x + 1000}}$$

✓

$$\therefore \text{Min occurs when } L'(x) = 0$$

✓

$$\frac{x}{\sqrt{x^2 + 16}} = \frac{30 - x}{\sqrt{x^2 - 60x + 1000}}$$

$$\therefore$$

(ii) $x = 8.57 \text{ m}$

✓

(b) $\delta\theta = -0.01$ and $\tan \theta = \frac{21.43}{10} = 2.143 \rightarrow \theta = 1.134$

✓

$$\tan \theta = \frac{30 - x}{10}$$

✓

$$\therefore x = 30 - 10 \tan \theta \rightarrow \frac{dx}{d\theta} = -\frac{10}{\cos^2 \theta}$$

✓

$$\therefore \delta x = -\frac{10}{\cos^2 \theta} \times (-0.01)$$

✓

$$\therefore \text{When } \theta = 1.134, \delta x = 0.559 \text{ m}$$

✓

[10]

12. (a) (i) $x < -2$ or $1 < x < 2$ ✓✓
 (ii) $x = -2, 1, 2$ ✓✓
 (iii) $x < -1.4$ or $x > 1.6$ ✓✓
 (b) $x = -1.4$ or $x = 1.6$ ✓✓ [8]
13. (a) $2\sin(4\pi) = 0$ GL/h ✓
 $W = \int 2\sin\left(\frac{\pi t}{3}\right) dt = -\frac{6}{\pi} \cos\left(\frac{\pi t}{3}\right) + c$
 (b) ✓
 $(0, 3) \rightarrow c = \frac{6}{\pi} + 3 \rightarrow W(t) = -\frac{6}{\pi} \cos\left(\frac{\pi t}{3}\right) + 3 + \frac{6}{\pi}$ ✓
 \therefore At midday $W(12) = 3$ GL ✓
 $\text{Amount} = \int_0^5 2\sin\left(\frac{\pi t}{3}\right) dt = 0.955$ GL ✓✓
 (c) ✓✓
 $W(t) = -\frac{6}{\pi} \cos\left(\frac{\pi t}{3}\right) + 3 + \frac{6}{\pi} = 4 \rightarrow t = 1.03$ ✓
 (d) ✓
 \therefore 1:02 am ✓ [8]
14. (a) DRV and involves success or failure ✓
 (b) (i) $(0.65)^5 = 0.1160$ ✓✓
 (ii) $(0.65)^3 (0.35)^2 = 0.0336$ ✓✓
 (iii) $X \sim B(5, 0.65)$
 $\therefore P(X \geq 2) = 0.9460$ ✓✓
 (iv) $(1 - h)^5$ ✓✓
 (c) $1 - P(0) = 0.99757$
 $\therefore P(0) = 0.00243$
 $(1 - h)^5 = 0.00243$ ✓
 $\therefore h = 0.7$ ✓
 (d) $\text{VAR} = 0.65 \times 0.35 = 0.2275$ ✓
 \therefore St dev = 0.4770 ✓ [13]

15. (a) $f'(x) = \frac{\sqrt{3}}{4} + \frac{1}{2} \cos\left(\frac{x}{2}\right)$ ✓
- TP occurs when $f'(x) = 0$
- $$\frac{1}{2} \cos\left(\frac{x}{2}\right) = -\frac{\sqrt{3}}{4} \rightarrow \cos\left(\frac{x}{2}\right) = -\frac{\sqrt{3}}{2}$$
- ✓
- $$\therefore \frac{x}{2} = \frac{5\pi}{6} \rightarrow x = \frac{5\pi}{3} \text{ km}$$
- ✓✓
- and height = 2.767 m ✓
- (b) Max gradient occurs when $f''(x) = 0$
- $$-\frac{1}{4} \sin\left(\frac{x}{2}\right) = 0$$
- ✓
- $$\therefore x = 0, 6.28, 12.57, 18.85$$
- ✓
- $$\therefore \text{Max gradient} = 0.93$$
- ✓ [8]
16. (a) $f'(x) = \sin x + x \cos x$ ✓✓
- (b) (i) $\int f'(x) dx = \int \sin x dx + \int x \cos x dx$ ✓
- $$f(x) = -\cos x + \int x \cos x dx$$
- ✓
- $$\therefore \int x \cos x dx = f(x) + \cos x + c$$
- ✓
- $$\therefore \int x \cos x dx = x \sin x + \cos x + c$$
- ✓
- (ii) $\int_0^{\pi} x \cos x dx = [x \sin x + \cos x]_0^{\pi}$ ✓
- $$= (0 - 1) - (0 + 1) = -2$$
- ✓
- (c) $\int_0^{\pi} |x \cos x| dx = 3.14$ ✓✓ [9]

17. (a) Uniform ✓
- (b) (i) $\frac{1}{5}$ ✓
- $\frac{\frac{2}{5}}{\frac{3}{5}} = \frac{2}{3}$
- (ii) $\frac{1}{5}$ ✓
- (iii) $\frac{1}{2}(2 + 6) = 4$ ✓
- (iv) $VAR = \frac{1}{12}(6 - 2 + 1)^2 = 2$ ✓
- (c) (i) $\therefore \text{Std Dev} = \sqrt{2}$ ✓
- (ii) $E(Y) = 3 - 2(4) = -5$ ✓
- $\text{Std Dev}(Y) = 2(\text{Std Dev}(X))$
- $\therefore = 2\sqrt{2}$ ✓
- $\therefore VAR(Y) = 8$ ✓ [9]
18. (a) $x(t) = \int e^{\sin 2t} \cos 2t \, dt$
- Since $\frac{d}{dx}(e^{\sin 2t}) = 2e^{\sin 2t} \cos 2t$
- $\therefore x(t) = \int e^{\sin 2t} \cos 2t \, dt = \frac{1}{2} e^{\sin 2t} + c$ ✓✓
- $\rightarrow c = -\frac{1}{2}$
- Since $x(0) = 0$
- $\therefore x\left(\frac{\pi}{4}\right) = \frac{1}{2} e - \frac{1}{2}$ ✓
- $\frac{d}{dt}(v(t)) = 0$ when $t = 0.3331$ ✓
- (b) $v(t) = 0$ when $t = 0.7854$
- $\therefore \text{Total distance} = \int_0^{0.3331} e^{\sin 2t} \cos 2t \, dt = 0.428$ ✓✓ [6]