

METHODS
MATHEMATICS

Calculator-assumed

Student number: In figures

SPIONI

Please place your student identification label in this box

Question/Answer booklet

Semester One Examination, 2018



ALL SAINTS' COLLEGE

ALL SAINTS' COLLEGE

Important note to candidates
No other items may be taken into the examination room. It is your responsibility to ensure that you do not have any unauthorised material if you have any unauthorised material with you, hand it to the supervisor before reading any further.

Important note to candidates

Special items: drawing instruments, templates, notes on two unfolded sheets of A4 paper, and up to three calculators approved for use in this examination

10 to be Provided By the Candidate
Standard items: pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

This Questionnaire Answer Sheet

Materrials required/recommended for this section

Time Showed To This Section
Reading time before commencing work:
ten minutes
one hundred minutes
Working time:

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	8	8	50	52	35
Section Two: Calculator-assumed	13	13	100	98	65
Total					100

Instructions to candidates

1. The rules for the conduct of examinations are detailed in the school handbook. Sitting this examination implies that you agree to abide by these rules.
2. Write your answers in this Question/Answer booklet.
3. You must be careful to confine your response to the specific question asked and to follow any instructions that are specified to a particular question.
4. Supplementary pages for the use of planning/continuing your answer to a question have been provided at the end of this Question/Answer booklet. If you use these pages to continue an answer, indicate at the original answer where the answer is continued, i.e. give the page number.
5. Show all your working clearly. Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
6. It is recommended that you do not use pencil, except in diagrams.
7. The Formula sheet is not to be handed in with your Question/Answer booklet.

- (a) Describe, with parameters, a suitable probability distribution to model the number of trout in a day's catch. (2 marks)
- (b) Determine the probability that there are fewer trout than fish of other species in a day's catch. (2 marks)

- (c) Calculate the probability that over two consecutive days, a total of exactly 19 trout are caught. (2 marks)

Question 9

Question 10**(8 marks)**

The population of a city can be modelled by $P = P_0 e^{kt}$, where P is the number of people living in the city, in millions, t years after the start of the year 2000.

At the start of years 2009 and 2013 there were 1945000 and 2061000 people respectively living in the city.

- (a) Determine the value of the constant k .

(2 marks)

- (b) Determine the value of the constant P_0 .

(2 marks)

- (c) Use the model to determine during which year the population of the city will first exceed 2500000.

(2 marks)


- (d) Determine the rate of change of the city's population at the start of 2013.

(2 marks)

Markers use only		
Question	Maximum	Mark
9	6	
10	8	
11	6	
12	8	
13	8	
14	7	
15	8	
16	9	
17	7	
18	11	
19	7	
20	6	
21	7	
S2 Total	98	
S2 Wt ($\times 0.6633$)	65%	

Question 11

The graph of $f(x) = \frac{6x+1}{5x+1}$ is shown below.



(6 marks)

(1 mark)

Two rectangles are also shown on the graph, with dotted lines, and they both have corners just touching the curve. The smaller is called the inscribed rectangle and the larger is called the circumscribed rectangle.

(a) Complete the missing values in the table below.

x	$f(x)$	$\frac{8}{3}$	$\frac{7}{2}$	$\frac{13}{3}$
0	0	0.5	1	1.5

(b) Complete the table of areas below and use the values to determine a lower and upper bound for $\int_0^2 f(x) dx$.

x interval	0 to 0.5	0.5 to 1	1 to 1.5	1.5 to 2
Area of inscribed rectangle	$\frac{3}{4}$			

(c) Explain how the bounds you found in (b) would change if a larger number of smaller intervals were used. (1 mark)

Question 12**(8 marks)**

A fairground shooting range charges customers \$6 to take 9 shots at a target. A prize of \$20 is awarded if a customer hits the target three times and a prize of \$40 is awarded if a customer hits the target more than three times. Otherwise no prize money is paid.

Assume that successive shots made by a customer are independent and hit the target with the probability 0.15.

- (a) Calculate the probability that the next customer to buy 9 shots wins

(i) a prize of \$20.

(2 marks)

(ii) a prize of \$40.

(1 mark)

- (b) Calculate the expected profit made by the shooting range from the next 30 customers who pay for 9 shots at the target.

(3 marks)

- (c) Determine the probability that more than 6 out of the next 8 customers will not win a prize.

(2 marks)


Supplementary page

Question number: _____

Three functions are defined by $f(x) = 9e^{-0.07x}$, $g(x) = 3e^{0.09x}$ and $h(x) = 3$.

(8 marks)

Question 13
One of the functions is shown on the graph above. Add the graphs of the other two functions.



(a)

Working to three decimal places throughout, determine the area of the region enclosed by all three functions.

(b)

Working to three decimal places throughout, determine the area of the region enclosed by all three functions.

Question 14

(7 marks)

A fuel storage tank, initially containing 3750 L, is being filled at a rate given by

$$\frac{dV}{dt} = \frac{t^2(240 - 3t)}{500}, 0 \leq t \leq 80$$

where V is the volume of fuel in the tank in litres and t is the time in minutes since filling began.
The tank will be completely full after 80 minutes.

- (a) Calculate the volume of fuel in the tank after half-an-hour. (3 marks)

- (b) Determine the time taken for the tank to fill to three-quarters of its maximum capacity. (4 marks)

METHODS UNIT 3		CALCULATOR-ASSUMED		CALCULATOR-ASSUMED		METHODS UNIT 3	
9							
Question 21	(8 marks)	Question 15	(5 marks)	The discrete random variable X has a mean of 5.28 and the following probability distribution.	(a) Determine the values of the constants a and b .	(3 marks)	

(iii) $\text{Var}(12 - 5X)$.

(ii) $E(100 - 15X)$.

- (c) Determine
(i) $\text{Var}(X)$.
(ii) $E(X)$.


(1 mark)

(1 mark)

(1 mark)

(3 marks)

End of questions




(c)

Sketch the graph of $y = F(x)$ on the axes below.

- (b) Another function, F , is defined as $F(x) = \int_x^0 f(t) dt$ over the domain $0 \leq x \leq 16$.

(b) Determine the value(s) of x for which $F(x)$ has a maximum and state the value of $F(x)$ at this location.

(a) Determine the exact area between the horizontal axis and the curve for $0 \leq t \leq 4$.



The graph of $y = f(t)$ is shown below, where $f(t) = 5 \sin\left(\frac{\pi}{4}t\right)$.

(7 marks)

- (a) Determine the values of the constants a and b .

Question 15
(5 marks)

The discrete random variable X has a mean of 5.28 and the following probability distribution.

(a) Determine the values of the constants a and b .

(3 marks)

Question 16

(9 marks)

A particle starts from rest at O and travels in a straight line.

Its velocity $v \text{ ms}^{-1}$, at time $t \text{ s}$, is given by $v=8t-t^2$ for $0 \leq t \leq 3$ and $v=135t^{-2}$ for $t > 3$.

- (a) Determine the initial acceleration of the particle.

(2 marks)

- (b) Calculate the change in displacement of the particle during the first three seconds.
(2 marks)

- (c) Determine, in terms of t , an expression for the displacement, $x \text{ m}$, of the particle from O for $t > 3$.
(2 marks)

- (d) Determine the distance of the particle from O when its acceleration is -1.25 ms^{-2} .
(3 marks)

Question 20

(6 marks)

The discrete random variable X is defined by

$$P(X=x)=\begin{cases} \frac{2k}{e^{1-x}} & x=0,1 \\ 0 & \text{elsewhere.} \end{cases}$$

- (a) Show that $k=\frac{e}{2+2e}$.

(3 marks)

- (b) Determine, in simplest form, the exact mean and standard deviation of X .
(3 marks)

A random sample of n components are selected at random from a factory production line. The proportion of components that are defective is p and the probability that a component is defective is independent of the condition of any other component.

The hourly cost of fuel to run a train is proportional to the square of its speed and is \$36 per hour when the train moves at a speed of 16 km $^{-1}$. Other costs amount to \$144 per hour, regardless of speed.

(a) Determine the values of n and p . (4 marks)

In the random variable X is the number of faulty components in the sample. The mean and standard deviation of X are 30.6 and 5.1, respectively.

(a) Show that when the train moves at a steady speed of $x \text{ km/h}$, where $x > 0$, the total cost per kilometre, C , is given by

$$C = \frac{9x}{144} + \frac{64}{x}.$$

 (3 marks)

(b) After changes are made to the manufacturing process, the proportion of defective components is now 3 %. Determine the smallest sample size required to ensure that the probability that the sample contains at least one defective component is at least 0.95.

Q3. Calculate the maximum speed of a train to travel 240 km, spending half the time travelling at a constant speed for the entire journey.

(11 marks)

Question 18

The air pressure, $P(h)$ in kPa, experienced by a weather balloon varies with its height above sea level h km and is given by

$$P(h) = 101.5 e^{-0.135h}, 0 \leq h \leq 20.$$

- (a) Determine $\frac{dP}{dh}$ when the height of the balloon is 5.5 km. (2 marks)

- (b) What is the meaning of your answer to (a). (1 mark)

- (d) Determine $\frac{dh}{dt}$ when the height of the balloon is 7.04 km.

(3 marks)

- (e) Determine $\frac{dP}{dt}$ when the height of the balloon is 7.04 km. (3 marks)

The height of the balloon above sea level varies with time t minutes and is given by

$$h(t) = \frac{t^2(120-t)}{12800}, 0 \leq t \leq 80.$$

- (c) Determine the air pressure experienced by the balloon when $t = 48$. (2 marks)