

- the end of week 1 of term 4, Fri October 12th 2018

The release date for this exam and marking scheme is

agreement/licence.

paper or marking guide until the agreed release date stipulated in the purchasing
form of showing them how the work is marked but students are not to retain a copy of the
available to anyone who is not a teacher at the school. Teachers may give feedback to students in
the items and the solutions/markings keys are to be kept confidential and not copied or made
They are not to be shared in any manner with a school which has not purchased their own licence.
are purchased.
The items that are contained in this examination are to be used solely in the school for which they
This examination is Copyright but may be freely used within the school that purchases this licence.

License Agreement

© MAWA, 2018

MATHEMATICS METHODS

MAWA Semester 2 (Unit 3&4) Examination 2018

Calculator-assumed

Marking Key

Section Two: Calculator-assumed**(100 Marks)****Question 8 (a)****(2 marks)**

Solution	
Bernoulli distribution with parameter	
$p = \frac{1}{300}$	
Mathematical behaviours	Marks
• states Bernoulli distribution	1
• states parameter of	1
$p = \frac{1}{300}$	

Question 8 (b)**(2 marks)**

Solution	
$E(X) = \frac{1}{300}$	
Variance(X)	
$= \frac{1}{300} \left(1 - \frac{1}{300} \right)$	
$= \frac{1}{300} \left(\frac{299}{300} \right)$	
$= \frac{299}{90000}$	
Mathematical behaviours	Marks
• states correct mean	1
• states variance	1

<p>Solution</p> <p>$F(x) = 0.75 \Rightarrow 0.75 = \frac{1}{1}(x - 2)$</p> <p>ie $x = 4.25$</p> <p>Hence the upper quartile is 4.25.</p>	(2 marks)
---	---

Question 9 (d)

<p>Marks</p> <p>Mathematical behaviours</p> <ul style="list-style-type: none"> • identifies need to integrate $f(x)$ • determines definite integral using correct limits of integration ($2, x$) • determines $F(x)$ and states it as a piecewise function 	1 1 1
---	---

Question 9 (c)

<p>Marks</p> <p>Mathematical behaviours</p> <ul style="list-style-type: none"> • determines numerator • determines denominator 	1 1
---	---

Question 9 (b)

<p>Marks</p> <p>Mathematical behaviours</p> <ul style="list-style-type: none"> • determines the expected value • determines the variance 	1 1
---	---

Question 9 (a)

<p>Marks</p> <p>Mathematical behaviours</p> <ul style="list-style-type: none"> • calculates n and deduces that sample size is appropriate 	1
--	---

Question 21 (e)

<p>Marks</p> <p>Mathematical behaviours</p> <ul style="list-style-type: none"> • substitutes $p = 0.7$ and equates true E to E for simple interval • solves inequality to determine E to E for simple interval • states 97% level of confidence 	1 1 1
--	---

Question 21 (d)

<p>Marks</p> <p>Mathematical behaviours</p> <ul style="list-style-type: none"> • uses the relationship between the errors to draw valid conclusion 	1
--	---

Question 21 (iii)

Mathematical behaviours	Marks
$0.75 = \frac{1}{3}(x - 2)$	1
• states • solves for x	1

Question 10 (a)

(2 marks)

Solution

$$P = P_0 e^{kt}$$

$$\text{ie } P = 35e^{0.03t}$$

$$P = 50 \Rightarrow 50 = 35e^{0.03t}$$

$$\text{ie } t = 11.88$$

ie the population will reach 50 million in roughly 11 years and 11 months

Mathematical behaviours	Marks
• substitutes $P_0 = 35$ and $P = 50$ to obtain required equation	1
• solves equation to obtain t	1

Question 10 (b)

(2 marks)

Solution

$$\text{In 15 years the country's population} = 35e^{15 \times 0.03}$$

$$\text{In 15 years the city's population} = (0.22)35e^{c \times 15} \text{ where } c \text{ represents its growth rate}$$

$$\text{Hence } (0.22)35e^{c \times 15} = (0.4)35e^{15 \times 0.03}$$

$$\text{Solving gives } c \approx 0.0699$$

Hence the continuous growth rate is approximately 7%.

Mathematical behaviours	Marks
• equates city's population to 40% of country's population in 15 years	1
• solves equation and states percentage growth rate	1

Question 21 (a)

(3 marks)

Solution

$$\hat{p} = \frac{465}{700} \approx 0.6643, E = 1.96 \sqrt{\frac{(0.6643)(1 - 0.6643)}{700}} \approx 0.0350$$

$$\text{CI is } 0.6643 - 0.0350 < p < 0.6643 + 0.0350$$

$$\text{ie } 0.6293 < p < 0.6993$$

So the 95% confidence interval is $0.629 < p < 0.699$

Mathematical behaviours	Marks
• calculates sample proportion correctly	1
• calculate standard error correctly	1
• calculates interval correctly	1

Question 21 (b)

(2 marks)

Solution

Because the old satisfaction rate (65%) lies within the new confidence interval, the recent survey does not provide conclusive evidence that the satisfaction rate has improved.

Mathematical behaviours	Marks
• states that survey is not conclusive	1
• states a valid reason	1

Question 21 (c) (i)

(3 marks)

Solution

$$\text{Since } \hat{p}(1 - \hat{p}) \leq \frac{1}{4}$$

$$E_{95\%} = 1.96 \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} \leq 1.96 \sqrt{\frac{1}{4n}} = \frac{1.96}{2} \left(\frac{1}{\sqrt{n}} \right) < \frac{1}{\sqrt{n}}$$

Mathematical behaviours	Marks
• substitutes $\hat{p}(1 - \hat{p}) = 0.25$	1
• expresses error as a constant $\times \frac{1}{\sqrt{n}}$	1
• deduces error $< \frac{1}{\sqrt{n}}$	1


Question 21 (c) (ii)

(1 mark)

Solution

$$E = \frac{1}{\sqrt{n}}$$

Mathematical behaviours	Marks
• states Margin of error	1

•	<p>indicates an appropriate expression involving an integral to determine required area</p> <p>evaluates x axis values, $\ln 2$ and $\ln 3$ to determine A_1</p> <p>substitutes correct bounds to determine A_2</p> <p>evaluates integral and A_2</p> <p>rearranges expression using log laws and simplifies</p>	1 1 1 1 1 1 1 1
Marks	<p>Mathematical behaviors</p>	3
Total area = $A_1 + A_2$	$ \begin{aligned} &= (\ln 2)(3 - 2) + 3(\ln 3 - \ln 2) - \int_{\ln 3}^{\ln 2} e^x dx \\ &= \ln 2 + 3\ln 3 - 3\ln 2 - \left[e^x \right]_{\ln 3}^{\ln 2} \\ &= \ln 2 + 3\ln 3 - 2\ln 2 - (\ln 2 - \left[e^x \right]_{\ln 2}^{\ln 3}) \\ &= \ln 3 - \ln 2 - \left[e^x \right]_{\ln 2}^{\ln 3} \\ &= \ln 3 - \ln 2 - 1 \\ &= \ln \frac{3}{2} - 1 \end{aligned} $	4
(6 marks)	 <p>Solution</p>	Question 11

SEMESTER 2 (UNIT 3&4) EXAMINATION
CALCULATOR-ASSUMED

SEMESTER 2 (UNIT 3&4) EXAMINATION

Question 12 (a)

Solution	
Confidence interval is $(\hat{p} - E, \hat{p} + E) = (0.53, 0.61)$	
So $\hat{p} = \frac{0.53+0.61}{2} = 0.57$	
Mathematical behaviours	Marks
• Obtains correct answer	1

Question 12 (b)

(2 marks)

Solution	
$E = z_{\alpha} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$	
ie $0.04 = 1.96 \sqrt{\frac{0.57 \times 0.43}{n}}$	
Solving for n gives $n \approx 588.5$	
So the sample size was 589 (approximately)	
Mathematical behaviours	Marks
• Uses $Z_{\alpha} \approx 1.96$	1
• Solves for n and rounds	1

Question 12 (c)

(3 marks)

Solution	
$E = z_{\alpha} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$	
0.07 = $z_{\alpha} \sqrt{\frac{0.57(0.43)}{589}}$	
$z_{\alpha} = 3.43$	
$P(-3.43 < Z < 3.43) = 0.9993964$	
Hence the level of confidence is 99.94%	
Mathematical behaviours	Marks
• Substitutes values into error equation	1
• Solves for z_{α}	1
• States level of confidence to at least 1 decimal place	1

Question 12 (d)

(2 marks)

Solution	
From the confidence interval in 10(c), there is a 99.94% probability that p lies between 0.5 and 0.64, and in particular $p > 0.5$. So the claim is justified.	
Mathematical behaviours	Marks
• States the claim is justified	1
• Gives a valid reason	1

Question 20 (a)

(6 marks)

Solution	
$y(9) = 0 \Rightarrow ae^{-bt} \sin 9c = 0 \Rightarrow 9c = \pi$	
$\Rightarrow c = \frac{\pi}{9} \approx 0.349$	
$\frac{dy}{dt} = -abe^{-bt} \sin ct + ace^{-bt} \cos ct = ae^{-bt} (c \cos ct - b \sin ct)$	
max at $t = 4$, $\Rightarrow \frac{dy}{dt} \Big _{t=4} = 0$	
$ie -abe^{-bt} \sin ct + ace^{-bt} \cos ct = 0$	
$ie -ae^{-bt} (b \sin ct - c \cos ct) = 0$	
$ie b \sin ct - c \cos ct = 0$	
$ie b \sin(4 \times 0.349) - 0.439 \cos(4 \times 0.349) = 0 \Rightarrow b = 0.0616$	
$y(4) = 60 \Rightarrow ae^{-(0.0616 \times 4)} \sin(4 \times 0.349) = 60 \Rightarrow a = 78.0$	
Mathematical behaviours	Marks
• uses (9,0) to conclude $9c = \pi$	1
• solves for c	1
• determines derivative function	1
• equates derivative function to 0 at $t = 4$ and solves for b	1
• uses (4,60) to solve for a	1
• gives all answers correct to 3 significant figures	1

Question 20 (b)

(3 marks)

Solution	
$y\left(t + \frac{\pi}{c}\right) = ae^{-b\left(t + \frac{\pi}{c}\right)} \sin\left(c\left(t + \frac{\pi}{c}\right)\right)$	
$\therefore ae^{-bt} e^{-\frac{b\pi}{c}} \sin(ct + \pi)$	
$\therefore -ae^{-bt} e^{\frac{-b\pi}{c}} \sin ct = -ry(t)$	
Mathematical behaviours	Marks
• replaces t with $t + \frac{\pi}{c}$ to obtain required function	1
• uses indices laws to factor out ae^{-bt}	1
• uses $\sin(ct + \pi) = -\sin ct$ to complete argument	1

Question 19 (a)		Solution	
Question 13 (a)		Solution	
Since velocity and acceleration are opposing one another, the particle is slowing down	Marks	$v(t) = 3t^2 + 2t - 9$ $v(0) = 9$ $a(t) = 6t - 2$ $a(0) = -2$	<ul style="list-style-type: none"> calculates $v(0)$ differentiates $v(t)$ to obtain $a(t)$ and $a(0)$ states particle is slowing down
Mathematical behaviours	Marks	$DB = 60$	<ul style="list-style-type: none"> solves the equation
(3 marks)			(2 marks)

Question 13 (b)		Solution	
(2 marks)			
Mathematical behaviours	Marks	$v(t) = 3t^2 - 2t + 9$	$v(0) = 9$ $a(t) = 6t - 2$ $a(0) = -2$
Since velocity and acceleration are opposing one another, the particle is slowing down	Marks	$DB = 60$	<ul style="list-style-type: none"> solves the equation
(2 marks)			

Question 13 (c)		Solution	
(2 marks)			
Mathematical behaviours	Marks	$s(t) = t^3 - t^2 + 9t + C$	$(0, -1) \Leftrightarrow C = -1$ $\therefore s(t) = t^3 - t^2 + 9t + C$ $\therefore s(5) = 5^3 - 5^2 + 9(5) - 1 = 144$
Hence its final position is 144m from the origin.	Marks	$I_g = 10^{8.5} I_0$	$I_g = 10^{8.5} I_0 = 10^{2.5} \approx 316$ $I_e = 10^6 I_0 = 10^{-2.5} \approx 316$ I_e ie the gunshot is approximately 316 times louder than the restaurant conversation
Mathematical behaviours	Marks	$I_e = 10^6 I_0 = 10^{-2.5} \approx 316$	<ul style="list-style-type: none"> evaluates the ratio $\frac{I_e}{I_g}$ correctly
(1 mark)			

Question 19 (c)		Solution	
(1 mark)			
Mathematical behaviours	Marks	$k = 10^{10} \log_{10}\left(\frac{I}{I_0}\right) \Leftrightarrow 0.1k = \log_{10}\left(\frac{I}{I_0}\right) \Leftrightarrow 10^{0.1k} = \frac{I}{I_0} \Leftrightarrow I = 10^{0.1k} I_0$	<ul style="list-style-type: none"> rearranges log expression correctly
Mathematical behaviours	Marks	$k = 85 \Leftrightarrow I_0 = 10^{0.1 \times 85} I_0 \Leftrightarrow I_0 = 10^{8.5} I_0$	<ul style="list-style-type: none"> substitutes $k = 85$ and states I_0 in terms of I_0
(1 mark)			

Question 19 (b)		Solution	
(2 marks)			
Mathematical behaviours	Marks	$I = 10^{-6} \text{ Watts/m}^2$	<ul style="list-style-type: none"> solves the equation
Mathematical behaviours	Marks	$dB = 60$	$60 = 10 \log_{10}\left(\frac{I}{I_0}\right) \Leftrightarrow 10^{6} = \frac{I}{I_0} \Leftrightarrow I = 10^6 I_0$
(2 marks)			

Question 13 (a)		Solution	
(3 marks)			
Mathematics Methods	Marks	$v(t) = 3t^2 + 2t - 9$	$v(0) = 9$ $a(t) = 6t - 2$ $a(0) = -2$
Calculator-assumed	Marks	$SEMESTER 2 (UNIT 3&4) EXAMINATION$	$SEMESTER 2 (UNIT 3&4) EXAMINATION$
Calculator-assumed	Marks	14	14

**CALCULATOR-ASSUMED
SEMESTER 2 (UNIT 3&4) EXAMINATION
(4 marks)**

Question 14

Solution	
$A(x) = 2xe^{-x^2}$	
$\frac{dA}{dx} = 2(-2x^2e^{-x^2} + e^{-x^2}) = 2e^{-x^2}(-2x^2 + 1)$	
$\frac{dA}{dx} = 0 \Rightarrow x = \pm \frac{1}{\sqrt{2}}$	
$x = \frac{1}{\sqrt{2}} \Rightarrow y = e^{-\frac{1}{2}}$	
Hence C has coordinates $\left(\frac{1}{\sqrt{2}}, e^{-\frac{1}{2}}\right)$	
Mathematical behaviours	Marks
• states $A(x)$	1
• differentiates $A(x)$	1
• solves $\frac{dA}{dx} = 0$	1
• states co-ordinates of C	1

**CALCULATOR-ASSUMED
SEMESTER 2 (UNIT 3&4) EXAMINATION**


Question 18 (a)

Solution	
$y = x \ln x$	
$\frac{dy}{dx} = (\ln x)(1) + x \cdot \frac{1}{x}$	
$\frac{dy}{dx} = \ln x + 1$	
$\frac{d^2y}{dx^2} = \frac{1}{x}$	
For stationary points, $\ln x + 1 = 0$ ie $\ln x = -1$ ie $x = e^{-1}$ $x = e^{-1} \Rightarrow y = -\frac{1}{e}$ $\frac{d^2y}{dx^2} = \frac{1}{e^{-1}} = e > 0$, hence a minimum turning point $\left(\frac{1}{e}, -\frac{1}{e}\right)$	
Mathematical behaviours	Marks
• differentiates correctly using the product rule	1
• equates first derivative to zero to determine x co-ordinate of stationary points	1
• uses second derivative test to determine nature of turning point	1
• determines correct coordinates of turning point	1

Question 18 (b)

(1 mark)

Solution	
Since $\frac{d^2y}{dx^2} = \frac{1}{x} \neq 0$ for any x value. Hence the curve cannot have a point of inflection.	
Mathematical behaviours	Marks
• states second derivative is never zero hence no P.O.I.	1

Question 15 (e)	
Solution	Using CAS, 
	• states $k=3$
	• Mathematical behaviours
	Marks 1

Question 15 (d)	
Solution	$P(X \leq 3)$
	Bincdf(0,3,20,0.101)=0.8634
	• states probability
	Marks 1

Question 15 (c)	
Solution	$P(X \leq 3) =$
	$(20 \choose 0)(0.101)^0(0.899)^{20} + (20 \choose 1)(0.101)^1(0.899)^{19} + (20 \choose 2)(0.101)^2(0.899)^{18} + (20 \choose 3)(0.101)^3(0.899)^{17}$
	• identifies that "no more than 3 prizes" means "can win 0,1,2, or 3 prizes"
	• states correct expression for the probability
	Marks 1

Question 15 (b)	
Solution	$X \sim Bin(20, 0.101)$
	• states Binomial distribution
	• states correct parameters
	Marks 1

Question 15 (a)	
Solution	$Prob\text{ility of winning a prize} = 0.1 + 0.001 = 0.101$
	• uses Addition Principle to calculate the correct probability
	Mathematical behaviours
	Marks 1

SEMESTER 2 (UNIT 3&4) EXAMINATION CALCULATOR-ASSUMED MATHEMATICS METHODS	
9	
	E(T) = $25E(X) - 0.15$
	$= 25(0.2305) - 0.15$
	$= 5.6125$
	$\approx \$5.61$
	$Sd Dev(T) = 25 \times Sd Dev(X)$
	$= 25 \times 0.0294$
	$= 0.735$
	$\approx 74c$
	• states linear transformation required
	• determines mean
	• determines standard deviation
	Marks 1

**CALCULATOR-ASSUMED
SEMESTER 2 (UNIT 3&4) EXAMINATION
(2 marks)**

Question 16 (a) (i)

Solution

$$\int_{-3}^2 f(x) dx = -3 + 4 - 4 = -3$$

Mathematical behaviours

- indicates addition of signed areas
- determines result

Marks

1
1**Question 16 (a) (ii)**

(2 marks)

Solution

Area = $3+4+4=11$

Mathematical behaviours

- expresses the area as a sum of areas
- determines result

Marks

1
1**Question 16 (b)**

(3 marks)

Solution

$$\int_2^0 (x - 2f(x))dx = \int_2^0 x dx + 2 \int_0^2 f(x) dx$$

$$= \left[\frac{x^2}{2} \right]_2^0 + 2(-4)$$

$$= \frac{1}{2}(0 - 4) - 8 = -10$$

Mathematical behaviours

- applies the additivity of integrals to split the integral

Marks

1

$$\int_2^0 (-2f(x))dx = 2 \int_0^2 f(x) dx$$

- applies the linearity of integrals to deduce
- determines result

1

1

Question 16 (c)

(3 marks)

Solution

F(x) $F'(x)=0$

Maximum value of occurs where

**CALCULATOR-ASSUMED
SEMESTER 2 (UNIT 3&4) EXAMINATION**

$$F'(x) = \frac{d}{dx} \int_{-3}^x f(t) dt = f(x)$$

$$f(x) = 0 \Rightarrow x = -2, 0, 2$$

$$F(-2) = -3, F(0) = 1, F(2) = -3$$

Hence, max is 1.

Mathematical behaviours		Marks
• applies the Fundamental Theorem		1
• solves $f(x) = 0$, stating $x = -2, 0, 2$		1
• determines maximum value		1

Question 17 (a)

(1 mark)

Solution			
$D \sim N(65, 4.9^2)$	small	medium	large

$$\text{Proportion of peaches } P(d < 57) = 0.0513 \quad P(57 \leq d \leq 70) = 0.7950 \quad 0.1538$$

Mathematical behaviours		Marks
• determines both probabilities		1

Question 17 (b)

(3 marks)

Solution			
----------	--	--	--

Let X = profit

$$\mu = \sum xp(x)$$

$$= 0.12 \times 0.0513 + 0.23 \times 0.795 + 0.27 \times 0.1538$$

$$= 0.2305$$

$$\approx 23c$$


$$\sigma^2 = \sum (x - \mu)^2 p(x)$$

$$= (0.12 - 0.23)^2 \times 0.0513 + (0.23 - 0.23)^2 \times 0.795 + (0.27 - 0.23)^2 \times 0.1538$$

$$= 0.00087$$

$$\sigma = 0.0294$$

$$\approx 2.9c$$



Mathematical behaviours		Marks
• states a calculation to determine the mean or variance		1
• determines mean		1
• determines standard deviation		1

Question 17 (c)

(3 marks)

Solution	
----------	--