

Copyright for test papers and marking guides remains with *West Australian Test Papers*.
Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.
Test papers should be withdrawn after use and stored securely in the school until 30 November 2018.



SEMESTER TWO

**MATHEMATICS
METHODS
UNITS 1 & 2**

2018

SOLUTIONS

Calculator-free Solutions

1. (a) $\frac{2^{3x}}{2^y} = 2^4 = 16$ ✓✓
- (b) $u^{\frac{7}{2} - (\frac{5}{2} + \frac{1}{2})}$ ✓
- $= u^{\frac{1}{2}} = \sqrt{u}$ ✓ [4]
2. $ab = 15$ and $a + b = 8$
 $a = 3, b = 5$ or $a = 5, b = 3$ ✓
- $abx^2 + (2b + 7a)x + 14 = 15x^2 + cx + 14$
 $2b + 7a = 2(5) + 7(3)$ or $2b + 7a = 2(3) + 7(5)$ ✓
 $c = 31$ or 41 ✓✓ [4]
3. (a) $m = \frac{4}{3}$ ✓
- $-5 = \frac{4}{3}(3) + c$
- $\therefore c = -9$ $\therefore y = \frac{4}{3}x - 9$ ✓
- $-4x + 3y = -27$ ✓
- $-8x + 6y = -14$
- (b) $9x - 6y = 12$
- $x = -2$ $y = -5$ ✓
- $D(-2, -5)$ ✓
- (c) $4(k - 2) - 3(2k - 3) = 7$ ✓
- $-2k + 1 = 7$ ✓
- $k = -3$ ✓ [7]
4. $\frac{1.2 \times 10^{-4}}{3 \times 10^{-7}} = 4 \times 10^2 = 400$ ✓✓ [2]
5. (a) $2(3xy) + 2(3x^2) + 2xy = 32$ ✓
- $3xy + 3x^2 + xy = 16$ ✓
- $3x^2 + 4xy = 16$
- (b) $V = 3x^2y$ and $y = \frac{16 - 3x^2}{4x}$ ✓
- $V = 3x^2 \left(\frac{16}{4x} - \frac{3x^2}{4x} \right)$ ✓
- $V = 12x - \frac{9x^3}{4}$
- (c) $V' = 12 - \frac{27x^2}{4}$
- $12 - \frac{27x^2}{4} = 0$ for stationary point ✓
- $x^2 = \frac{16}{9}$ ✓
- $x = \frac{4}{3}$ (discard $-\frac{4}{3}$) ✓

(d)

$V(x)$	\uparrow	$\frac{4}{3}$	\downarrow
$V'(x)$	$+$	0	$-$

 V has a maximum value

✓✓

[9]

6. (a) $T_{n+1} = T_n + 10 \quad T_1 = -3$

✓✓

(b) $T_2 = -8$

✓

$T_3 = -32$

✓

[4]

7. (a) $y = 10x^2 - 2x^3 - 16x + c$

✓

$3 = 10(2)^2 - 2(2)^3 - 16(2) + c$

✓

$c = 11$

$y = 10x^2 - 2x^3 - 16x + 11$

✓

(b) Gradient of x -axis = 0

$\frac{dy}{dx} = 20(2) - 6(2)^2 - 16 = 0$

✓

 $\therefore m = 0$ Tangent is horizontal and parallel to the x -axis.

✓

[5]

8. (a) Vertical translation 12 units down

✓

Horizontal dilation by factor $\frac{1}{2}$.

✓

(b) $7^x = 7^{2x} - 12$

Let 7^x be k . $k^2 - k - 12 = 0$

✓

$(k - 4)(k + 3) = 0$

✓

$7^x = 4 \quad 7^x \neq -3$

✓

One solution, therefore intersects at only one point.

[5]

9. (a) $y = \frac{x^3}{2} + 2x - 7$

✓

$\frac{dy}{dx} = \frac{3}{2}x^2 + 2$

✓

$m = 8$


✓

$1 = 8(2) + c$

$y = 8x - 15$

✓

(b)




✓✓

[6]

10. (a) Period = 4π
Amplitude = 2
(b)

✓
✓



✓✓✓

[5]

Calculator-assumed Solutions

11. (a) The number of phones she is given to repair for the week.
(b) She fixes 23 per day, for 4 days $23 \times 4 = 92$ phones
 $\frac{108}{23} = 4.69565$
(c) 23 days
 $0.69565 \times 8 = 5.5652$ hours
5 hours and 34 minutes
(d) $T_1 = 155$ $T_2 = 128$ $T_3 = 99$ $T_4 = 68$
 $T_{n+1} = T_n - (25 + 2n)$ $T_0 = 180$

✓
✓

✓

✓

✓✓

[6]

12. (a) $x^2 + y^2 = 4$
(b) $\tan \theta = \sqrt{3} \therefore \theta = \frac{\pi}{3}$ radians
radius = 2 units
 $OB = \frac{2}{\cos \frac{\pi}{3}} = 4$
(c)

✓

✓

✓

Therefore area of $\triangle AOB = \frac{1}{2}(4)\sqrt{3} = 2\sqrt{3}$ units²

✓

Area of sector = $\frac{1}{2}r^2\theta = \frac{1}{2}(2)^2\left(\frac{\pi}{3}\right)$
 $= \frac{2\pi}{3}$ units²

✓

Area of shaded part = triangle – sector

$= 2\sqrt{3} - \frac{2\pi}{3} = 1.37$ units² (3 sig fig)

✓

[6]

13. (a) 5.1 seconds $v(t) = -9.8t + 25$ ✓
 (b) $v(0) = 25$ ✓✓
 (c) Maximum height is 31.89 m when $t = 2.55$ sec
 $\frac{31.89 \times 2}{2.55 \times 2} = 12.5$ m/s ✓✓

[5]

14. (a) $f(x) = (2 + x)^2 = 4 + 4x + x^2$ ✓
 $\lim_{h \rightarrow 0} \frac{(2 + x + h)^2 - (2 + x)^2}{h} = 2x + 4$ ✓
 (b) (i) -7.5 ✓
 (ii) 18 ✓
 (c) $p'(x) = 3x^2 - 3a$ ✓
 $0 = 3(\sqrt{2})^2 - 3a$ ✓
 $a = 2$ ✓
 $-\sqrt{2} = (\sqrt{2})^3 - 3(2)(\sqrt{2}) + b$ ✓
 $b = 3\sqrt{2}$ ✓

[7]

15. (a) $3x^2 - \frac{4}{x^2} - 11 = 0$
 $x = -2$ or $x = 2$ ✓✓

(b)

		-2		2	
y'	+	0	-	0	+
y	↑	-	↓	-	↑

✓✓


 $x = -2$ Maximum $x = 2$ Minimum

✓

- (c) $y = x^3 - 11x + \frac{4}{x} + c$ ✓
 $c = 13$ ✓

$y = x^3 - 11x + \frac{4}{x} + 13$ ✓

(d)




✓✓✓

[10]

16. (a) (i) $T_{100} = 23 + (99)(9)$ ✓
 $= 914$ ✓
- (ii) $357980 = \frac{n}{2}(23 + 2534)$ ✓
 $n = 280$ ✓
- (b) (i) $T_{n+1} = \frac{1}{4}T_n$ $T_1 = 12$ ✓✓
 $S_{\infty} = \frac{12}{1 - \frac{1}{4}} = 16$
- (ii) ✓✓ [8]
17. (a) 92°C (initial temp of tea) ✓
 22°C (room temp) ✓
- (b) After 4.12 mins and before 7.45 mins ✓✓
 $4.12 \leq t \leq 7.45$
- (c) Horizontal asymptote $y = 22$ ✓
The tea will cool at a decreasing rate as it approaches room temperature which is 22° . ✓ [6]
18. (a) $y = \frac{2}{x-3} + 2$ ✓✓
- (b) $y = -3\sqrt{x+4}$ ✓✓
- (c) $y = \left(\frac{1}{2}\right)^x - 4$ ✓✓ [6]
19. (a) 0.2 ✓
(b) 0.5 ✓
(c) 0.3 ✓
- (d) $\Pr(X \cap Y) = \Pr(X) \times \Pr(Y)$ if independent ✓
 $\Pr(X \cup Y) = \Pr(X) + \Pr(Y) - \Pr(X \cap Y)$ ✓
 $\Pr(X) + \Pr(Y) - \Pr(X \cap Y) = \Pr(X) \times \Pr(Y)$ ✓
Let $\Pr(Y) = k$
 $0.5 + k - 0.8 = 0.5k$
 $k = 0.6 = \Pr(Y)$ ✓ [6]
20. (a) 1, 3, 7, 15, ... ✓✓
 $T_{n+1} = T_n + 2^n$ $T_1 = 1$
- (b) 20 sets ✓ [3]

21. (a) (i) $y = 16 - 6 - w$
 $y = 10 - w$ ✓
(ii) $y^2 = 36 + w^2 - 12w \cos Y$ ✓
(iii) $(10 - w)^2 = 36 + w^2 - 12w \cos Y$ ✓
 $64 - 20w = -12w \cos Y$ ✓
 $\therefore \cos Y = \frac{5w - 16}{3w}$
- (b) (i) $A = \frac{6w \sin Y}{2} = 3w \sin Y$ ✓
 $A^2 = 9w^2 \sin^2 Y$ ✓
(ii) $9w^2 \sin^2 Y = 9w^2(1 - \cos^2 Y)$ ✓
 $A^2 = 9w^2 \left(1 - \left(\frac{5w - 16}{3w} \right)^2 \right)$ ✓
 $A^2 = -16w^2 + 160w - 256$ ✓
- (c) (i) $A = \sqrt{-16w^2 + 160w - 256}$
 $A' = 0$ when $w = 5$ ✓
Maximum area = 12 units² ✓
(ii) $y = 10 - w = 5$
The triangle is isosceles. ✓ [12]
22. $y' = 3x^2 - 12x + k$ ✓
 $b^2 - 4ac = 0$ for one solution
 $144 - 4(3)k = 0$ ✓
 $k = 12$ ✓ [3]
23. (a) $W = W_0 (1.085)^t$
 $R = R_0 (0.95)^t$
 $10W_0 = R_0$
 $W_0 (1.085)^t = 10 \times W_0 (0.95)^t$ ✓
 $(1.085)^t = 10(0.95)^t$
 $t = 17.329$ years ✓
After 18 years there will be more wallabies. ✓
- (b) $W_{n+1} = 1.085 W_n$ $W_0 = 655$ ✓
 $W_5 = 985$ ✓ [5]
24. (a) $r = -\frac{3}{2}$ ✓
 $\frac{-2 \left(1 - \left(-\frac{3}{2} \right)^n \right)}{1 - \left(-\frac{3}{2} \right)} = 30$ ✓
 $n = 9.003$
 $\therefore 10$ terms ✓
- (b) $T_5 = a + 4d$ and $T_7 = a + 6d$
 $\therefore 2a + 10d = 38$ (eq 1) ✓
 $= 375 = \frac{15}{2}(2a + 14d)$ (eq 2) ✓
 $a = 4$ $d = 3$ ✓
 $S_{30} = 15(2(4) + 29(3)) = 1425$
Sum of next 15 terms = $S_{30} - S_{15} = 1050$ ✓ [7]

25. (a)



- (b) (i) 0.5323
 (ii) 0.2726
 (iii) 0.51
 (iv) 0.8026

✓✓✓✓
 ✓
 ✓
 ✓
 ✓

[7]

26. The particle's initial displacement is 5 m to the right of the origin.

✓

$$v = 3t^2 - 12t \therefore \text{Initial velocity} = 0$$

✓

[2]