# PERTH MODERN SCHOOL

Exceptional schooling. Exceptional students.



INDEPENDENT PUBLIC SCHOOL

# WAEP Semester One Examination, 2018

**Question/Answer booklet** 



OwT goitae
↑ TINU
<b>NETHODS</b>
<b>SOLTAMENTAN</b>

Calculator-assumed

Student number:

ten minutes one hundred minutes	Time allowed for this section seading time before commencing work: Vorking time:
	Your nam
	SDIOW UI

In figures

# Materials required/recommended for this section

To be provided by the supervisor This Question/Answer booklet

Formula sheet (retained from Section One)

To be provided by the candidate

Standard items: pens (blue/black preferred), pencils (including coloured), sharpener, correction fluid/tape, eraser, ruler, highlighters

Special items: drawing instruments, templates, notes on two unfolded sheets of A4 paper, and up to three calculators approved for use in this examination

# Important note to candidates

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised material. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

© 2018 WA Exam Papers. Perth Modern School has a non-exclusive licence to copy and communicate this document for non-commercial, educational use within the school. No other copying, communication or use is permitted without the express written permission other copying, communication of WA Exam Papers, SMO78-112-4.

METHODS UNIT 1 2 CALCULATOR-ASSUMED

# Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	8	8	50	52	35
Section Two: Calculator-assumed	13	13	100	98	65
				Total	100

# Instructions to candidates

- The rules for the conduct of examinations are detailed in the school handbook. Sitting this
  examination implies that you agree to abide by these rules.
- 2. Write your answers in this Question/Answer booklet.
- 3. You must be careful to confine your response to the specific question asked and to follow any instructions that are specified to a particular question.
- 4. Supplementary pages for the use of planning/continuing your answer to a question have been provided at the end of this Question/Answer booklet. If you use these pages to continue an answer, indicate at the original answer where the answer is continued, i.e. give the page number.
- 5. Show all your working clearly. Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
- 6. It is recommended that you do not use pencil, except in diagrams.
- 7. The Formula sheet is not to be handed in with your Question/Answer booklet.

See next page SN078-112-4

CALCULATOR-ASSUMED	10	METHODS UNIT 1
CALCULATUR-ASSUMED	19	METHODS UNIT I

Supplementary page

Question number: \_\_\_\_\_

65% (98 Marks)

8١

Supplementary page

Question number:

Working time: 100 minutes.


Section Two: Calculator-assumed

provided.


#### (4 marks) Question 9

This section has thirteen (13) questions. Answer all questions. Write your answers in the spaces

The graphs of  $y=2x^2+2x+c$ ,  $y=a(x-2)^2+1$  and y=(x+b)(x+3) are shown below.



Determine the values of the constants a,b and  $\varepsilon.$ 



**CALCULATOR-ASSUMED** 

Question 10 (7 marks)

An online cosmetics company is offering new customers the opportunity to select 6 different products for just \$6 each. They can select from a range of 11 different moisturisers, 15 different lipsticks and 9 different eye-shadows.

Determine how many different selections can be made. (2 marks)

Solution	_
$^{35}C_6 = 1623160$	
Specific behaviours	_
✓ indicates choosing 6 from 35	
✓ correct number	

Determine how many different selections can be made that just include moisturisers.

(2 marks)

	Solution
	$^{11}C_6 = 462$
	C <sub>6</sub> — 402
	Specific behaviours
✓	indicates choosing 6 from 11
✓	correct number

Determine the probability that if a new customer randomly makes a selection, it will not include an eye-shadow. (3 marks)

Solution		
11 + 15 = 26		
$^{26}C_6 = 230\ 230$		
$P = \frac{230230}{1623160} = \frac{299}{2108} \approx 0.142$		
$P = \frac{1623160}{1623160} = \frac{2108}{2108} \approx 0.142$		
1025100 2100		
Specific behaviours		
✓ indicates choosing 6 from 26		
✓ forms probability		
✓ simplifies or approximates with decimal		

See next page SN078-112-4 **CALCULATOR-ASSUMED** 17 **METHODS UNIT 1** 

Question 21 (9 marks)

Injury events in a state were grouped as follows: (D) injury events resulting in death; (H) injury events requiring hospitalisation; and (E) injury events requiring treatment at an emergency department. Injury events were further grouped according to whether prior use of alcohol was a contributing factor. The number of injury events for one month is summarised in the table below.

	D	Н	Е	Total
Alcohol	25	581	5 605	6 211
No Alcohol	115	4 422	11 912	16 449
Total	140	5 003	17 517	22 660

Calculate the percentage of all injury events that involved prior use of alcohol.

Solution		
$\frac{6211}{20000000000000000000000000000000000$		
$\frac{1}{22660} \times 100 = 27.4\%$		
Specific behaviours		
√ identifies correct totals		
√ correct percentage		

If one injury event was selected at random from the data, determine the probability that

it required hospitalisation.

	$P(H) = \frac{5003}{22660} \approx 0.221$	(1 mark
Ì	Specific behaviours	
	✓ correct probability	

it did not require treatment at an emergency department given that prior use of alcohol was a contributing factor. (2 marks)

$P(E A) = \frac{25 + 581}{6211} = \frac{606}{6211} \approx 0.0976$				
Specific behaviours				
✓ numerator				
✓ denominator				

prior use of alcohol was a contributing factor given that it required hospitalisation

salion.
2 marks

Use your answers above to explain whether the type of injury event (D, H or E) appears to be independent of prior use of alcohol. (2 marks)

Solution
From (a), $P(A) = 0.274$ but from (b)(iii) $P(A H) = 0.116$
and since $P(A) \neq P(A H)$ then events NOT independent.
Chasifia bahayiayra
Specific behaviours
✓ states not independent
✓ explains using existing answers

SN078-112-4 End of questions

**METHODS UNIT 1** 


9 CALCULATOR-ASSUMED

(6 marks) Auestion 11

the angle of elevation to the top of the pole, T, is 18°. A thin pole stands vertically in the middle of a level playing ground. From point  ${\mathbb A}$  on the ground,


angle of elevation to the top of the pole is 15°. From point B, also on the ground but 5.35 metres further from the foot of the pole than A, the

(1 mark)



(2 marks)

Showing use of trigonometry, determine the height of the post.




## (7 marks) Question 20 9١

METHODS UNIT 1

CALCULATOR-ASSUMED

respectively of the unit circles shown below, where  $\theta$  and  $\gamma$  are acute angles. Consider the points with coordinates (p,q) and (r,s) that lie in the first and second quadrants



Determine the following in terms of p, q, r and s, simplifying your answers where possible.

(a) 
$$\tan \theta$$
. (b)  $\frac{q}{q}$  (c)  $\sin(180^{\circ} - \theta)$ . (c)  $\sin(180^{\circ} - \theta)$ . (d)  $\sin(180^{\circ} - \theta)$ . (ii)  $-r$  (iii)  $-r$  (iv)  $-s$  (c)  $-r$  (iv)  $-s$  (d)  $-r$  (iv)  $-s$  (each correct response

(a) 
$$\cos(\gamma - \theta)$$
.  $\cos(\gamma - \theta) = \cos\gamma \cos\theta + \sin\gamma \sin\theta$  (b)  $\cos(\gamma - \theta) = \cos\gamma \cos\theta + \sin\gamma \sin\theta$  (c)  $\cos(\gamma - \theta) = \cos\gamma \cos\theta + \sin\gamma \sin\theta$  (d)  $\cos(\gamma - \theta) = \cos\gamma \cos\theta + \sin\gamma \sin\theta$  (e)  $\cos(\gamma - \theta) = \cos\gamma \cos\theta + \cos\gamma \cos\theta$  (figures)  $\cos(\gamma - \theta) = \cos(\gamma - \theta)$  (figures)  $\cos(\gamma - \theta) =$ 

√ correct expression


**CALCULATOR-ASSUMED** 

**METHODS UNIT 1** 

Question 12 (7 marks)

Let  $f(x) = \frac{4}{3-x}$  and  $g(x) = \frac{1}{x+p} + q$ , where p and q are constants.

The graph of y = g(x) is shown below.



(a) Sketch the graph of y = f(x) on the axes above.

(3 marks)

(2 marks)

(b) Determine the values of p and q.

Solution	
p = 2, $q = -3$	
Specific behaviours	
value of p	
value of a	

(c) Solve the equation f(x) = g(x), giving your solution(s) to one decimal place. (2 marks)

Solution
$x = -1.7, \qquad x = 4.4$
Specific behaviours
✓ one solution
✓ second solution
(Rounding for guidance only but
penalise answers given as coordinates)

 State the minimum altitude of the sun at noon in Melbourne and on which day of the year this occurred. (2 marks)

15

Solution
$\theta_{MIN} = 28.7^{\circ} \text{ on day } 173$
Cassifia habayiayra
Specific behaviours
✓ altitude, correct to 1dp

- (d) Solar panels on the roof of a Melbourne business are designed to meet its entire power needs on cloudless days when the altitude of the sun is at least 36° at noon.
  - (i) Draw a straight line on your graph to represent this requirement. (1 mark)

Solution
See graph - horizontal line $\theta = 36^{\circ}$
Specific behaviours
√ neat, straight line

(ii) Determine the number of days the panels are expected to achieve this aim during 2018, ignoring the possibility of cloud cover. (3 marks)

Solution
$\theta > 36^{\circ} \Rightarrow 126 \le n \le 220$
220 - 126 = 94
365 - 94 = 271  days
Specific behaviours
$\checkmark$ one value of $n$
✓ second value of n
√ maximum number of days

(6 marks) Question 13 L

(a) Determine the equation of the axis of symmetry for the graph of  $y = 3x^2 + 12x + 40$ .

Solution
$$Solution
$$S = -\frac{1}{2x} - \frac{1}{2} - \frac{1}{2} - x$$

$$S = -2$$

$$Specific behaviours
$$Specific behaviours$$

$$Specific behaviours$$

$$Specific behaviours$$

$$Specific behaviours$$

$$Specific contents
$$Solvers$$$$$$$$

(4 marks) Determine the values of the constants a and b. (b) The graph of  $y = ax^2 + bx + 13$  passes through the points (-3, -23) and (4, 5).

Solution
$$-23 = (-3)^2 a - 3b + 13$$

$$-23 = 9a - 3b + 13$$

$$5 = 4^2 a + 4b + 13$$

$$5 = 16a + 4b + 13$$
Solve simultaneously using CAS
$$a = -2, \qquad b = 6$$

$$a = -2, \qquad b = 6$$

$$\sqrt{authority}$$

(12 marks) **4 Restion 19** カレ


can be modelled by the equation During 2018, the altitude of the sun, heta degrees, at noon in Melbourne on the  $n^{th}$  day of the year

$$S.SZ + \left(\frac{(n+101)\pi 8}{1841}\right) \text{nis } S.SZ = \theta$$

(z marks) On the 26th of January, the altitude was 71.4°. Calculate the altitude ten days earlier.

Solution
$$n = 16, \quad \theta = 73.4^{\circ}$$
Specific behaviours
 $\checkmark$  indicates  $n$ 
 $\checkmark$  correct angle

(4 marks) Graph the altitude on the axes below for  $0 \le n \le 365$ .



✓ maximum close to right endpoint √ minimum close to (173,29°)  $^{\circ}$  endpoints,  $\theta \approx 75^{\circ}$ Specific behaviours See Graph Solution


√ smooth curve

**Question 18** 

(6 marks)

Question 14 (10 marks)

A logo with triangular outline ABC contains a shaded segment bounded by the straight line PM and the circular arc PM with centre B and radius BM = 18 cm, as shown below.



Given that  $\angle ABC = \frac{5\pi}{12}$ ,  $\angle BCA = 2\angle BAC$  and M is the midpoint of BC, determine

(a) the size of  $\angle ABC$  in degrees.

(1 mark)

Solution
$\frac{5\pi}{12} \times \frac{180}{\pi} = 75^{\circ}$
Specific behaviours
✓ converts angle

(b) the area of the shaded segment.

(2 marks)

Solution
$$A = \frac{1}{2} (18)^2 \left( \frac{5\pi}{12} - \sin\left(\frac{5\pi}{12}\right) \right) \approx 55.6 \text{ cm}^2$$

## Specific behaviours

- ✓ indicates substitution into segment area formula ✓ evaluates area
- (c) the perimeter of the shaded segment.

(3 marks)

# Solution $PM_{arc} = 18 \times \frac{5\pi}{12} = \frac{15\pi}{2} \approx 23.56$ $b = \sqrt{18^2 + 18^2 - 2(18)(18)\cos 75} \approx 21.92$


Perimeter = 
$$23.56 + 21.92 \approx 45.5$$
 cm

## Specific behaviours

- ✓ calculates arc length
- ✓ indicates use of cosine rule to find PM
- ✓ evaluates *PM* and states perimeter

Part of the graph of y = f(x) is shown below, where  $f(x) = -2(x - b)^3 + c$ , and b and c are constants.

13



(i) State the degree of f(x).

Solution
3
Specific behaviours
✓ correct degree

Determine the value of b.

(1 mark)

(1 mark)

Solution
b = 4
Specific behaviours
√ correct value

(iii) Determine f(0).

(2 marks)

Solution
$f(x) = -2(x-4)^3 + 2$
_
$f(0) = -2(-4)^3 + 2 = 130$
Specific behaviours
✓ indicates value of c
✓ evaluates

(b) Another function is given by g(x) = f(x + 8).

Describe how to obtain the graph of y = g(x) from the graph of y = f(x). (2 marks)

Solution
Translate graph 8 units to the left.
Specific behaviours
✓ uses translation
✓ indicates distance and direction

the area of triangle ABC. (4 marks) 6

27 - 081 = 32 + 62Solution

$$2S = A\Delta = 10S = A\Delta + A\Delta$$

$$\frac{81 \times 2}{81 \times 3} = \frac{3A}{87 \text{ nis}}$$

$$89.09 = 34$$

Area = 
$$\frac{1}{2}(36)(60.63)$$
 sin(2 × 35)  $\approx 1025$  cm<sup>2</sup>

## Specific behaviours

- ✓ evaluates second angle and indicates use in sin rule √ indicates use of equation to find second angle
- √ evaluates triangle area √ evaluates second side

(9 marks) Auestion 17 15

probabilities, using any additional information only within that part of the question. For two events, X and Y, it is known that P(X)=0.64 and P(Y)=0.55. Determine the following

(1 mark) (a) P(X).

 $80.0 = 10.0 - 1 = (\overline{X})^{1}$ Solution

√ correct probability Specific behaviours

 $.81.0 = (\overline{X} \cap \overline{Y}) \text{ when } P(Y \cap \overline{X})$  (d) (2 marks)

√ correct probability  $\checkmark$  calculates  $P(X \cap Y)$ Specific behaviours  $72.0 = 78.0 - 40.0 = (\overline{Y} \cap X)q$  $75.0 = 81.0 - 25.0 = (Y \cap X)q$ Solution

(3 marks)  $P(X \cup Y)$  when X and Y are independent.

Specific behaviours 888.0 = $225.0 - 25.0 + 40.0 = (Y \cup X)q$  $225.0 = 25.0 \times 40.0 = (Y \cap X)q$ Solution

√ correct probability  $\checkmark$  indicates use of addition rule  $\checkmark$  uses independence to calculate  $P(X \cap Y)$  ∨

P(Y|X) when P(X|Y) = 0.96. (3 marks)

 $\checkmark$  calculates  $P(X \cap Y)$ √ indicates use of conditional probability rule Specific behaviours  $2.28.0 = \frac{822.0}{4.0.0} = (X|Y)q$  $822.0 = 69.0 \times 22.0 = (Y \cap X)q$ Solution

√ correct probability

See next page See next page 5-211-870NS 5-211-870NS

10

**CALCULATOR-ASSUMED** 

**Question 15** (9 marks)

In a group of 150 students, it is known that 40% are in Year 11 and the rest in Year 12. 60% of those in Year 11 and 50% of those in Year 12 study maths, while the remainder do not.

Let E be the subset of students in Year 11 and M be the subset of students who study maths.

Use set language and notation to describe the subset of students who

mark)
mark)

- Determine the probability that a randomly chosen student from the group
  - is in Year 12 and studies math.

(1 mark)

Solution	
$P(\bar{E} \cap M) = 0.6 \times 0.5 = 0.3 \ \left(\equiv \frac{3}{10}\right)$	
Specific behaviours	
√ correct probability	

studies math.

(2 marks)

Solution
$P(E \cap M) + P(\bar{E} \cap M) = 0.4 \times 0.6 + 0.3 = 0.24 + 0.3 = 0.54 \ \left( \equiv \frac{27}{50} \right)$
Specific behaviours
✓ indicates suitable method
√ correct probability

is in Year 12, given that they study math.

(2 marks)

Solution
$P(\bar{E} M) = \frac{0.3}{0.54} = \frac{5}{9} \ (\approx 0.5556)$
Specific behaviours
✓ uses (i) and (ii)
✓ expresses <i>P</i> correctly (no need to fully simplify)

is in Year 11 or studies math.

(2 marks)


Solution 
$$P(E) + P(M) - P(E \cap M) = 0.4 + 0.54 - 0.24 = 0.7 \quad \left( \equiv \frac{7}{10} \right)$$
Specific behaviours

✓ indicates suitable method
✓ correct probability

See next page SN078-112-4

11 Question 16 (6 marks)

The graph of y = f(x) is shown in bold below. The graphs of y = -f(x), y = f(x + p)and y = f(x) + q are also shown, where p and q are constants.



Clearly label the remaining graphs with y = -f(x), y = f(x + p) or y = f(x) + q. (3 marks)

The one-to-one relation y = 7 - 3x has domain and range given by  $\{x: x = -2, 3, a\}$  and  $\{y: y = -8, -2, b\}$  respectively. Determine the values of constants a and b.

Solution		
x = 3, y = -2		
x = -2, y = 13 = b		
$x = a, y = 7 - 3a = -8 \Rightarrow a = 5$		
Specific behaviours		
✓ value of b		
√ indicates a is mapped onto −8		
✓ solves for value of a		