

Mathematics Methods Units 3 & 4 Solutions					
20. (a) Binomial (100, 0.02)	$n = np = 2$				
	$\sigma = \sqrt{np(1-p)} = \sqrt{0.98} = 1.4$				
	Copyright for test papers and marking guides remains with West Australian Test Papers.				
	Test papers may only be reproduced within the purchasing school according to the advertised Conditions of Sale.				
	Test papers should be withdrawn after use and stored securely in the school until Wednesday 11th October 2017.				
(b)	$P(X \geq 5) = 1 - P(X \leq 4) = 0.0508$	✓			
(c)	$n = 2000, p = 0.02, X = 40$	✓✓			
(d)	$P(X \geq 2) = C_2(0.9)^2(0.1)^0 = 0.243$ or from CAS 90% interval is 0.0149 to 0.0251 from CAS	✓✓			
(e)	$0.0149 \times 2000 \approx 30$	✓✓			
(f)	$0.0251 \times 2000 \approx 50$	✓✓			
	Interval is from 30 to 50.				
	Sample 2 is outside. ( $57 > 50$ )				
	Sample 3 is outside. ( $28 < 30$ )				

# SOLUTIONS

## 2017

### Semester Two

#### MATHEMATICS

#### METHODS

#### UNITS 3 & 4



**Calculator-free Solutions**

1. (a)  $\frac{d}{dx}(e^{\cos x} + 5)$

(b)  $= -\sin x e^{\cos x}$

✓✓

$\int (\sin x \cdot e^{\cos x}) dx$   
 $= - \int (-\sin x \cdot e^{\cos x}) dx$   
 $= -e^{\cos x} + C$

✓✓

[4]

2. (a)  $f'(x) = 2e^{2x} - x$

✓

For max/min,  $2e^{2x} - x = 0$   
 $\therefore 2xe^{2x} - 1 = 0$

✓

$x = 0.5 e^{-2x}$

✓

(b)  $f''(x) = 4e^{2x} + \frac{1}{x^2}$

Since expression > 0 for all x values,  
then stationary point is a minimum.

✓

✓ [5]

3. (a)  $A = \int_0^k (2 - e^{-x}) - x dx$

✓✓

(b)  $A = \left[ 2x - \frac{e^{-x}}{-1} - \frac{x^2}{2} \right]_0^k$   
 $= (2k + e^{-k} - \frac{k^2}{2}) - 1$

✓

✓✓ [5]

4. (a)  $\frac{5x^2}{2} - \frac{\sin 5x}{5} + C$

✓✓

(b)  $\left[ \frac{e^{2x}}{2} - \frac{2x^{1.5}}{3} \right]_0^4$

✓

$= \left[ \frac{e^8}{2} - \frac{16}{3} \right] - \left[ \frac{1}{2} - 0 \right] = 0.5e^8 - \frac{35}{6}$

✓✓

(c)  $2 \sin 2x$

✓✓

[7]

5. (a)  $x = \sin 2t + e^{-2t} + C$   
 $x(0) = 0+1+C=1 \therefore C=0$

✓

(b)  $x = \sin 2t + e^{-2t}$

✓

(c) Assume  $a = -k^2 x$

✓✓

Then  $-4 \sin 2t + 4 e^{-2t} = -k^2(\sin 2t + e^{-2t})$   
This leads to the result that  $k^2 = 4$  and  $k^2 = -4$ .  
Hence, relationship is false.

Or  $a = -4(\sin 2t - 4e^{-2t}) \neq -4(\sin 2t + 4e^{-2t}) = -2^2 x$

✓✓

[7]

**Calculator-free Solutions**

16. (a)  $\frac{1}{3}$  ✓  
 $\frac{1}{3} = \frac{2}{3}$

(b)  $\frac{1}{2}$  ✓✓

(c)  $\mu = 6$  and  $\text{Var}(X) = \int_3^9 \frac{1}{6} (x - 6)^2 dx = 3$   
 $s(x) = \sqrt{3}$  or  $s(X) = \frac{\sqrt{12}}{6}$  by formula  
(d)  $\mu(y) = 2(6) + 5 = 17$   
 $s(y) = 2\sqrt{3}$

[7]

17. (a)  $P(390 < X < 410) = 0.15852$

✓✓

(b)  $P(X < 400 + k) = 0.96$

✓

$\therefore 400 + k = 487.534$

✓

(c)  $m = 810$   $s = 100$

[6]

18. (a)  $\dot{x} = 16 - 4t^3 + t^2$

✓

$\ddot{x} = -12t^2 + 2t = 0$

✓

(b)  $\therefore t = 0$  or  $\frac{1}{6}$

✓

$16 - 4t^3 + t^2 = 0$

✓

$t = 1.68$

✓

(c)  $-12t^2 + 2t$  is a maximum when  $-24t + 2 = 0$

✓

ie when  $t = \frac{1}{12}$

✓

$x = 5.33$

✓

(d)  $\int_0^2 |16 - 4t^3 + t^2| dt = 22.323 \text{ m}$

[8]

19. (a) Normal curve

✓

(b) mean = 21

✓

standard deviation =  $\sqrt{(0.07)(300)(0.93)} = 4.4193$

✓

(c)  $P = 0.07$

✓

standard deviation =  $\sqrt{\frac{(0.07)(0.93)}{300}} = 0.0147$

✓✓

(d) This is a proportion of 0.15

$Z = \frac{0.15 - 0.07}{0.0147} = 5.44$

5.44 standard deviations above the mean is very unlikely.

The testing method may need reviewing.

✓✓

[8]

$\log d$	2.11	2.34	2.69	2.90	3.15
$\log z$	0.0414	0.1139	0.2305	0.3010	0.3802

$\log d$	1.1	1.3	1.7	2.0	2.4
$d$	130	220	490	800	1400

14. (a)  $\log (a) + \log (b) + \log (c) = \log (abc)$

6. (a)  $\log \left( \frac{a}{b} \cdot \frac{b}{c} \cdot a \right) = \log \left( \frac{a^2 b}{b c} \right) = \log \left( \frac{a^2}{c} \right)$


(b)  $y = 1 - x$   
 $2^x = 3^{1-x}$   
 $x \log 2 = (1-x) \log 3$   
 $x \log 2 + x \log 3 = \log 3$

(c)  $x = \log_2 + \log_3$   
 $x = \log_3$   
 $x = \log_6$  as required  
 $y = 1 - \log_6$

7. (a) (i)  $14$  (ii)  $14 + 6 - 6 = 14$   
 $2 \int_0^{10} f(x) dx = 40$   
 $\int_6^4 f(x) dx = -6$   
 $\int_8^4 f(x) dx = 6$   
 $\therefore a = 4, b = 8, c = 10$

8. (a)  $A = \text{Length} \times \text{Width} = 3x \cos 2x$   
 $A' = 3x \cos 2x$   
 $6x \sin 2x = 3 \cos 2x = 0 \text{ for max/min}$   
 $2x \tan 2x = 1 \text{ as required.}$   
 $\therefore \sin 2x + x \cos 2x > 0 \text{ as required}$

15. (a)  $A = \frac{1}{2} \cdot 5 \sin \theta = 7.5 \sin \theta$   
 $dA = 7.5 \cos \theta$   
 $\therefore \frac{dA}{dt} = \frac{d\theta}{dt} \cdot 7.5 \cos \theta = 0$



(c) Area has reached a maximum value.

$$\therefore \frac{dA}{dt} = \frac{d\theta}{dt} \cdot 7.5 \cos \theta = 0$$

[5]

**Calculator-assumed Solutions**

9. (a) Solve  $0.9 = e^{-2k}$   
 $\therefore k = 0.05268 = 0.0527$  (3 s.f.)

✓✓

(b) Solve  $0.5 = e^{-0.05268 t}$   
 $t = 13.153$

✓✓

Half life is 13.153 years.

 $dM$ 

(c)  $\frac{dM}{dt} = M_0 e^{-kt} \cdot (-k)$

✓

At  $t = 2$ ,  $\frac{dM}{dt} = 20 \cdot e^{-0.0527 \cdot 2} \cdot (-0.0527)$   
 $= -0.9486$  units of mass per year.

✓✓ [7]

10. (a) Only 2 results for each trial—single or married.

✓

(b)  $0.6$

✓

(c)  $\sqrt{\frac{(0.6)0.4}{40}} = \sqrt{0.006} = 0.07746$

✓✓

(d) We can be 95% confident that the true proportion is  $p$   
 where  $0.6 - (1.96)(0.07746) < p < 0.6 + (1.96)(0.07746)$   
 ie  $0.4482 < p < 0.7518$

✓✓

(e)  $n = \frac{(1.96)^2 \times 0.6 \times 0.4}{(0.05)^2}$

✓

$n = 368.8$

✓

Sample size needs to be 369.


✓ [9]

11. (a) From calculator, 0.142

✓

(b)  $\sin x$  has a minimum of  $-1$ . So,  $2 + \sin x$  has a minimum of 1  
 So  $\log(2 + \sin x)$  has a minimum of 0.

✓✓



✓✓✓

(d) By inspecting the graph, all of this curve is above the x axis.

✓

∴ since  $\log \sqrt{2 + \sin x} = \frac{1}{2} \log(2 + \sin x)$

✓

Area =  $0.5(0.142) = 0.071$

✓ [9]

12. (a) Minimum when  $f'(x) = 0$ 

$$\therefore 2e^{2x} - 2ke^{-2x} = 0$$

✓

$$\therefore e^{4x} = k$$

✓

$$\therefore x = \frac{1}{4} \ln k$$

✓

$$\therefore \text{Minimum value is } e^{\frac{1}{2} \ln k} + k e^{-\frac{1}{2} \ln k}$$

✓

$$= \sqrt{k} + \frac{k}{\sqrt{k}} = 2\sqrt{k}$$

✓

$$\therefore \text{Range is } y \geq 2\sqrt{k}$$

✓

$$\delta y \approx \frac{dy}{dx} \cdot \Delta x = (2e^{2x} - 6e^{-2x})(0.01)$$

✓✓

$$= 1.09 \text{ (2 decimal places)}$$

✓

$$(c) f(2) = 54.653$$

✓

$$f(2.01) = 55.755$$

✓

Change is 1.10 (2 decimal places)

✓

[9]

13. (a) Not equally likely outcomes, so biased.

✓

$$(b) \frac{7}{8}$$

✓

$$(c) E(X) = 1.875 \quad \text{Var}(X) = 1.0533^2 = 1.109$$

✓✓

$$(d) P(Y = 4) = {}^5C_4 (0.25)^4 (0.75) = \frac{15}{4^5} = 0.0146$$

✓✓

$$(e) P(\text{five 4s}) = {}^5C_5 \left(\frac{1}{8}\right)^5 \left(\frac{7}{8}\right)^0 = 0.00003$$

✓✓

[8]