

ALL SAINTS' COLLEGE

Semester One Examination, 2018

Question/Answer booklet



MATHEMATICS
METHODS
Section One:
Calculator-free

Time allowed for this section		
Your name		
 ln words		
ln figures	Student number:	

sətunim əvit sətunim (titi)

Materials required/recommended for this section

To be provided by the supervisor

Reading time before commencing work:

This Question/Answer booklet

Formula sheet

To be provided by the candidate

Standard items: pens (blue/black preferred), pencils (including coloured), sharpener,

correction fluid/tape, eraser, ruler, highlighters

Special items: nil

Important note to candidates

No other items may be taken into the examination room. It is **your** responsibility to ensure that you do not have any unauthorised material. If you have any unauthorised material with you, hand it to the supervisor **before** reading any further.

© 2018 WA Exam Papers. All Saints' College has a non-exclusive licence to copy and communicate this document for non-commercial, educational use within the school. No other copying, communication or use is permitted without the express written permission of WA Exam Papers. SM001-115-3.

METHODS UNIT 3 2 CALCULATOR-FREE

Structure of this paper

Section	Number of questions available	Number of questions to be answered	Working time (minutes)	Marks available	Percentage of examination
Section One: Calculator-free	8	8	50	52	35
Section Two: Calculator-assumed	13	13	100	98	65
				Total	100

Instructions to candidates

- 1. The rules for the conduct of examinations are detailed in the school handbook. Sitting this examination implies that you agree to abide by these rules.
- 2. Write your answers in this Question/Answer booklet.
- 3. You must be careful to confine your response to the specific question asked and to follow any instructions that are specified to a particular question.
- 4. Supplementary pages for the use of planning/continuing your answer to a question have been provided at the end of this Question/Answer booklet. If you use these pages to continue an answer, indicate at the original answer where the answer is continued, i.e. give the page number.
- 5. Show all your working clearly. Your working should be in sufficient detail to allow your answers to be checked readily and for marks to be awarded for reasoning. Incorrect answers given without supporting reasoning cannot be allocated any marks. For any question or part question worth more than two marks, valid working or justification is required to receive full marks. If you repeat any question, ensure that you cancel the answer you do not wish to have marked.
- 6. It is recommended that you do not use pencil, except in diagrams.
- 7. The Formula sheet is not to be handed in with your Question/Answer booklet.

See next page SN001-115

CALCULATOR-FREE	11	METHODS UNIT 3
CALCULATOR-FREE	11	METHODS UNIT 3

Supplementary page

Question number: _____

32% (25 Marks)

Section One: Calculator-free

This section has **eight (8)** questions. Answer **all** questions. Write your answers in the spaces provided.

Working time: 50 minutes.

Question 1 (6 marks)

A box contains five balls numbered 2,3,4,5 and 6. Three balls are randomly drawn from the box at the same time and the random variable X is the largest of the three numbers drawn.

(a) By listing all possible outcomes (234, 235, etc.), determine $P(X \le 5)$.

Solution Solution $P(X \le 5) = \frac{4}{10}$ Specific behaviours $P(X \le 5) = \frac{4}{10}$ Specific behaviours $\frac{\text{Specific behaviours}}{\text{Vorteet probability}}$

Construct a table to show the probability distribution of X.

		(x=X)	√ values of P
			x values of x
Specific behaviours			
10	10	10	(v-v) 1
9	3	ī	(x=X)d
9	S	₹ v	Х
noitulos			

(c) Calculate $\mathbb{E}\left(X\right)$.


Solution
$$E(X) = \frac{\text{Solution}}{10} = 5.5$$
 Specific behaviours
$$\frac{\text{Specific behaviours}}{\text{indicates products } x \cdot P(X = x)}$$
 \checkmark correct value

Question 8 (7 marks)

0τ

CALCULATOR-FREE

Two houses, P and Q, are 900 m apart on either side of a straight railway line AC. AC is the perpendicular bisector of PQ and the midpoint of PQ is B. A small train, R, leaves station C and



travels towards B, 1200 m from C.

METHODS UNIT 3

Let $\angle PRB = \angle QRB = \theta$, where $0 < \theta < 90^{\circ}$, and X = PR + QR + CR, the sum of the distances of the train from the houses and station.

a) By forming expressions for PR, BR and CR, show that
$$X = 1200 + 450 \frac{(2-\cos\theta)}{\sin\theta}$$
. (3 marks)

Solution
$$PR = \frac{450 \cos \theta}{\sin \theta}, BR = PR \cos \theta = \frac{450 \cos \theta}{\sin \theta}, CR = 1200 - BR = 1200 - \frac{450 \cos \theta}{\sin \theta}$$

$$X = 2 \times \frac{450}{\sin \theta} + 1200 - \frac{450 \cos \theta}{\sin \theta} = 1200 + \frac{450[2 - \cos \theta]}{\sin \theta}$$

$$Specific behaviours
$$Specific behaviours$$

$$Specific behaviours
$$Specific behaviours
$$Specific behaviours$$

$$Specific behaviours
$$Specific behaviours$$

$$Specific beha$$$$$$$$$$$$$$$$$$$$$$$$$$$$

(b) Use a calculus method to determine the minimum value of *X*. (4 marks)

Solution
$$\frac{dX}{d\theta} = 450 \left(\frac{\sin x + \sin x + \cos x + \cos x}{\sin x + \cos x} \right) = 450 \left(\frac{\sin x + \cos x}{\sin x + \cos x} \right) = 450 \left(\frac{\cos x + \cos x}{\sin x + \cos x} \right) = 450 \left(\frac{1 - 2\cos \theta}{\sin x} \right) = 450 \left(\frac{1 - 2\cos \theta}{\sin x} \right) = 450 \left(\frac{1 - 2\cos \theta}{\sin x} \right) = 1200 + 450 \left(\frac{3}{2} \right) \times \frac{2}{\sqrt{3}} = 1200 + 450 \sqrt{3} \text{ m}$$

$$X_{MIN} = 1200 + \frac{450 \left(2 - \cos \theta \right)}{\sin \theta} = 1200 + 450 \left(\frac{3}{2} \right) \times \frac{2}{\sqrt{3}} = 1200 + 450 \sqrt{3} \text{ m}$$

$$\Rightarrow \cos x + \cos x + \cos x + \cos x = 30$$

$$\Rightarrow \cos x + \cos x + \cos x = 30$$

$$\Rightarrow \cos x + \cos x + \cos x = 30$$

$$\Rightarrow \cos x + \cos x + \cos x = 30$$

$$\Rightarrow \cos x + \cos x$$


METHODS UNIT 3

CALCULATOR-FREE

Question 2 (5 marks)

A function defined by $f(x)=13+18x-6x^2-2x^3$ has stationary points at (1,23) and (-3,-41).

(a) Use the second derivative to show that one of the stationary points is a local maximum and the other a local minimum. (3 marks)



$$f''(1) = -12(1) - 12 = -24 < 0 \Rightarrow (1,23)$$
 is a maximum

$$f''(-3) = -12(-3) - 12 = 24 > 0 \Rightarrow (-3, -41)$$
 is a minimum

Specific behaviours

- √ differentiates twice
- \checkmark shows f''(1)<0 and interprets
- ✓ shows and interprets

(b) Determine the coordinates of the point of inflection of the graph of y=f(x). (2 marks)

Solution

$$f''(x)=0 \Rightarrow x=-1$$

$$f(-1)=13-18-6+2=-9$$

$$At(-1,-9)$$

Specific behaviours

- ✓ correct x-coordinate
- √ correct y-coordinate

CALCULATOR-FREE 9 METHODS UNIT 3

Question 7 (9 marks)

The function g is such that $g'(x) = ax^2 + 18x + b$, it has a point of inflection at (-1, 29) and a stationary point at (1, -19).

(a) Determine q(2). (5 marks)


Solution		
g''(x)=2ax+18		
$g''(-1) = -2a + 18 = 0 \Rightarrow a = 9$		
$g'(1)=0 \Rightarrow 9+18+b=0$ b=-27		
$g'(x)=9x^2+18x-27$ $g(x)=3x^3+9x^2-27x+c$		
$g(1)=-19 \Rightarrow 3+9-27+c=-19$ c=-4		
g(2)=24+36-54-4=2		
Specific behaviours		

- √ value of a
- ✓ value of b
- ✓ antiderivative
- ✓ constant of integration
- √ value
- (b) Determine

(i)
$$\int_{1}^{2} g'(x) dx$$

lx.	Solution
	$\int g'(x) = \Delta y = 2 - (-19) = 21$
	Specific behaviours
	✓ uses total change
	✓ correct value

(ii)
$$\int_{1}^{2} 4g'(x) + 16 dx.$$



(2 marks)

See next page SN001-115-3 See next page

(2 marks) Question 3 G

seconds is given by A particle travels in a straight line so that its distance x cm from a fixed point O on the line after t

$$.0 \le 1, \frac{35}{1+12} = x$$

Calculate the acceleration of the particle when t=1.

$$v = \frac{9t^{2}(2t+1)-3t^{3}(2)}{(2t+1)^{2}}, \frac{12t^{3}+9t^{2}}{(2t+1)^{2}}$$

$$a(1) = \frac{(36t^{2}+18t)(2t+1)-3t^{3}(2)}{(2t+1)^{3}} = \frac{3^{2}}{3^{2}} = \frac{9}{9} \text{ cm/s}^{2}$$

$$\sqrt{\cotect \text{ form of quotient rule}}$$

$$\sqrt{\cotect \text{ form of quotient rule}}$$

$$\sqrt{\cotect \text{ correct use of chain rule in second derivative}}$$

- √ correct expression for acceleration
- ✓ substitutes and simplifies

(2 marks) Question 6 8

The height, in metres, of a lift above the ground t seconds after it starts moving is given by

$$h = 2 \cos^2\left(\frac{t}{5}\right)$$
.

Use the increments formula to estimate the change in height of the lift from $t=\frac{\pi}{\delta}$ to $t=\frac{\pi}{\delta}$.

Solution
$$\frac{dh}{dt} = 2 \times 2 \times \cos\left(\frac{t}{S}\right) \sin\left(\frac{t}{S}\right) \left(\cos\left(\frac{t}{S}\right)\right)$$

$$\delta t = \frac{4}{S} \cos\left(\frac{t}{S}\right) \sin\left(\frac{Sn}{S}\right) \left(\frac{Sn}{S}\right) \left(\frac{S$$

See next page


See next page E-9TT-T00NS E-STT-T00NS

6

CALCULATOR-FREE

(8 marks)

Question 4The graph of $y=[3-2x]^3$ is shown below.



(a) Determine the area of the region enclosed by the curve and the coordinates axes.

(4 marks)

Solution		
$3-2x=0 \Rightarrow x=1.5$		
$A = \int_{0}^{1.5} (3 - 2x)^{3} dx \dot{\zeta} \left[\frac{(3 - 2x)^{4}}{-8} \right]_{0}^{1.5} \dot{\zeta}(0) - \left(\frac{81}{-8} \right)$ $\dot{\zeta} \frac{81}{8} \text{ sq units}$		
Specific behaviours		
✓ writes integral with limits		
✓ antidifferentiates		

✓ expression with both limits substituted

√ correct area

(b) Given that the area of the region bounded by the curve, the x-axis and the line x=k is 8 square units, determine the value of k, where 0 < k < 1.5. (4 marks)

t	letermine the value of k , where $0 < k < 1.5$.	
	Solution	
	$A = \int_{k}^{1.5} (3 - 2x)^3 dx \Rightarrow 8 = \left[\frac{(3 - 2x)^4}{-8} \right]_{k}^{1.5}$	
	$8 = (0) - \left(\frac{(3 - 2k)^4}{-8}\right)$	
	$(3-2k)^4 = 64$ $3-2k = \sqrt{8}$	
	$2k=3-2\sqrt{2}$	
	$k = \frac{3}{2} - \sqrt{2}$	
	Specific behaviours	
	✓ equation with antiderivative	
	✓ equation with both limits substituted	

See next page

✓ simplifies equation ✓ value of ' See CALCULATOR-FREE 7 METHODS UNIT 3

Question 5 (7 marks)

Determine $\frac{dy}{dx}$ for the following, simplifying each answer.

(a) $y = \sqrt{5 - 6x}$.

Specific behaviours

✓ indicates use of chain rule

✓ correct derivative, simplified

(b) $y=3x^4\sin(4x)$.

Solution	
$\frac{dy}{dx} = 12x^3 \sin(4x) + 3x^4(4) \cot(4x)$	os(4 x)

$$\frac{12}{6} 12 x^3 \sin(4x) + 12 x^4 \cos(4x)$$

Specific behaviours

- ✓ indicates use of product rule
- \checkmark correct derivative of $\sin(4x)$
- ✓ correct derivative

(c)
$$y = \int_{x}^{2} \frac{1}{\sqrt{1+t}} dt$$
.

(2 marks)

(3 marks)

(2 marks)

$$\frac{dy}{dx} = -\int_{2}^{x} \frac{1}{\sqrt{1+t}} dt \dot{c} - \frac{1}{\sqrt{1+x}}$$

Specific behaviours

- √ reverse limits
- ✓ correct derivative