

Data Management

Table of contents

```
library(tidyverse)
```

```
-- Attaching core tidyverse packages ----- tidyverse 2.0.0 --
```

```
v dplyr      1.1.4      v readr      2.1.5
```

```
v forcats   1.0.0      v stringr    1.5.1
```

```
v ggplot2    3.5.1      v tibble     3.2.1
```

```
v lubridate  1.9.4      v tidyr      1.3.1
```

```
v purrr      1.0.4
```

```
-- Conflicts ----- tidyverse_conflicts() --
```

```
x dplyr::filter() masks stats::filter()
```

```
x dplyr::lag()     masks stats::lag()
```

```
i Use the conflicted package (<http://conflicted.r-lib.org/>) to force all conflicts to become
```

```
library(descr)
```

```
library(knitr)
```

```
library(dplyr)
```

```
library(readr)
```

```
library(readxl)
```

```
file_path <- "/cloud/project/data/data.xls"
```

```
data.xls <- read_excel(file_path)
```

```
options(width=80)
```

```
head(data.xls)
```

```
# A tibble: 6 x 138
```

	RespondentID	Country	A3.1	A3.2	A3.3	A3.4	A3.5	A3.6	A3.7	A3.8	A3.9
	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>
1	1	1	2	3	52	52	4	1	5	12	1
2	2	1	2	5	52	52	3	1	8	2	5
3	3	1	1	4	94	94	4	1	5	2	1
4	4	1	1	2	221	221	1	1	10	3	4

```

5          5          1          1          3          52          52          3          1          8          2          0
6          6          1          1          1          52          52          1          4          9          3          4
# i 127 more variables: A3.10 <dbl>, A5.1 <dbl>, A5.2 <dbl>, `A5.3*` <dbl>,
#   A5.4 <dbl>, A5.5 <dbl>, A6.1 <dbl>, A7.1 <dbl>, A7.2 <dbl>, A9.1 <dbl>,
#   A9.2 <dbl>, `A9.3*` <dbl>, A9.4 <dbl>, A9.5 <dbl>, A9.6 <dbl>, A9.7 <dbl>,
#   `A9.8*` <dbl>, A11.1 <dbl>, A11.2 <dbl>, A11.3 <dbl>, A11.4 <dbl>,
#   A12.1 <dbl>, A12.2 <dbl>, A12.3 <dbl>, A12.4 <dbl>, A13.1 <dbl>,
#   A13.2 <dbl>, A13.3 <dbl>, A13.4 <dbl>, A14.1 <dbl>, A14.2 <dbl>,
#   A14.3 <dbl>, A14.4 <dbl>, A16.1 <dbl>, A16.2 <dbl>, `A16.3*` <dbl>, ...

```

```
freq(as.ordered(data.xls$A3.8), plot = FALSE)
```

```
as.ordered(data.xls$A3.8)
```

	Frequency	Percent	Cum Percent
1	293	14.4264	14.43
2	566	27.8680	42.29
3	87	4.2836	46.58
4	22	1.0832	47.66
5	11	0.5416	48.20
6	10	0.4924	48.70
7	11	0.5416	49.24
8	15	0.7386	49.98
9	55	2.7080	52.68
10	155	7.6317	60.32
11	75	3.6928	64.01
12	731	35.9921	100.00
Total	2031	100.0000	

This frequency table shows the many different religious orientations that a sample included. What this is showing me is how many people are a part of a Catholic, Protestant or any other religious orientations. This will be important as I need to gather a variable in which excludes those who are not religious.

```
freq(as.ordered(data.xls$A3.10), plot = FALSE)
```

```
as.ordered(data.xls$A3.10)
```

	Frequency	Percent	Cum Percent
1	49	2.413	2.413
2	323	15.903	18.316
3	330	16.248	34.564
4	565	27.819	62.383
5	441	21.713	84.097
6	252	12.408	96.504
7	37	1.822	98.326
8	34	1.674	100.000
Total	2031	100.000	

This frequency table shows the levels of education within a sample. It shows me how many people don't have a high school degree to how many people have a doctorates. This will help me relate the info to people who are religious, to their levels of education. I need this info because I will eventually compare the information I gather to how many took COVID-19 as a serious threat.

```
freq(as.ordered(data.xls$`A5.3*`), plot = FALSE)
```

```
as.ordered(data.xls$`A5.3*`)
      Frequency Percent Cum Percent
1           592  29.148      29.15
2           401  19.744      48.89
3           259  12.752      61.64
4           168   8.272      69.92
5           148   7.287      77.20
6           139   6.844      84.05
7           122   6.007      90.05
8            94   4.628      94.68
9            60   2.954      97.64
10           48   2.363     100.00
Total        2031 100.000
```

This frequency table shows the levels in which how many people thought COVID-19 was being exaggerated. It goes from level 1 = Not at all to level 10= Extremely. This will help me see how seriously people took COVID-19 and who underestimated it.

```
freq(as.ordered(data.xls$A31.4), plot = FALSE)
```

```
as.ordered(data.xls$A31.4)
      Frequency Percent Cum Percent
1           874  43.033      43.03
2           121   5.958      48.99
3           120   5.908      54.90
4           185   9.109      64.01
5           184   9.060      73.07
6           195   9.601      82.67
7           352  17.331     100.00
Total        2031 100.000
```

This frequency table shows how many people had frequently put on a mask during COVID-19. It goes from level 1, Never, to level 7, Always. This'll help aid my question on who was taking COVID-19 seriously.

```
freq(as.ordered(data.xls$A31.6), plot = FALSE)
```

```
as.ordered(data.xls$A31.6)
```

	Frequency	Percent	Cum Percent
1	24	1.1817	1.182
2	20	0.9847	2.166
3	36	1.7725	3.939
4	106	5.2191	9.158
5	138	6.7947	15.953
6	496	24.4215	40.374
7	1211	59.6258	100.000
Total	2031	100.0000	

This table shows the frequency in which a person kept a distance of 1.5 meters away from others. The chart goes from level 1, Never, to level 7, Always. This will help me see who didn't wear their masks during COVID-19, which relates to my question of who didn't follow the guidelines provided to everyone across the globe.

```
data.xls$HS_NBELOW <- factor(ifelse(data.xls$A3.10 %in% c("1", "2"), "Yes", "No"))
summary(data.xls$HS_NBELOW)
```

No	Yes
1659	372

I am creating a variable named HS_NBELOW to help me see who have educations from High School and below.

```
data.xls$SOME_CLGNABOVE <- factor(ifelse(data.xls$A3.10 %in% c("3", "4", "5", "6", "7", "8"), "Yes", "No"))
summary(data.xls$SOME_CLGNABOVE)
```

No	Yes
372	1659

I am creating a variable named SOME_CLGNABOVE to help me see who has had some college education all the way to a doctoral degree.

```
data.xls$ONLY_R <- factor(ifelse(data.xls$A3.8 %in% c("1", "2", "3", "4", "5", "6", "7", "8", "9", "10", "11", "12"), "Yes", "No"))
summary(data.xls$ONLY_R)
```

No	Yes
886	1145

I am creating a variable named ONLY_R to only include those of religious orientation.