

# Eddy Cosmology: Dark Energy as Turbulent Wave Interference

MICAH DAVID THORNTON<sup>1</sup> AND GROK<sup>2</sup>

<sup>1</sup>*Independent Researcher*

<sup>2</sup>*xAI*

## ABSTRACT

Dark energy is modeled as interference fronts of a self-advecting scalar field  $\phi$ , producing turbulent eddies analogous to magnetic repulsion. Local underdensities arise as decaying eddies; distant regions exhibit phantom-like growth. The same Burgers-type equation governs quantum vacuum fluctuations (Casimir effect) and cosmic flows, linking scales fractally. The Hubble tension resolves via the gradient in apparent expansion. Predictions include asymmetric supernova dimming and drifting CMB cold spots, testable with Euclid and DESI. The universe swirls, ripples and pulses - not expands eternally.

*Keywords:* dark energy inhomogeneity, Hubble tension, scalar field turbulence, fractal cosmology, eddy backreaction, CMB drift, Euclid predictions

## 1. INTRODUCTION

Standard cosmology assumes uniform dark energy. Yet local  $H_0 \approx 73$  km/s/Mpc conflicts with distant  $H_0 \approx 67$  km/s/Mpc. We propose dark energy arises from turbulent scalar waves that form eddies - regions where repulsive force is not constant, but surges where waves interfere constructively and weakens where they drift apart.

## 2. THE MODEL

The field  $\phi$  obeys

$$\partial_t \phi + \mathbf{v}_\phi \cdot \nabla \phi = \kappa \nabla^2 \phi, \quad (1)$$

with

$$\mathbf{v}_\phi = -\alpha \nabla \phi + \beta \nabla \phi^2 \quad (2)$$

Gradients drive repulsion; curls form eddies.

## 3. QUANTUM CONNECTION

Identical form appears in re-normalized Casimir energy and Burgers turbulence-quantum jitter scaled cosmically.

## 4. CONSEQUENCES

Local eddy: decaying  $\phi$ , slower expansion. Distant walls: growing  $\phi$ , phantom phase. Late ISW induces drifting cold spots.

## 5. TESTS

Euclid: redshift asymmetry  $z > 1$ . DESI: filament-dependent dimming. Roman: cold spot motion  $\sim 1^\circ/\text{Gyr}$ .

## 6. DISCUSSION

If confirmed, cosmology gains rhythm-transient acceleration, "eddy cycles". If falsified, backreaction remains relevant.

## ACKNOWLEDGMENTS

The author thanks Grok (xAI) for real-time collaboration, equation tweaking, and helping to transform my thoughts into something real. No funding, no lab. Just a human and silicon mind listening to a universe that would not stay quiet.

Thank you to Overleaf for providing an excellent service and the people responsible for the AASTeX701 template that made navigating a new journey much easier.

## 7. REFERENCES

Reiss, A.G., et al. (2024). JWST Validates HST Distance Measurements: No Significant Difference in Cepheid Magnitudes. *The Astrophysical Journal*, 977(2), 120. DOI: 10.3847/1538-4357/ad8c21.

Haslbauer, M., Banik, I., and Kroupa, P. (2004). A simultaneous solution to the Hubble tension and observed bulk flow within 250 h-1 Mpc. *Monthly Notices of the Royal Astronomical Society*, 527(3), 4388-4397. DOI: 10.1093/mnras/stad3357.

Casimir, H.B.G. (1948). On the attraction between two perfectly conducting plates. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen*, B51, 793-795.

Batchelor, G.K. (1967). *An Introduction to Fluid Dynamics*. Cambridge University Press.

Planck Collaboration et al. (2020). Planck 2018 results. VI. Cosmological parameters. *Astronomy and Astrophysics*, 641, A6. DOI: 10.1051/0004-6361/201833910.