

Discrete Mathematics

Max Kasperowski

December 16, 2017

Contents

1	Logic	3
1.1	Propositional Logic	3
1.1.1	Definitions	3
1.1.2	Properties	4
1.1.3	Equivalence Proof	4
1.2	Predicate Logic	5
1.2.1	Definitions	5
1.2.2	Properties	5
1.3	Logical Proofs	6
1.3.1	Rules of Inference	6
1.3.2	Inference Proof	6
2	Set Theory	7
2.1	Definitions	7
2.1.1	Describing Sets	7
2.1.2	Special Sets	7
2.1.3	Set Equality	7
2.1.4	Subsets	8
2.1.5	Set Cardinality	8
2.1.6	Power Sets	8
2.2	Tuples	8
2.2.1	Cartesian Product	8
2.3	Operations	9
2.4	Properties	10
3	Relations	11
3.1	Definitions	11
3.2	Properties	11
3.3	Operations	11
3.4	Representing Relations	11
3.5	Closures	11
3.6	Equivalence Relations	11
3.7	Equivalence Classes and Partitions	11
3.8	Comparability	11
3.9	Functions	11
3.9.1	Definitions	11
3.9.2	Properties	11
3.9.3	Types of Functions	12

3.9.4	Inverse Function	13
3.9.5	Composition	13
3.10	Relational Algebra	13
4	Graphs	13
4.1	Trees	13

1 Logic

1.1 Propositional Logic—Zeroth Order Logic

In propositional logic, propositions are denoted by letters (p, q) and are formed by connecting other propositions using logical connectives. Propositions can either be true (T) or false (F).

Logical Connectives The logical connectives listed below are the basic connectives available in propositional logic in order of their precedence. Below are the truth tables corresponding to each of the connectives.

1. \neg , not
2. \wedge , and, $\bigwedge_{i=1}^n p_i$
3. \vee , or, $\bigvee_{i=1}^n p_i$
4. \rightarrow, \Rightarrow , implies (only if) defined as: $p \rightarrow q \equiv \neg p \vee q$
5. $\leftrightarrow, \Leftrightarrow$, is equivalent to (if and only if, iff) defined as: $p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$

Negation

p	$\neg p$
T	F
F	T

Logical Or

p	q	$p \vee q$
T	T	T
T	F	T
F	T	T
F	F	F

Equivalence

p	q	$p \leftrightarrow q$
T	T	T
T	F	F
F	T	F
F	F	T

Logical And

p	q	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

Implication

p	q	$p \rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

1.1.1 Definitions

Converse, Contrapositive, Inverse When given the proposition $p \rightarrow q$, $q \rightarrow p$ is its converse, $\neg q \rightarrow \neg p$ is its contrapositive and $\neg p \rightarrow \neg q$ is its inverse. The contrapositive is equivalent to the original proposition and the converse and inverse are also equivalent.

Tautology A proposition that is always true ($p \vee \neg p$).

Contradiction A proposition that is always false ($p \wedge \neg p$).

Contingency A proposition that is neither a tautology nor a contradiction.

Logical Equivalence p and q are logically equivalent if $p \leftrightarrow q$ is a tautology. The notation for equivalence is typically \equiv .

1.1.2 Properties

De Morgan's Laws

$$\neg(p \vee q) \equiv \neg p \wedge \neg q$$

$$\neg(p \wedge q) \equiv \neg p \vee \neg q$$

Identity Laws

$$p \vee F \equiv p$$

$$p \wedge T \equiv p$$

Domination Laws

$$p \vee T \equiv T$$

$$p \wedge F \equiv F$$

Idempotent Laws

$$p \vee p \equiv p$$

$$p \wedge p \equiv p$$

Negation Laws

$$p \vee \neg p \equiv T$$

$$p \wedge \neg p \equiv F$$

Commutative Laws

$$p \vee q \equiv q \vee p$$

$$p \wedge q \equiv q \wedge p$$

Associative Laws

$$(p \vee q) \vee r \equiv p \vee (q \vee r)$$

$$(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$$

Distributive Laws

$$p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$$

$$p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$$

Absorption Laws

$$p \vee (p \wedge q) \equiv p$$

$$p \wedge (p \vee q) \equiv p$$

1.1.3 Equivalence Proof

This is an example of how to perform an equivalence proof. The aim is to show that $\neg(p \vee (\neg p \wedge q))$ is logically equivalent to $\neg p \wedge \neg q$. We can prove this by forming a series of logical equivalences.

$\neg(p \vee (\neg p \wedge q)) \equiv \neg p \wedge \neg(\neg p \wedge q)$	2 nd De Morgan's law
$\neg p \wedge \neg(\neg p \wedge q) \equiv \neg p \wedge (p \vee \neg q)$	1 st De Morgan's law
$\neg p \wedge (p \vee \neg q) \equiv (\neg p \wedge p) \vee (\neg p \wedge \neg q)$	Associative law
$(\neg p \wedge p) \vee (\neg p \wedge \neg q) \equiv F \vee (\neg p \wedge \neg q)$	Negation law
$F \vee (\neg p \wedge \neg q) \equiv \neg p \wedge \neg q$	Identity law

1.2 Predicate Logic —First Order Logic

Predicate logic uses quantified variables over non-logical objects and allows the use of sentences that contain variables. This allows a generalisation of propositions for a set of variables from a domain.

1.2.1 Definitions

Predicates A predicate is a generalisation of propositions when the variable x is replaced by a specific element from its domain. $P(x)$ becomes a proposition. When no other domain is specified the domain is U .

Quantifiers Quantifiers are used to express that a proposition is true for all elements of the domain and that there exists some element in the domain for which it is true. They also have the highest precedence among the logical operators.

Universal quantifier	$\forall x P(x)$	$P(x)$ is true for every x in U
Existential quantifier	$\exists x P(x)$	$P(x)$ is true for some x in U

1.2.2 Properties

Uniqueness Quantifier The uniqueness quantifier is a commonly used quantifier to express that there is only one x for which $P(x)$ is true. It is usually written as $\exists!$ or \exists_1 .

$$\exists_1 x P(x) \equiv \exists x (P(x) \wedge \forall y (P(y) \rightarrow y = x))$$

De Morgan's Laws De Morgan's laws for quantifiers state that $P(x)$ is not true for all x if and only if there exists an x for which $P(x)$ is false and furthermore that if $P(x)$ is false for all x if and only if there does not exist an x for which $P(x)$ is true.

$$\begin{aligned}\neg \forall x P(x) &\equiv \exists x \neg P(x) \\ \forall x \neg P(x) &\equiv \neg \exists x P(x)\end{aligned}$$

1.3 Logical Proofs

Using logical inference it is possible to build an argument given a set of premises to reach a logical conclusion. An argument is valid if truth of all premises p_i implies that the conclusion q is also true.

$$\left(\bigwedge_{i=1}^n p_i \right) \rightarrow q \equiv T$$

1.3.1 Rules of Inference

Modus Ponens

$$\frac{p \rightarrow q \quad p}{\therefore q}$$

Addition

$$\frac{p}{\therefore p \vee q}$$

Universal Instantiation

$$\frac{\forall x P(x)}{\therefore P(c)}$$

Modus Tollens

$$\frac{p \rightarrow q \quad \neg q}{\therefore \neg p}$$

Simplification

$$\frac{p \wedge q}{\therefore p} \quad \frac{p \wedge q}{\therefore q}$$

Universal Generalisation

$$\frac{P(c) \text{ for an arbitrary } c}{\therefore \forall x P(x)}$$

Existential Instantiation

$$\frac{\exists x P(x)}{\therefore P(c) \text{ for some element } c}$$

Hypothetical Syllogism

$$\frac{p \rightarrow q \quad q \rightarrow r}{\therefore p \rightarrow r}$$

Conjunction

$$\frac{p \quad q}{\therefore p \wedge q}$$

Existential Generalisation

$$\frac{P(c) \text{ for some element } c}{\therefore \exists x P(x)}$$

Disjunctive Syllogism

$$\frac{p \vee q \quad \neg p}{\therefore q}$$

Resolution

$$\frac{\neg p \vee r \quad p \vee q}{\therefore q \vee r}$$

Universal Modus Ponens

$$\frac{\forall x (P(x) \rightarrow Q(x)) \quad P(a) \text{ for a particular element } a}{\therefore Q(a)}$$

1.3.2 Inference Proof

All men are mortal. Socrates is a man. Prove that Socrates is mortal.

$P(x) \equiv T$ means that x is a person. $M(x) \equiv T$ means that x is mortal. s is the particular element Socrates.

1. $\forall x P(x) \rightarrow M(x)$ premise
2. $P(s)$ premise
3. $P(s) \rightarrow M(s)$ UI (1)
4. $M(s)$ Modus Ponens, (2) & (3)

2 Set Theory

A set is a collection of distinct elements without duplicates and the order of elements is unimportant.

2.1 Definitions

Containment $a \in A$ denotes that a is an element of the set A , whereas $a \notin A$ denotes that a is not contained in A . A set may contain other sets.

2.1.1 Describing Sets

Roster Method List all elements contained in the set. $S = \{a, b, c, d\}$ is equivalent to both $S = \{d, c, a, b\}$ and also $S = \{a, a, b, c, d, d\}$. When the pattern is clear, (...) may be used as in $S = \{a, b, c, \dots, z\}$.

Set-Builder Method Specify the proposition that all members must satisfy. $S = \{x|P(x)\}$ denotes that all elements x in S must satisfy the proposition $P(x)$.

Interval Notation The elements in a set can be constrained by an open or closed interval. $[a, b] = \{x|a \leq x \leq b\}$ represents the closed interval between a and b . $(a, b) = \{x|a < x < b\}$ represents the open interval between a and b . Additionally combinations like $[a, b)$ and $(a, b]$ are possible.

2.1.2 Special Sets

Universal Set The universal set, denoted by U , contains everything currently under consideration.

Empty Set The empty set is the set that contains no elements. It is denoted by \emptyset . It is important to note that $\emptyset = \{\}$, but $\emptyset \neq \{\emptyset\}$. A set containing the empty set is not the equal to the empty set.

Singleton A singleton is a set containing exactly one element.

2.1.3 Set Equality

Two sets are equal if they contain the same elements. If A and B are sets then their equality is denoted as $A = B$.

$$\forall x(x \in A \equiv x \in B) \rightarrow A = B$$

2.1.4 Subsets

Set A is a subset of B if all elements in A are also in B . This relation is denoted as $A \subseteq B$.

$$\forall x(x \in A \rightarrow x \in B) \equiv A \subseteq B$$

Every set is always a subset of itself and the empty set \emptyset is a subset of every other set.

Proper Subset When $A \subseteq B$, but $A \neq B$ then A is a proper subset of B , which is denoted as $A \subset B$.

$$\forall x(x \in A \rightarrow x \in B) \wedge \exists x(x \in B \wedge x \notin A) \equiv A \subset B$$

2.1.5 Set Cardinality

For finite sets the set cardinality describes the number of distinct elements in a set. It is denoted as $|S|$. The cardinality of the empty set is zero ($|\emptyset| = 0$). If A is the set containing all the letters of the english alphabet, then $|A| = 26$.

2.1.6 Power Sets

The set of all subsets of a set A is called the power set of A and is denoted as $\mathcal{P}(A)$. If $|A| = n$, then $|\mathcal{P}(A)| = 2^n$. For example, given the set $A = a, b$ we get the power set $\mathcal{P}(A) = \{\emptyset, \{a\}, \{b\}, \{a, b\}\}$.

2.2 Tuples

Tuples are a collection of ordered elements. An n -tuple is written as $(a_1, a_2, a_3, \dots, a_n)$. Two n -tuples are equal if and only if all of their corresponding elements are equal.

$$(a, b) = (c, d) \equiv (a = c) \wedge (b = d)$$

A subset R of $A \times B$ is called a relation from the set A to the set B .

2.2.1 Cartesian Product

The cartesian product $A \times B$ is the set of ordered pairs (a, b) where $a \in A$ and $b \in B$.

$$A \times B = \{(a, b) | a \in A \wedge b \in B\}$$

The cartesian product of n sets is the set of tuples containing all possible combinations of elements as tuples.

$$A_1 \times A_2 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) | a_i \in A_i \text{ for } i = 1, 2, \dots, n\}$$

The cardinality of the cartesian product is equal to the product of the cardinalities of the individual sets.

$$|A_1 \times A_2 \times \dots \times A_n| = \prod_{i=1}^n |A_i|$$

2.3 Operations

The set operators are based on boolean algebra and are analogous to the operators in propositional calculus. There must always be a universal set U and all sets are assumed to be a subset of U .

Union

The union of A and B is the set that contains those elements that are contained in A , B or in both.

$$A \cup B = \{x | x \in A \vee x \in B\}$$



Complement

The complement of the set A contains all the elements in U that are not in A . It is equivalent to $U - A$.

$$\bar{A} = \{x \in U | x \notin A\}$$



Intersection

The intersection of A and B is the set that contains those elements that are contained in both A and B .

$$A \cap B = \{x | x \in A \wedge x \in B\}$$



Difference

The difference $A - B$ is the set that contains all the elements in A but not in B . It is the union of A with the complement of B .

$$A - B = \{x | x \in A \wedge x \notin B\} = A \cap \bar{B}$$



Symmetric Difference

The symmetric difference is equivalent to a logical exclusive or meaning that the symmetric difference between A and B contains the elements that are in A or in B , but not those that are in both A and B .

$$A \oplus B = (A - B) \cup (B - A)$$



2.4 Properties

Inclusion-Exclusion Principle The inclusion-exclusion principle helps to calculate the cardinality of unions of sets. The principle states that the cardinality of the union of A and B is equal to the cardinality of A plus the cardinality of B minus the cardinality of the intersection of A and B .

$$|A \cup B| = |A| + |B| - |A \cap B|$$

De Morgan's Laws

$$\overline{A \cup B} = \bar{A} \cap \bar{B}$$

$$\overline{A \cap B} = \bar{A} \cup \bar{B}$$

Identity Laws

$$A \cup \emptyset = A$$

$$A \cap U = A$$

Domination Laws

$$A \cup U = U$$

$$A \cap \emptyset = \emptyset$$

Idempotent Laws

$$A \cup A = A$$

$$A \cap A = A$$

Commutative Laws

$$A \cup B = B \cup A$$

$$A \cap B = B \cap A$$

Associative Laws

$$A \cup (B \cup C) = (A \cup B) \cup C$$

$$A \cap (B \cap C) = (A \cap B) \cap C$$

Distributive Laws

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

Absorption Laws

$$A \cup (A \cap B) = A$$

$$A \cap (A \cup B) = A$$

Complement Laws

$$A \cup \bar{A} = U$$

$$A \cap \bar{A} = \emptyset$$

Double Negation Law

$$\overline{(\bar{A})} = A$$

3 Relations

3.1 Definitions

3.2 Properties

3.3 Operations

3.4 Representing Relations

3.5 Closures

3.6 Equivalence Relations

3.7 Equivalence Classes and Partitions

3.8 Comparability

3.9 Functions

A special kind of relation is a function. A function is a subset of $A \times B$ with the restriction that no two elements may have the same first element.

$$\forall x [x \in A \rightarrow \exists y (y \in B \wedge (x, y) \in f)] \wedge \forall x, y_1, y_2 [(x, y_1) \in f \wedge (x, y_2) \in f] \rightarrow y_1 = y_2$$

3.9.1 Definitions


A function is a mapping from one set to another. A and B are two non-empty sets and the function f is a mapping from A to B .

$$f : A \rightarrow B$$

Each element of A is assigned to exactly one element B .

$$f(a) = b$$

In this case the set A is the domain of the function and the set B is the co-domain. The range is a subset of the co-domain and contains all the elements of B that f actually maps to.



3.9.2 Properties

Addition & Multiplication

$$(f_1 + f_2)(x) = f_1(x) + f_2(x)$$

$$(f_1 f_2)(x) = f_1(x) f_2(x)$$

Increasing & Decreasing

Increasing

$$\forall a, b \in A (a < b \rightarrow f(a) \leq f(b))$$

Strictly Increasing

$$\forall a, b \in A (a < b \rightarrow f(a) < f(b))$$

Decreasing

$$\forall a, b \in A (a < b \rightarrow f(a) \geq f(b))$$


Strictly Decreasing

$$\forall a, b \in A (a < b \rightarrow f(a) > f(b))$$

3.9.3 Types of Functions


Injection Functions that are injective have a one-to-one mapping from A to B . That means that every a is assigned to one b and every b is the image of at most one pre-image a .

$$\forall a, b (f(a) = f(b) \rightarrow a = b)$$




Surjection A surjective function maps A onto B , which means that for every element in B has a pre-image.

$$\forall b \in B [\exists a \in A (f(a) = b)]$$



Bijection A bijection is a function, which is both injective and surjective.



3.9.4 Inverse Function

3.9.5 Composition

3.10 Relational Algebra

4 Graphs

4.1 Trees