

DOMINION BUREAU OF STATISTICS

HEALTH AND WELFARE DIVISION

Summary of Cases of Notifiable Diseases in Canada, as reported by the Provincial Health Departments

for the week ended September 1, 1956, in comparison with the previous week¹, and the 5-year (1951-55) Median

No.	DISEASE	CANADA ²						NFLD.			P.E.I.			N.S.			N.B.			QUE.			ONT.			MAN.			SASK.			ALTA.			B.C.			No.
		Report Week	Previous Week	Median 1951-55	Cumulative Total ³			Report Week	Previous Week	Median 1951-55	Report Week	Previous Week	Median 1951-55	Report Week	Previous Week	Median 1951-55	Report Week	Previous Week	Median 1951-55	Report Week	Previous Week	Median 1951-55	Report Week	Previous Week	Median 1951-55	Report Week	Previous Week	Median 1951-55	Report Week	Previous Week	Median 1951-55	Report Week	Previous Week	Median 1951-55				
					1956	1955	Median 1951-55																															
1	Chickenpox	98	235	86	26,680	24,412	32,611	-	3	-	2	-	-	13	5	1	..	..	..	7	111	21	53	96	30	13	5	8	..	..	12	..	..	2	10	15	17	1
2	Diphtheria	2	-	1	62	68	91	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-	2	
3	Dysentery: ⁴	18	11	11	435	432	432	-	-	-	-	-	-	-	-	-	2	2	..	..	3	1	2	-	2	3	-	-	2	-	-	3	-	-	6	6	6	3
4	(a) Amoebic	-	-	-	-	2	7	-	-	-	-	-	-	-	-	-	..	..	..	..	..	..	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	
5	(b) Bacillary	18	11	11	435	430	430	-	-	-	-	-	-	-	-	-	2	2	..	..	3	1	2	-	2	3	-	-	-	-	3	-	-	6	6	6	5	
6	Encephalitis, infectious	2	2	2	17	18	23	-	-	-	..	..	..	-	-	-	-	-	-	2	-	..	..	-	-	-	-	-	-	2	-	1	-	-	-	-	6	
7	Influenza, epidemic	126	55	59	8,805	36,652	9,049	..	..	..	39	9	..	86	43	28	-	-	-	-	-	..	..	..	..	1	2	1	-	-	..	..	..	..	1	-	7	
8	Measles	152	246	121	41,208	43,678	47,327	2	-	1	1	-	1	17	13	6	..	..	1	52	40	60	156	28	5	9	11	..	..	7	10	7	16	30	9	16	8	
9	Meningitis, meningococcal	31	2	3	229	209	189	1	2	-	..	..	..	-	-	-	-	-	-	-	-	3	-	-	-	-	-	-	-	-	..	..	..	-	-	1	9	
10	Mumps	96	127	98	21,647	19,928	28,321	-	-	-	-	-	6	12	1	..	..	..	3	14	20	67	60	62	5	9	11	..	..	6	..	..	2	15	32	15	10	
11	Poliomyelitis, epidemic: ⁴	31	26	203	245	599	1,261	-	-	1	-	-	3	-	-	8	1	3	4	7	7	19	7	10	61	-	2	4	1	-	18	9	2	12	6	2	11	
12	(a) With paralysis	15	17	83	134	318	513	-	-	1	-	-	1	-	-	5	-	1	-	7	5	7	5	6	23	-	2	3	(6)	-	4	3	1	1	-	2	6	12
13	(b) Without paralysis	15	7	101	91	227	695	-	-	-	-	-	-	-	-	1	2	4	-	-	4	2	4	38	-	-	1	(6)	-	4	6	1	3	6	-	5	13	
14	Rubella	103	73	37	49,134	17,409	9,035	1	1	-	..	..	..	3	2	1	..	..	..	1	25	5	74	25	12	4	-	-	..	1	1	8	6	7	12	13	9	14
15	Scarlet fever, etc. ⁵	83	77	60	8,877	6,422	9,419	1	-	-	31	3	2	32	35	8	1	1	-	6	7	20	8	19	11	2	-	5	-	3	6	-	6	8	2	3	6	15
16	Smallpox	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	16	
17	Tuberculosis: ⁴	184	213	175	6,218	6,543	7,110	11	6	1	-	1	1	8	-	1	9	7	10	87	58	68	17	56	11	6	40	12	13	11	6	11	18	17	22	16	21	17
18	(a) Pulmonary	154	142	156	5,039	5,356	5,717	11	6	1	-	1	-	7	-	1	9	5	9	86	56	66	(6)	(6)	(6)	5	38	12	8	8	6	11	13	14	17	15	20	18
19	(b) Non-pulmonary	13	15	10	390	419	365	-	-	-	-	-	-	1	-	-	-	2	-	1	2	2	(6)	(6)	(6)	1	2	-	5	3	-	..	..	5	1	3	19	
20	Typhoid and paratyphoid	7	14	10	287	232	314	-	-	-	-	-	-	-	-	-	-	-	5	10	5	2	4	1	-	-	-	-	-	-	-	-	-	-	-	-	20	
21	Undulant fever	-	4	2	103	90	90	-	-	-	-	-	-	-	-	-	..	..	..	..	2	1	-	1	1	-	1	-	-	-	..	..	..	-	-	-	21	
22	Venereal diseases:	365	378	333	10,929	11,395	11,963	10	4	9	3	-	-	6	8	11	3	5	3	64	78	88	49	47	59	28	25	29	27	25	23	79	73	55	96	113	51	22
23	(a) Syphilis, primary	-	2	4	66	75	145	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	1	-	1	-	-	-	-	-	1	-	1	-	-	23	
24	(b) " secondary	1	1	3	30	32	78	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-	1	-	1	-	-	-	24	
25	(c) " other	37	28	36	1,231	1,613	1,778	-	1	2	-	-	-	3	-	2	-	-	1	19	14	9	3	8	11	2	1	1	1	-	1	2	-	1	7	4	4	25
26	(d) Gonorrhoea	327	347	289	9,599	9,660	9,753	10	3	6	3	-	-	3	8	8	3	5	2	45	63	68	46	39	45	26	23	27	26	25	21	76	73	52	89	108	42	26
27	(e) Other venereal diseases ⁷	-	-	-	3	15	15	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	27	
28	Whooping cough	137	130	211	5,636	9,088	5,572	-	-	-	9	1	-	3	-	2	-	-	-	34	61	69	25	14	70	8	6	9	13	5	13	6	3	17	39	40	11	28

(1) Figures adjusted for delayed reports. (2) Exclusive of Yukon and Northwest Territories. (3) Total accumulated to the end of the current week from the first week ending in January of the year stated. (4) Includes cases where type was not specified. (5) Scarlet fever includes cases of streptococcal sore throat; streptococcal sore throat not reportable in New Brunswick and Quebec. (6) Type not specified. (7) Includes chancroid, granuloma inguinale and lymphogranuloma venereum. .. Not reportable.

THE SITUATION IN THE UNITED STATES (Source: "MORBIDITY AND MORTALITY-WEEKLY REPORT", U.S. National Office of Vital Statistics): Number of reported cases for the current week (previous week in parentheses):

Diphtheria 12 (18); Measles 1,015 (1,244); Meningitis, other 33 (31); Poliomyelitis, with paralysis 288 (375); Smallpox - (-); Undulant fever 20 (10); Encephalitis infectious 81 (35); Meningococcal infections 24 (28); Poliomyelitis, acute 866 (943); Poliomyelitis, without paralysis 411 (377); Typhoid fever 48 (48);

The number of cases of poliomyelitis reported for the current week is 866 as compared with 943 for the previous week. This number is less than the figure for the same week in all prior years since 1947. This also is the first week in the current polio season that the number of reported cases is less than the number reported for the previous week. The peak week last year was the week ended August 27 and the 5-year median peak week, the second week in September. The number reported in Chicago for the current week is 62 as compared with 100 for the previous week. This is the third successive week in which reported cases have been fewer in number. Forty of the 62 cases were paralytic. Incidence in the state of Illinois exclusive of Chicago dropped slightly this week from 67 cases to 58 cases.

To date typhoid fever cases have been reported as follows: Missouri, 1 fatal case confirmed by culture (phage type C-2), 2 other clinically confirmed cases, and 4 suspect cases; Oklahoma, 1 serologically confirmed case, and 4 possible cases; Kansas, 1 suspect clinical case; California, 2 cases confirmed by culture (phage type C-2); Louisiana, 2 cases confirmed by culture (1 phage type C-2), 3 suspect cases, and 1 death -- a young man with chronic infection not confirmed as a case of typhoid fever; Oregon, 1 possible case.