

Discrete Math for Computer Science

Peter Schaefer

Freshman Fall

Contents

1	Logic	5
1.1	Propositions and Logical Operations	5
1.2	Evaluating Compound Propositions	5
1.3	Conditional Statements	5
1.4	Logical Equivalence	5
1.5	Laws of Propositional Logic	6
1.6	Predicates and Quantifiers	6
1.7	Quantified Statements	6
1.8	DeMorgan's law for Quantified Statements	7
1.9	Nested Quantifiers	7
1.10	More Nested Quantifiers	7
1.10.1	Expressing Uniqueness in Quantified Statements	7
1.10.2	Moving Quantifiers in Logical Statements	8
1.11	Logical Reasoning	8
1.11.1	The Form of an Argument	8
1.12	Rules of Inference with Propositions	9
1.13	Rules of Inference with Quantifiers	9
1.13.1	Example of using the Laws of Inference for Quantified Statements	10
1.13.2	Showing an Argument with Quantified Statements is Invalid	10
2	Proofs	11
2.1	Mathematical Definitions	11
2.2	Introduction to Proofs	11
2.3	Writing Proofs: Best Practices	11
2.4	Writing Direct Proofs	11
2.5	Proof by Contrapositive	11
2.6	Proof by Contradiction	11
2.7	Proof by Cases	11
3	Sets	12
3.1	Sets and Subsets	12
3.2	Sets of sets	13
3.3	Union and Intersection	13
3.4	More set operations	13
3.5	Set identities	13
3.6	Cartesian products	13
3.7	Partitions	13

4	Functions	14
4.1	Definition of functions	14
4.2	Floor and Ceiling functions	14
4.3	Properties of functions	14
4.4	The inverse of a function	14
4.5	Composition of functions	14
4.6	Logarithms and exponents	14
5	Boolean Algebra	15
5.1	An introduction to Boolean Algebra	15
5.2	Boolean functions	15
5.3	Disjunctive and conjunctive normal form	15
5.4	Functional completeness	15
5.5	Boolean satisfiability	15
5.6	Gates and circuits	15
6	Relation and Digraphs	16
6.1	Introduction to binary relations	16
6.2	Properties of binary relations	16
6.3	Directed graphs, paths, and cycles	16
6.4	Composition of relations	16
6.5	Graph powers and the transitive closure	16
6.6	Matrix multiplication and graph powers	16
6.7	Partial orders	16
6.8	Strict orders and directed acyclic graphs	16
6.9	Equivalence relations	16
6.10	N-ary relations and relational databases	16
7	Computation	17
7.1	An introduction to algorithms	17
7.2	Asymptotic growth of functions	17
7.3	Analysis of algorithms	17
7.4	Finite state machines	17
7.5	Turing machines	17
7.6	Decision problems and languages	17
8	Induction and Recursion	18
8.1	Sequences	18
8.2	Recurrence relations	18
8.3	Summations	18
8.4	Mathematical induction	18
8.5	More inductive proofs	18
8.6	Strong induction and well-ordering	18
8.7	Loop invariants	18
8.8	Recursive definitions	18
8.9	Structural induction	18
8.10	Recursive algorithms	18
8.11	Induction and recursive algorithms	18
8.12	Analyzing the time complexity of recursive algorithms	18
8.13	Divide-and-conquer algorithms: Introduction and mergesort	18
8.14	Divide-and-conquer algorithms: Binary Search	18
8.15	Solving linear homogeneous recurrence relations	18
8.16	Solving linear non-homogeneous recurrence relations	18
8.17	Divide-and-conquer recurrence relations	18

9 Integer Properties	19
9.1 The Division Algorithm	19
9.2 Modular arithmetic	19
9.3 Prime factorizations	19
9.4 Factoring and primality testing	19
9.5 Greatest common factor divisor and Euclid's algorithm	19
9.6 Number representation	19
9.7 Fast exponentiation	19
9.8 Introduction to cryptography	19
9.9 The RSA cryptosystem	19
10 Introduction to Counting	20
10.1 Sum and Product Rules	20
10.2 The Bijection Rules	20
10.3 The generalized product rule	20
10.4 Counting permutations	20
10.5 Counting subsets	20
10.6 Subset and permutation examples	20
10.7 Counting by complement	20
10.8 Permutations with repetitions	20
10.9 Counting multisets	20
10.10 Assignment problems: Balls in bins	20
10.11 Inclusion-exclusion principle	20
11 Advanced Counting	21
11.1 Generating permutations	21
11.2 Binomial coefficients and combinatorial identities	21
11.3 The pigeonhole principle	21
11.4 Generating functions	21
12 Discrete Probability	22
12.1 Probability of an event	22
12.2 Unions and complements of events	22
12.3 Conditional probability and independence	22
12.4 Bayes' Theorem	22
12.5 Random variables	22
12.6 Expectation of random variables	22
12.7 Linearity of expectations	22
12.8 Bernoulli trials and the binomial distribution	22
13 Graphs	23
13.1 Introduction to Graphs	23
13.2 Graph representations	23
13.3 Graph isomorphism	23
13.4 Walks, trails, circuits, paths, and cycles	23
13.5 Graph connectivity	23
13.6 Euler circuits and trails	23
13.7 Hamiltonian cycles and paths	23
13.8 Planar coloring	23
13.9 Graph coloring	23

14 Trees	24
14.1 Introduction to trees	24
14.2 Tree application examples	24
14.3 Properties of trees	24
14.4 Tree traversals	24
14.5 Spanning trees and graph traversals	24
14.6 Minimum spanning trees	24

1 Logic

1.1 Propositions and Logical Operations

Proposition: a statement that is either true or false.

Some examples include "It is raining today" and " $3 \cdot 8 = 20$ ".

However, not all statements are propositions, such as "open the door"

Name	Symbol	alternate name	p	q	$\neg p$	$p \wedge q$	$p \vee q$	$p \oplus q$
NOT	\neg	negation	T	T	F	T	T	F
AND	\wedge	conjunction	T	F	F	F	T	T
OR	\vee	disjunction	F	T	T	F	T	T
XOR	\oplus	exclusive or	F	F	T	F	F	F

XOR is very useful for encryption and binary arithmetic.

1.2 Evaluating Compound Propositions

p : The weather is bad.

$p \wedge q$: The weather is bad *and* the trip is cancelled

q : The trip is cancelled.

$p \vee q$: The weather is bad *or* the trip is cancelled

r : The trip is delayed.

$p \wedge (q \oplus r)$: The weather is bad *and* either the trip is cancelled *or* delayed

Order of Evaluation \neg , then \wedge , then \vee , but parenthesis always help for clarity.

Example Truth Table:

p	q	$p \wedge q$	$\neg q$	$(p \wedge q) \oplus \neg q$
T	T	T	F	T
T	F	F	T	T
F	T	F	F	F
F	F	F	T	T

1.3 Conditional Statements

$p \rightarrow q$ where p is the hypothesis and q is the conclusion

Format	Terminology	
$p \rightarrow q$	given	given
$\neg q \rightarrow \neg p$	contrapositive	$p \rightarrow q \equiv \neg q \rightarrow \neg p$ contrapositive
$q \rightarrow p$	converse	inverse
$\neg p \rightarrow \neg q$	inverse	$\neg p \rightarrow \neg q \equiv q \rightarrow p$ converse

p	q	$p \rightarrow q$		Phrase	Logic
T	T	T	p is a <u>sufficient</u> condition for q	q if p	$p \rightarrow q$
T	F	F	q is a <u>necessary</u> condition for p	q only if p	$q \rightarrow p$
F	T	T		q if and only if p	$p \leftrightarrow q$
F	F	T			

Order of Operations: $p \wedge q \rightarrow r \equiv (p \wedge q) \rightarrow r$

1.4 Logical Equivalence

Tautology: a proposition that is always true

Contradiction: a proposition that is always false

Logically equivalent: same truth value regardless of the truth values of their individual propositions

DeMorgan's Laws:

$$\neg(p \vee q) \equiv \neg p \wedge \neg q$$

$$\neg(p \wedge q) \equiv \neg p \vee \neg q$$

Verbally,

It is not true that the patient has migraines *or* high blood pressure \equiv
 \equiv The patient does not have migraines *and* does not have high blood pressure

It is not true that the patient has migraines *and* high blood pressure \equiv
 \equiv The patient does not have migraines *or* does not have high blood pressure

1.5 Laws of Propositional Logic

You can use **substitution** on logically equivalent propositions.

Law Name	\vee or	\wedge and
Idempotent	$p \vee p \equiv p$	$p \wedge p \equiv p$
Associative	$(p \vee q) \vee r \equiv p \vee (q \vee r)$	$(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$
Commutative	$p \vee q \equiv q \vee p$	$p \wedge q \equiv q \wedge p$
Distributive	$p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$	$p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$
Identity	$p \vee F \equiv p$	$p \wedge T \equiv p$
Domination	$p \vee T \equiv T$	$p \wedge F \equiv F$
Double Negation	$\neg \neg p \equiv p$	
Complement	$p \vee \neg p \equiv T$	$p \wedge \neg p \equiv F$
DeMorgan	$\neg(p \vee q) \equiv \neg p \wedge \neg q$	$\neg(p \wedge q) \equiv \neg p \vee \neg q$
Absorption	$p \vee (p \wedge q) \equiv p$	$p \wedge (p \vee q) \equiv p$
Conditional	$p \rightarrow q \equiv \neg p \vee q$	$p \leftrightarrow q \equiv (p \rightarrow q) \wedge (q \rightarrow p)$

1.6 Predicates and Quantifiers

Predicate: a logical statement where truth value is a function of a variable.

$P(x)$: x is an even number. $P(5)$: false $P(2)$: true

Domain: the set of all possible values for a variable in a predicate.

Ex. \mathbb{Z}^+ is the set of all positive integers.

*If domain is not clear from context, it should be given as part of the definition of the predicate.

Quantifier	Symbol	Meaning
Universal	\forall	"for all"
Existential	\exists	"there exists"

Quantifier: converts a predicate to a proposition.

$\exists x(x + 1 < x)$ is false.

Counter Example: universally quantified statement where an element in the domain for which the predicate is false. Useful to prove a \forall statement false.

1.7 Quantified Statements

Consider the two following two predicates:

$P(x)$: x is prime, $x \in \mathbb{Z}^+$

$O(x)$: x is odd

Proposition made of predicates: $\exists x(P(x) \wedge \neg O(x))$

Verbally: there exists a positive integer that is prime but is not odd.

Free Variable: a variable that is free to be any value in the domain.

Bound Variable: a variable that is bound to a quantifier.

			$P(x)$	$S(x)$	$\neg S(x)$
$P(x)$:	x came to the party	$P(x) \stackrel{?}{\equiv} \neg S(x)$	Joe	T	F
$S(x)$:	x was sick	$P(x) \not\equiv \neg S(x)$	Theo	F	T
			Gert	T	F
			Sam	F	T

1.8 DeMorgan's law for Quantified Statements

Consider the predicate: $F(x) : "x \text{ can fly}"$, where x is a bird. According to the DeMorgan Identity for Quantified Statements,

$$\neg \forall x F(x) \equiv \exists x \neg F(x)$$

"not every bird can fly \equiv "there exists a bird that cannot fly"

Example using DeMorgan Identities:

$$\begin{aligned} \neg \exists x (P(x) \rightarrow \neg Q(x)) &\equiv \forall x \neg (P(x) \rightarrow \neg Q(x)) \\ &\equiv \forall x (\neg \neg P(x) \wedge \neg \neg Q(x)) \\ &\equiv \forall x (P(x) \wedge Q(x)) \end{aligned}$$

1.9 Nested Quantifiers

A logical expression with more than one quantifier that binds different variables in the same predicate is said to have **Nested Quantifiers**.

Logic	Variable Boundedness	Logic	Meaning
$\forall x \exists y P(x, y)$	x, y bound	$\forall x \forall y M(x, y)$	"everyone sent an email to everyone"
$\forall x P(x, y)$	x bound, y free	$\forall x \exists y M(x, y)$	"everyone sent an email to someone"
$\exists x \exists y T(x, y, z)$	x, y bound, z free	$\exists x \forall y M(x, y)$	"someone sent an email to everyone"
		$\exists x \exists y M(x, y)$	"someone sent an email to someone"

There is a two-player game analogy for how quantifiers work:

Player	Action	Goal
Existential Player \exists	selects value for existentially-bound variables	tries to make expression <u>true</u>
Universal Player \forall	selects value for universally-bound variables	tries to make expression <u>false</u>

Consider the predicate $L(x, y) : "x \text{ likes } y"$.

$\exists x \forall y L(x, y)$ means "there is a student who likes everyone in the school".

$\neg \exists x \forall y L(x, y)$ means "there is no student who likes everyone in the school".

After applying DeMorgan's Laws,

$\forall x \exists y \neg L(x, y)$ means "there is no student who likes everyone in the school".

1.10 More Nested Quantifiers

$M(x, y) : "x \text{ sent an email to } y"$. Consider $\forall x \forall y M(x, y)$. It means that "email sent an email to everyone including themselves". Using $(x \neq y \rightarrow M(x, y))$ can fix this quirk.

$\forall x \forall y (x \neq y \rightarrow M(x, y))$ means "everyone sent an email to everyone else"

1.10.1 Expressing Uniqueness in Quantified Statements

Consider $L(x)$: x was late to the meeting. If someone was late to the meeting, how could you express that that someone was the only person late to the meeting? You want to express that there is someone where everyone else was not late, which can be done with

$$\exists x (L(x) \wedge \forall y (x \neq y \rightarrow \neg L(y)))$$

1.10.2 Moving Quantifiers in Logical Statements

Consider $M(x, y)$: " x is married to y " and $A(x)$: " x is an adult". One way of expressing "For every person x , if x is an adult, then there is a person y to whom x is married to" is by this statement:

$$\forall x(A(x) \rightarrow \exists y M(x, y))$$

Since y does not appear in $A(x)$, " $\exists y$ " can be moved so that it appears just after the " \forall ", resulting with

$$\forall x \exists y(A(x) \rightarrow M(x, y))$$

When doing this, keep in mind that $\forall x \exists y \neq \exists y \forall x$:

$\forall x \exists y(A(x) \rightarrow M(x, y))$ means

for every x , if x is an adult, there exists y who is married to x .

$\exists y \forall x(A(x) \rightarrow M(x, y))$ means

There exists a y , such that every x who is an adult is also married to y

1.11 Logical Reasoning

Argument: a sequence of propositions, called hypothesis, followed by a final proposition, called the conclusion.

An argument is **valid** if the conclusion is true whenever the hypothesis are all true, otherwise the argument is **invalid**.

$$\begin{array}{c} p_1 \\ p_2 \\ \vdots \\ p_n \\ \hline \therefore c \end{array} \quad \text{where } \begin{array}{l} p_1, p_2, \dots, p_n \text{ are hypothesis} \\ c \text{ is the conclusion} \end{array}$$

The argument is valid whenever the proposition $(p_1 \wedge p_2 \wedge \dots \wedge p_n) \rightarrow c$ is a tautology. Additionally, because of the commutative law, hypothesis can be reordered without changing the argument.

$$\frac{p}{p \rightarrow q} \quad \equiv \quad \frac{p \rightarrow q}{p} \quad \frac{p}{\therefore q}$$

1.11.1 The Form of an Argument

$$\begin{array}{l} \text{It is raining today.} \\ \text{If it is raining today, then I will not ride my bike to school.} \\ \hline \therefore \text{I will not ride my bike to school.} \end{array} \quad \begin{array}{l} p \\ p \rightarrow q \\ \hline \therefore q \end{array}$$

The argument is valid because its form, $\frac{p}{p \rightarrow q} \therefore q$ is an valid argument.

$$\begin{array}{l} 5 \text{ is not an even number.} \\ \text{If 5 is an even number, then 7 is an even number.} \\ \hline \therefore 7 \text{ is not an even number.} \end{array} \quad \begin{array}{l} p \\ p \rightarrow q \\ \hline \therefore q \end{array}$$

The argument is invalid because its form, $\frac{\neg p}{p \rightarrow q} \therefore \neg q$ is an invalid argument.

1.12 Rules of Inference with Propositions

Using truth tables to establish the validity of an argument can become tedious, especially if an argument uses a large number of variables.

$\frac{p}{p \rightarrow q} \quad \text{Modus Ponens}$	$\frac{p}{q} \quad \text{Conjunction}$
$\frac{\neg q}{p \rightarrow q} \quad \text{Modus Tollens}$	$\frac{p \rightarrow q}{q \rightarrow r} \quad \text{Hypothetical Syllogism}$
$\frac{p}{\therefore p \vee q} \quad \text{Addition}$	$\frac{p \vee q}{\neg p} \quad \text{Disjunctive Syllogism}$
$\frac{p \wedge q}{\therefore p} \quad \text{Simplification}$	$\frac{p \rightarrow q}{q \rightarrow r} \quad \text{Resolution}$
	$\frac{q \rightarrow r}{\therefore q \vee r}$

Example expressed in English:

If it is raining or windy or both, the game will be cancelled.	$(r \vee w) \rightarrow c$
The game will not be cancelled	$\neg c$
\therefore It is not windy.	$\therefore \neg w$

Steps to Solve:

$(r \vee w) \rightarrow c$	Hypothesis	(1)
$\neg c$	Hypothesis	(2)
$\neg(r \vee w)$	Modus Tollens: 1, 2	(3)
$\neg r \wedge \neg w$	DeMorgan's Law: 3	(4)
$\neg w \wedge \neg r$	Commutative Law: 4	(5)
$\neg w$	Simplification: 5	(6)

1.13 Rules of Inference with Quantifiers

In order to apply the rules of quantified expressions, such as $\forall x \neg (P(x) \wedge Q(x))$, we need to remove the quantifier by plugging in a value from the domain to replace the variable x .

For example:

Every employee who received a large bonus works hard.	$\forall x (B(x) \rightarrow H(x))$
Linda is an employee at the company.	$Linda \in x$
Linda received a large bonus.	$B(Linda)$
\therefore Some employee works hard.	$\therefore \exists x H(x)$

Arbitrary Element: has no special properties other than those shared by all elements of the domain.

Particular Element: may have special properties that are not shared by all the elements of the domain. For example, if the domain is the set of all integers, \mathbb{Z} , a particular element is 3, because it is odd, which is not true for all integers.

Rules of Inference for Quantified Statements

c is an element $\frac{\forall x P(x)}{\therefore P(c)}$	Universal Instantiation	$\frac{\exists x P(x)}{\therefore c \text{ is particular} \wedge P(c)}$	Existential Instantiation*
c is arbitrary $\frac{P(c)}{\therefore \forall P(x)}$	Universal Generalization	$\frac{c \text{ is an element} \quad P(c)}{\therefore \exists x P(x)}$	Existential Generalization

*Each use of Existential Instantiation must define a new element with its own symbol or name.

1.13.1 Example of using the Laws of Inference for Quantified Statements

Consider the following argument:

$$\frac{\begin{array}{l} \forall x(P(x) \vee Q(x)) \\ 3 \text{ is an integer} \\ \neg P(3) \end{array}}{\therefore Q(3)}$$

Steps to Solve:

$\forall x(P(x) \vee Q(x))$	Hypothesis	(1)
3 is an integer	Hypothesis	(2)
$(P(3) \vee Q(3))$	Universal Instantiation: 1, 2	(3)
$\neg P(3)$	Hypothesis	(4)
$Q(3)$	Disjunctive Syllogism: 3, 4	(5)

1.13.2 Showing an Argument with Quantified Statements is Invalid

Consider the following argument:

$$\frac{\begin{array}{l} \exists x P(x) \\ \exists x Q(x) \end{array}}{\therefore \exists x(P(x) \wedge Q(x))}$$

Using a supposed domain $\{c, d\}$, with truth values of

	P	Q
c	T	F
d	F	T

, the argument is invalid.

2 Proofs

2.1 Mathematical Definitions

2.2 Introduction to Proofs

2.3 Writing Proofs: Best Practices

2.4 Writing Direct Proofs

2.5 Proof by Contrapositive

2.6 Proof by Contradiction

2.7 Proof by Cases

3 Sets

3.1 Sets and Subsets

A **set** is a collection of objects. Objects in a set are called **elements**. Order does not matter, and there are no duplicates.

Roster notation:

$$\begin{aligned} A &= \{2, 4, 6, 10\} \\ B &= \{4, 6, 10, 2\} \\ A &= B \end{aligned}$$

To show membership, use the \in symbol. For example, $2 \in A$, while $7 \notin A$. The empty set, which contains nothing, typically uses the \emptyset symbol, or $\{\}$. Sets can be finite, or infinite. **Cardinality** of a set is the number of elements in a set. For example, the cardinality of A is 4.

$$|A| = 4$$

Cardinality can be infinite. Consider the set of all the integers, \mathbb{Z} . $|\mathbb{Z}| = \infty$

$$\begin{aligned} \mathbb{N} &: \text{set of natural numbers} \\ &= \{0, 1, 2, 3, \dots\} \end{aligned}$$

$$\begin{aligned} \mathbb{Z} &: \text{set of integers} \\ &= \{\dots, -2, -1, 0, 1, 2, \dots\} \end{aligned}$$

$$\begin{aligned} \mathbb{Q} &: \text{set of rational numbers} \\ &= \{x | x = \frac{a}{b} \text{ where } a, b \in \mathbb{Z}, b \neq 0\} \end{aligned}$$

$$\begin{aligned} \mathbb{R} &: \text{set of real numbers} \\ &= \{x | x \text{ has a decimal representation}\} \end{aligned}$$


The subset operator is \subseteq

$$\begin{aligned} A &\subseteq B \text{ if } \forall x(x \in A \rightarrow x \in B) \\ A &\subseteq A \text{ is true for any set} \\ \emptyset &\subseteq A \text{ is true for any set} \end{aligned}$$

Sometimes it is easier to define a set by defining properties that all the elements have. That is easy to do in **set builder notation**.

$$\begin{aligned} A &= \{x \in S : P(x)\}, \text{ where } S \text{ is another set} \\ C &= \{x \in \mathbb{Z} : 0 < x < 100 \text{ and } x \text{ is prime}\}. \\ D &= \{x \in \mathbb{R} : |x| < 1\} \end{aligned}$$

The **Universal Set**, usually called 'U', is a set that contains all elements mentioned in a particular context. For example, a discussion about certain types of real numbers, it would be understood that any element in the discussion is a real number. Sets are often represented pictorially with **Venn Diagrams**.



If $A \subseteq B$ and there is an element of B that is not an element of A , meaning $A \neq B$, then A is a **proper subset** of B , denoted as $A \subset B$. An important fact is that

$$\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R}$$

3.2 Sets of sets

3.3 Union and Intersection

3.4 More set operations

3.5 Set identities

3.6 Cartesian products

3.7 Partitions

4 Functions

- 4.1 Definition of functions
- 4.2 Floor and Ceiling functions
- 4.3 Properties of functions
- 4.4 The inverse of a function
- 4.5 Composition of functions
- 4.6 Logarithms and exponents

5 Boolean Algebra

5.1 An introduction to Boolean Algebra

5.2 Boolean functions

5.3 Disjunctive and conjunctive normal form

5.4 Functional completeness

5.5 Boolean satisfiability

5.6 Gates and circuits

6 Relation and Digraphs

- 6.1 Introduction to binary relations
- 6.2 Properties of binary relations
- 6.3 Directed graphs, paths, and cycles
- 6.4 Composition of relations
- 6.5 Graph powers and the transitive closure
- 6.6 Matrix multiplication and graph powers
- 6.7 Partial orders
- 6.8 Strict orders and directed acyclic graphs
- 6.9 Equivalence relations
- 6.10 N-ary relations and relational databases

7 Computation

- 7.1 An introduction to algorithms
- 7.2 Asymptotic growth of functions
- 7.3 Analysis of algorithms
- 7.4 Finite state machines
- 7.5 Turing machines
- 7.6 Decision problems and languages

8 Induction and Recursion

8.1 Sequences

8.2 Recurrence relations

8.3 Summations

8.4 Mathematical induction

8.5 More inductive proofs

8.6 Strong induction and well-ordering

8.7 Loop invariants

8.8 Recursive definitions

8.9 Structural induction

8.10 Recursive algorithms

8.11 Induction and recursive algorithms

8.12 Analyzing the time complexity of recursive algorithms

8.13 Divide-and-conquer algorithms: Introduction and mergesort

8.14 Divide-and-conquer algorithms: Binary Search

8.15 Solving linear homogeneous recurrence relations

8.16 Solving linear non-homogeneous recurrence relations

8.17 Divide-and-conquer recurrence relations

9 Integer Properties

9.1 The Division Algorithm

9.2 Modular arithmetic

9.3 Prime factorizations

9.4 Factoring and primality testing

9.5 Greatest common factor divisor and Euclid's algorithm

9.6 Number representation

9.7 Fast exponentiation

9.8 Introduction to cryptography

9.9 The RSA cryptosystem

10 Introduction to Counting

- 10.1 Sum and Product Rules
- 10.2 The Bijection Rules
- 10.3 The generalized product rule
- 10.4 Counting permutations
- 10.5 Counting subsets
- 10.6 Subset and permutation examples
- 10.7 Counting by complement
- 10.8 Permutations with repetitions
- 10.9 Counting multisets
- 10.10 Assignment problems: Balls in bins
- 10.11 Inclusion-exclusion principle

11 Advanced Counting

11.1 Generating permutations

11.2 Binomial coefficients and combinatorial identities

11.3 The pigeonhole principle

11.4 Generating functions

12 Discrete Probability

- 12.1 Probability of an event
- 12.2 Unions and complements of events
- 12.3 Conditional probability and independence
- 12.4 Bayes' Theorem
- 12.5 Random variables
- 12.6 Expectation of random variables
- 12.7 Linearity of expectations
- 12.8 Bernoulli trials and the binomial distribution

13 Graphs

13.1 Introduction to Graphs

13.2 Graph representations

13.3 Graph isomorphism

13.4 Walks, trails, circuits, paths, and cycles

13.5 Graph connectivity

13.6 Euler circuits and trails

13.7 Hamiltonian cycles and paths

13.8 Planar coloring

13.9 Graph coloring

14 Trees

14.1 Introduction to trees

14.2 Tree application examples

14.3 Properties of trees

14.4 Tree traversals

14.5 Spanning trees and graph traversals

14.6 Minimum spanning trees