

1 Homework 5

Section 4.2

7

Which of the following are linear combinations of $\vec{u} = (0, -2, 2)$ and $\vec{v} = (1, 3, -1)$?

a. $(2, 2, 2)$

Proof. Let $k_1, k_2 \in \mathbb{R}$ such that $k_1\vec{u} + k_2\vec{v} = (2, 2, 2)$. That is, $k_1(0, -2, 2) + k_2(1, 3, -1) = (2, 2, 2)$. From this equation, we get a linear system of equations.

$$\begin{aligned} 0k_1 + 1k_2 &= 2 \\ -2k_1 + 3k_2 &= 2 \\ 2k_1 - 1k_2 &= 2 \end{aligned}$$

$$\begin{aligned} \left(\begin{array}{cc|c} 0 & 1 & 2 \\ -2 & 3 & 2 \\ 2 & -1 & 2 \end{array} \right) &\xrightarrow{-\frac{1}{2}R_2} \left(\begin{array}{ccc} 0 & 1 & 2 \\ 1 & -\frac{3}{2} & -1 \\ 2 & -1 & 2 \end{array} \right) &\xrightarrow{R_1 \leftrightarrow R_2} \left(\begin{array}{ccc} 1 & -\frac{3}{2} & -1 \\ 0 & 1 & 2 \\ 2 & -1 & 2 \end{array} \right) &\xrightarrow[\begin{smallmatrix} R_3-2R_1 \\ (-2, 3, 2) \end{smallmatrix}]{R_3-2R_1} \left(\begin{array}{ccc} 1 & -\frac{3}{2} & -1 \\ 0 & 1 & 2 \\ 0 & 2 & 4 \end{array} \right) &\xrightarrow[\begin{smallmatrix} R_3-2R_2 \\ (0, -2, -4) \end{smallmatrix}]{R_3-2R_2} \\ &\left(\begin{array}{ccc} 1 & -\frac{3}{2} & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{array} \right) &\xrightarrow[\begin{smallmatrix} R_1+\frac{3}{2}R_2 \\ (0, \frac{3}{2}, 3) \end{smallmatrix}]{R_1+\frac{3}{2}R_2} \left(\begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{array} \right) \end{aligned}$$

This augmented matrix represents the following equations:

$$\begin{aligned} k_1 + 0k_2 &= 2 & k_1 &= 2 \\ 0k_1 + k_2 &= 2 & k_2 &= 2 \\ 0 + 0 &= 0 \end{aligned}$$

This means that $(2, 2, 2)$ is a linear combination of $\{\vec{u}, \vec{v}\}$, when $k_1 = 2$ and $k_2 = 2$. □

c. $(0, 4, 5)$

Proof. Let $k_1, k_2 \in \mathbb{R}$ such that $k_1\vec{u} + k_2\vec{v} = (0, 4, 5)$. That is, $k_1(0, -2, 2) + k_2(1, 3, -1) = (0, 4, 5)$. From this equation, we get a linear system of equations.

$$\begin{aligned} 0k_1 + 1k_2 &= 0 \\ -2k_1 + 3k_2 &= 4 \\ 2k_1 - 1k_2 &= 5 \end{aligned}$$

$$\left(\begin{array}{cc|c} 0 & 1 & 0 \\ -2 & 3 & 4 \\ 2 & -1 & 5 \end{array} \right) \xrightarrow{-\frac{1}{2}R_2} \left(\begin{array}{ccc} 0 & 1 & 0 \\ 1 & -\frac{3}{2} & -2 \\ 2 & -1 & 5 \end{array} \right) \xrightarrow[\begin{smallmatrix} R_3-2R_1 \\ (-2, 3, 4) \end{smallmatrix}]{R_3-2R_1} \left(\begin{array}{ccc} 0 & 1 & 0 \\ 1 & -\frac{3}{2} & -2 \\ 0 & 2 & 9 \end{array} \right) \xrightarrow[\begin{smallmatrix} R_3-2R_2 \\ (0, -2, 0) \end{smallmatrix}]{R_3-2R_2} \left(\begin{array}{cc|c} 0 & 1 & 0 \\ 1 & -\frac{3}{2} & -2 \\ 0 & 0 & 9 \end{array} \right)$$

The last row from this matrix provides the equation $0k_1 + 0k_2 = 9$, meaning $0 + 0 = 9$, which is impossible. Therefore, $(0, 4, 5)$ is not spanned by $\{\vec{u}, \vec{v}\}$. □

8

Express the following combinations of $\vec{u} = (2, 1, 4)$, $\vec{v} = (1, -1, 3)$, and $\vec{w} = (3, 2, 5)$

a. $(-9, -7, -15)$

Proof. Let $k_1, k_2, k_3 \in \mathbb{R}$ such that $k_1\vec{u} + k_2\vec{v} + k_3\vec{w} = (-9, -7, -15)$. That is $k_1(2, 1, 4) + k_2(1, -1, 3) + k_3(3, 2, 5) = (-9, -7, -15)$. From this equation, we get a linear system of equations.

$$\begin{aligned} 2k_1 + k_2 + 3k_3 &= -9 \\ 1k_1 - k_2 + 2k_3 &= -7 \\ 4k_1 + 3k_2 + 5k_3 &= -15 \end{aligned}$$

$$\begin{aligned} \left(\begin{array}{ccc|c} 2 & 1 & 3 & -9 \\ 1 & -1 & 2 & -7 \\ 4 & 3 & 5 & -15 \end{array} \right) &\xrightarrow[(-4, -2, -6, 18)]{R_3 - 2R_1} \left(\begin{array}{ccc|c} 2 & 1 & 3 & -9 \\ 1 & -1 & 2 & -7 \\ 0 & 1 & -1 & 3 \end{array} \right) \xrightarrow[(-2, 2, -4, 14)]{R_1 - 2R_2} \left(\begin{array}{ccc|c} 0 & 3 & -1 & 5 \\ 1 & -1 & 2 & -7 \\ 0 & 1 & -1 & 3 \end{array} \right) \xrightarrow{R_2 + R_1} \\ \left(\begin{array}{ccc|c} 0 & 3 & -1 & 5 \\ 1 & 0 & 1 & -4 \\ 0 & 1 & -1 & 3 \end{array} \right) &\xrightarrow[(0, -3, 3, -9)]{R_1 - 3R_3} \left(\begin{array}{ccc|c} 0 & 0 & 2 & -4 \\ 1 & 0 & 1 & -4 \\ 0 & 1 & -1 & 3 \end{array} \right) \xrightarrow{\frac{1}{2}R_1} \left(\begin{array}{ccc|c} 0 & 0 & 1 & -2 \\ 1 & 0 & 1 & -4 \\ 0 & 1 & -1 & 3 \end{array} \right) \xrightarrow[(0, 0, -1, 2)]{R_2 - R_1} \\ \left(\begin{array}{ccc|c} 0 & 0 & 1 & -2 \\ 1 & 0 & 0 & -2 \\ 0 & 1 & -1 & 3 \end{array} \right) &\xrightarrow[(0, 0, 1, -2)]{R_3 + R_1} \left(\begin{array}{ccc|c} 0 & 0 & 1 & -2 \\ 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & 1 \end{array} \right) \xrightarrow[R_2 \leftrightarrow R_3]{R_1 \leftrightarrow R_2} \left(\begin{array}{ccc|c} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & -2 \end{array} \right) \end{aligned}$$

This augmented matrix represents the following equations:

$$\begin{aligned} k_1 &= -2 \\ k_2 &= 1 \\ k_3 &= -2 \end{aligned}$$

This means that $(-9, -7, -15)$ is a linear combination of $\{\vec{u}, \vec{v}, \vec{w}\}$, when $k_1 = -2, k_2 = 1$, and $k_3 = -2$. \square

c. $(0, 0, 0)$

Proof. Let $k_1, k_2, k_3 \in \mathbb{R}$ such that $k_1\vec{u} + k_2\vec{v} + k_3\vec{w} = (0, 0, 0)$. That is $k_1(2, 1, 4) + k_2(1, -1, 3) + k_3(3, 2, 5) = (0, 0, 0)$. From this equation, we get a linear system of equations.

$$\begin{aligned} 2k_1 + k_2 + 3k_3 &= 0 \\ 1k_1 - k_2 + 2k_3 &= 0 \\ 4k_1 + 3k_2 + 5k_3 &= 0 \end{aligned}$$

$$\begin{aligned} \left(\begin{array}{ccc|c} 2 & 1 & 3 & 0 \\ 1 & -1 & 2 & 0 \\ 4 & 3 & 5 & 0 \end{array} \right) &\xrightarrow[(-4, -2, -6, 0)]{R_3 - 2R_1} \left(\begin{array}{ccc|c} 2 & 1 & 3 & 0 \\ 1 & -1 & 2 & 0 \\ 0 & 1 & -1 & 0 \end{array} \right) \xrightarrow[(-2, 2, -4, 0)]{R_1 - 2R_2} \left(\begin{array}{ccc|c} 0 & 3 & -1 & 0 \\ 1 & -1 & 2 & 0 \\ 0 & 1 & -1 & 0 \end{array} \right) \xrightarrow{R_2 + R_1} \\ \left(\begin{array}{ccc|c} 0 & 3 & -1 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 0 \end{array} \right) &\xrightarrow[(0, -3, 3, 0)]{R_1 - 3R_3} \left(\begin{array}{ccc|c} 0 & 0 & 2 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 0 \end{array} \right) \xrightarrow{\frac{1}{2}R_1} \left(\begin{array}{ccc|c} 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & -1 & 0 \end{array} \right) \xrightarrow[(0, 0, -1, 0)]{R_2 - R_1} \\ \left(\begin{array}{ccc|c} 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 \end{array} \right) &\xrightarrow[(0, 0, 1, 0)]{R_3 + R_1} \left(\begin{array}{ccc|c} 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{array} \right) \xrightarrow[R_2 \leftrightarrow R_3]{R_1 \leftrightarrow R_2} \left(\begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right) \end{aligned}$$

This augmented matrix represents the following equations:

$$\begin{aligned} k_1 &= 0 \\ k_2 &= 0 \\ k_3 &= 0 \end{aligned}$$

This means that $(0, 0, 0)$ is a linear combination of $\{\vec{u}, \vec{v}, \vec{w}\}$, when $k_1 = 0, k_2 = 0$, and $k_3 = 0$. \square

9

Which of the following are linear combinations of

$$A = \begin{bmatrix} 4 & 0 \\ -2 & -2 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix}, \quad C = \begin{bmatrix} 0 & 2 \\ 1 & 4 \end{bmatrix}?$$

a. $\begin{bmatrix} 6 & -8 \\ -1 & -8 \end{bmatrix}$

Proof. Let $k_1, k_2, k_3 \in \mathbb{R}$ such that $k_1A + k_2B + k_3C = \begin{bmatrix} 6 & -8 \\ -1 & -8 \end{bmatrix}$.

That is, $k_1 \begin{bmatrix} 4 & 0 \\ -2 & -2 \end{bmatrix} + k_2 \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} + k_3 \begin{bmatrix} 0 & 2 \\ 1 & 4 \end{bmatrix} = \begin{bmatrix} 6 & -8 \\ -1 & -8 \end{bmatrix}$. From this equation, we get a linear system of equations.

$$\begin{aligned} 4k_1 + 1k_2 + 0k_3 &= 6 \\ 0k_1 - 1k_2 + 2k_3 &= -8 \\ -2k_1 + 2k_2 + 1k_3 &= -1 \\ -2k_1 + 3k_2 + 4k_3 &= -8 \end{aligned}$$

$$\begin{aligned} &\left(\begin{array}{ccc|c} 4 & 1 & 0 & 6 \\ 0 & -1 & 2 & -8 \\ -2 & 2 & 1 & -1 \\ -2 & 3 & 4 & -8 \end{array} \right) \xrightarrow[(-4, 4, 2, -2)]{R_1 + 2R_3} \left(\begin{array}{ccc|c} 0 & 5 & 2 & 4 \\ 0 & -1 & 2 & -8 \\ -2 & 2 & 1 & -1 \\ -2 & 3 & 4 & -8 \end{array} \right) \xrightarrow[(2, -2, -1, 1)]{R_4 - R_3} \left(\begin{array}{ccc|c} 0 & 5 & 2 & 4 \\ 0 & -1 & 2 & -8 \\ -2 & 2 & 1 & -1 \\ 0 & 1 & 3 & -7 \end{array} \right) \xrightarrow{-\frac{1}{2}R_3} \\ &\left(\begin{array}{ccc|c} 0 & 5 & 2 & 4 \\ 0 & -1 & 2 & -8 \\ 1 & -1 & -\frac{1}{2} & \frac{1}{2} \\ 0 & 1 & 3 & -7 \end{array} \right) \xrightarrow[R_2 + R_4]{R_3 + R_4} \left(\begin{array}{ccc|c} 0 & 5 & 2 & 4 \\ 0 & 0 & 5 & -15 \\ 1 & 0 & 2\frac{1}{2} & -6\frac{1}{2} \\ 0 & 1 & 3 & -7 \end{array} \right) \xrightarrow[(0, -5, -15, 35)]{R_1 - 5R_4} \left(\begin{array}{ccc|c} 0 & 0 & -13 & 39 \\ 0 & 0 & 5 & -15 \\ 1 & 0 & 2\frac{1}{2} & -6\frac{1}{2} \\ 0 & 1 & 3 & -7 \end{array} \right) \xrightarrow[\frac{1}{5}R_2]{R_1 + 2R_2} \\ &\left(\begin{array}{ccc|c} 0 & 0 & -3 & 9 \\ 0 & 0 & 1 & -3 \\ 1 & 0 & 2\frac{1}{2} & -6\frac{1}{2} \\ 0 & 1 & 3 & -7 \end{array} \right) \xrightarrow[R_1 \leftrightarrow R_4]{R_1 + 3R_2} \left(\begin{array}{ccc|c} 0 & 1 & 3 & -7 \\ 0 & 0 & 1 & -3 \\ 1 & 0 & 2\frac{1}{2} & -6\frac{1}{2} \\ 0 & 0 & 0 & 0 \end{array} \right) \xrightarrow[(0, 0, -3, 9)]{R_1 - 3R_2} \left(\begin{array}{ccc|c} 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & -3 \\ 1 & 0 & 2\frac{1}{2} & -6\frac{1}{2} \\ 0 & 0 & 0 & 0 \end{array} \right) \xrightarrow[(0, 0, -2\frac{1}{2}, 7\frac{1}{2})]{R_3 - 2\frac{1}{2}R_2} \\ &\left(\begin{array}{ccc|c} 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & -3 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right) \rightarrow \left(\begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & -3 \\ 0 & 0 & 0 & 0 \end{array} \right) \end{aligned}$$

This augmented matrix represents the following equations:

$$\begin{aligned} k_1 &= 1 \\ k_2 &= 2 \\ k_3 &= -3 \end{aligned}$$

This means that $\begin{bmatrix} 6 & -8 \\ -1 & -8 \end{bmatrix}$ is a linear combination of $\{A, B, C\}$, when $k_1 = 1, k_2 = 2$, and $k_3 = -3$. □

c. $\begin{bmatrix} 6 & 0 \\ 3 & 8 \end{bmatrix}$

Proof. Let $k_1, k_2, k_3 \in \mathbb{R}$ such that $k_1A + k_2B + k_3C = \begin{bmatrix} 6 & 0 \\ 3 & 8 \end{bmatrix}$.

That is, $k_1 \begin{bmatrix} 4 & 0 \\ -2 & -2 \end{bmatrix} + k_2 \begin{bmatrix} 1 & -1 \\ 2 & 3 \end{bmatrix} + k_3 \begin{bmatrix} 0 & 2 \\ 1 & 4 \end{bmatrix} = \begin{bmatrix} 6 & 0 \\ 3 & 8 \end{bmatrix}$. From this equation, we get a linear system of equations.

$$\begin{aligned} 4k_1 + 1k_2 + 0k_3 &= 6 \\ 0k_1 - 1k_2 + 2k_3 &= 0 \\ -2k_1 + 2k_2 + 1k_3 &= 3 \\ -2k_1 + 3k_2 + 4k_3 &= 8 \end{aligned}$$

$$\begin{aligned} &\left(\begin{array}{cccc|c} 4 & 1 & 0 & 0 & 6 \\ 0 & -1 & 2 & 0 & 0 \\ -2 & 2 & 1 & 0 & 3 \\ -2 & 3 & 4 & 0 & 8 \end{array}\right) \xrightarrow[(-4, 4, 2, 6)]{R_1 + 2R_3} \left(\begin{array}{cccc|c} 0 & 5 & 2 & 12 & 6 \\ 0 & -1 & 2 & 0 & 0 \\ -2 & 2 & 1 & 0 & 3 \\ -2 & 3 & 4 & 0 & 8 \end{array}\right) \xrightarrow[(2, -2, -1, -3)]{R_4 - R_3} \left(\begin{array}{cccc|c} 0 & 5 & 2 & 12 & 6 \\ 0 & -1 & 2 & 0 & 0 \\ -2 & 2 & 1 & 0 & 3 \\ 0 & 1 & 3 & 5 & 5 \end{array}\right) \xrightarrow{-\frac{1}{2}R_3} \\ &\left(\begin{array}{cccc|c} 0 & 5 & 2 & 12 & 6 \\ 0 & -1 & 2 & 0 & 0 \\ 1 & -1 & -\frac{1}{2} & -\frac{1}{2} & \frac{3}{2} \\ 0 & 1 & 3 & 5 & 5 \end{array}\right) \xrightarrow[R_2 + R_4]{R_3 + R_4} \left(\begin{array}{cccc|c} 0 & 5 & 2 & 12 & 6 \\ 0 & 0 & 5 & 5 & 5 \\ 1 & 0 & 2\frac{1}{2} & 3\frac{1}{2} & 3\frac{1}{2} \\ 0 & 1 & 3 & 5 & 5 \end{array}\right) \xrightarrow[(0, -5, -15, -25)]{R_1 - 5R_4} \left(\begin{array}{cccc|c} 0 & 0 & -13 & -13 & -13 \\ 0 & 0 & 5 & 5 & 5 \\ 1 & 0 & 2\frac{1}{2} & 3\frac{1}{2} & 3\frac{1}{2} \\ 0 & 1 & 3 & 5 & 5 \end{array}\right) \xrightarrow[(0, 0, 10, 10)]{R_1 + 2R_2} \\ &\left(\begin{array}{cccc|c} 0 & 0 & -3 & -3 & -3 \\ 0 & 0 & 5 & 5 & 5 \\ 1 & 0 & 2\frac{1}{2} & 3\frac{1}{2} & 3\frac{1}{2} \\ 0 & 1 & 3 & 5 & 5 \end{array}\right) \xrightarrow[-\frac{1}{3}R_1]{R_4 + R_1} \left(\begin{array}{cccc|c} 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 5 & 5 & 5 \\ 1 & 0 & 2\frac{1}{2} & 3\frac{1}{2} & 3\frac{1}{2} \\ 0 & 1 & 0 & 2 & 2 \end{array}\right) \xrightarrow[R_2 - 5R_1]{R_3 - 2\frac{1}{2}R_1} \left(\begin{array}{cccc|c} 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 2 & 2 \end{array}\right) \rightarrow \\ &\left(\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 2 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{array}\right) \end{aligned}$$

This augmented matrix represents the following equations:

$$\begin{aligned} k_1 &= 1 \\ k_2 &= 2 \\ k_3 &= 1 \end{aligned}$$

This means that $\begin{bmatrix} 6 & 0 \\ 3 & 8 \end{bmatrix}$ is a linear combination of $\{A, B, C\}$, when $k_1 = 1, k_2 = 2$, and $k_3 = 1$. \square

10

In each part express the vector as a linear combination of $\vec{p}_1 = 2 + x + 4x^2$, $\vec{p}_2 = 1 - x + 3x^2$, and $\vec{p}_3 = 3 + 2x + 5x^2$.

a. $-9 - 7x - 15x^2$

d. $7 + 8x + 9x^2$

Section 1.2

4

In each part, suppose that the augmented matrix for a system of linear equations has been reduced by row operations to the given reduced row echelon form. Solve the system.

a. $\left[\begin{array}{cccc|c} 1 & 0 & 0 & -3 & \\ 0 & 1 & 0 & 0 & \\ 0 & 0 & 1 & 7 & \end{array}\right]$

b.
$$\begin{bmatrix} 1 & 0 & 0 & -7 & 8 \\ 0 & 1 & 0 & 3 & 2 \\ 0 & 0 & 1 & 1 & -5 \end{bmatrix}$$

c.
$$\begin{bmatrix} 1 & -6 & 0 & 0 & 3 & -2 \\ 0 & 0 & 1 & 0 & 4 & 7 \\ 0 & 0 & 0 & 1 & 5 & 8 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

d.
$$\begin{bmatrix} 1 & -3 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$