


SURVIVAL ANALYSIS


Peter Hall


Recruitment and follow-up





Average (mean) survival = 9 months


Recruitment and follow-up








Recruitment and follow-up




Recruitment and follow-up - reality



Average survival = 6 months


Analysis



Option 1 – ignore patients at lost-to-follow-up

7 6 5 4 3
■ ★ ■ ■ ★


2 1
■ ★



Option 1 – ignore patients at lost-to-follow-up

7 6 5 4 3
■ ★ ■ ■ ★


2 1
■ ★



Option 2 – assume patients experience event at loss-to-follow-up

7 6 5 4 3
■ ★ ★ ■ ★


2 1
■ ★



Option 2 – assume patients experience event at loss-to-follow-up

7 6 5 4 3
■ ★ ★ ■ ★


2 1
■ ★



Option 3 – Kaplan-Meier estimator

7 6 5 4 3
■ ★ ★ ■ ★


2 1
■ ★




Option 3 – Kaplan-Meier estimator

7 6 5 4 3
■ ★ ★ ■ ★

2 1
■ ★





Number at risk

No. at risk 850

463

207

94

27

9

0

Events

0

12

24

36

48

60

72

Time since diagnosis (months)


Tabulated K-M function

Interval (Start- End)	# At Risk at Start of Interval	# Censored During Interval	# At Risk at End of Interval	# Who Died at End of Interval	Proportion Surviving This Interval	Cumulative Survival at End of Interval
0-1	7	0	7	1	$6/7 = 0.86$	0.86
1-4	6	2	4	1	$3/4 = 0.75$	$0.86 * 0.75 = \mathbf{0.64}$
4-10	3	1	2	1	$1/2 = 0.5$	$0.86 * 0.75 * 0.5 = \mathbf{0.31}$
10-12	1	0	1	0	$1/1 = 1.0$	$0.86 * 0.75 * 0.5 * 1.0 = \mathbf{0.31}$

Life table

Time	Survival
0	1.0000
1	0.86
2	0.86
3	0.86
4	0.64
5	0.64
10	0.32
12	0.32

Cumulative incidence




Terminology

- survival analysis,
- survival modeling,
- survival data analysis,
- actuarial analysis,
- lifetable,
- life-history analysis,
- life data analysis,
- life-testing,
- failure-time data,
- event-history analysis,
- censored data analysis
- product limit estimator


= all the same thing

Summary measures


□ Median or mean or mode?




Summary measures




Summary measures




Summary measures



Summary measures



Comparing survival




Comparing survival


- Significant difference?
- Logrank test
 - gives a p-value
 - tests equality of survival function
- Stratified log-rank test
- Relies on proportional hazards

Sums the observed-minus-expected differences across all event times, then standardizes by the variance. If no difference, this follows a chi-squared distribution.


Proportional Hazards?




Proportional Hazards?




Summary measures




Proportional hazards?




Proportional hazards



Proportional hazards?



Proportional hazards?



Proportional hazards?





Fig. 3 - Survival curve for ejection fraction greater or lower than 0.30.

Proportional hazards?



Proportional hazards?






Figure 2: Effect of Therapy on Survival in Chronic Myelogenous Leukemia—Hydroxyurea or interferon-alpha vs related-donor transplantation. Adapted, with permission, from Gale et al (200).


Proportional hazards?




Proportional hazards?




Proportional hazards?



Proportional hazards?




Proportional hazards?



Proportional hazards?

□ Schoenfeld residuals

	chisq	df	p
age_at_diagnosis	0.140	1	0.708
ecog_ps	2.605	3	0.457
hormone_receptor	0.496	1	0.481
her2_status	4.851	1	0.028
n_metastatic_sites	2.461	1	0.117
liver_mets	3.673	1	0.055
brain_mets	1.496	1	0.221
GLOBAL	14.928	9	0.093



Size of difference?

- Cox-proportional hazards model
 - gives hazard ratio (= relative risk)
 - semi-parametric
 - doesn't specify underlying hazards but does assume proportionality
 - multivariable regression

Cox-regression

