



Overhang set report




EcoFlex level 2->3

There are 5 overhangs in this set. The restriction enzyme used for this set is **Bsal**.

Error! Palindromic overhang(s): GTAC; TCGA

Please see the Appendix on the last page for an explanation of details.





ATAG

CTAT

GC content: **25 %**.

Can form the following amino acids in 6 translation frames:

I[GEVDA]

[SPTG*ELKQVIRA]*


[NYDH][SR]

L[SF*LCYW]

[ISTFPGLCYVHNRDA]Y

[SPTA][IM]

Misannealing overhangs:





GTAC

GTAC

GC content: 50 %.

The overhang is palindromic, cannot be used for DNA assembly.

Can form the following amino acids in 6 translation frames:

V[PLQHR]

[SPTGA*ELKQVMRW]Y


[SRGC][T]

V[PLQHR]

[SPTGA*ELKQVMRW]Y

[SRGC][T]

Misannealing overhangs:





GGAC

GTCC

GC content: **75 %**.

Can form the following amino acids in 6 translation frames:

G[PLQHR]

[SPTGA*ELKQVMRW]D


[RGW][T]

V[PLQHR]

[SPTGA*ELKQVMRW]S

[SRGC][P]

Misannealing overhangs:





TCGA

TCGA


GC content: 50 %.

The overhang is palindromic, cannot be used for DNA assembly.

Can form the following amino acids in 6 translation frames:

S[ISTKMNR]
[ISTFPGLCYVHNRDA]R
[VIFL][DE]
S[ISTKMNR]
[ISTFPGLCYVHNRDA]R
[VIFL][DE]

Misannealing overhangs:





AACA

TGTT

GC content: **25 %**.

Can form the following amino acids in 6 translation frames:

N[ISTKMNR]

[SPTG*ELKQVIRA]T


[K*QE][QH]

C[SF*LCYW]

[ISTFPGLCYVHNRDA]V

[MVL][FL]

Misannealing overhangs:



Appendix

The report consists of 3 sections: results, overhangs, appendix.

Result page(s)

The first page describes the overhang set. The result is also summarised with a symbol:

☑ : good overhang set

⚠ : warning; there are ways to significantly improve the set

☒ : error: the set cannot be used for DNA assembly

Overhang pages

Each overhang is also analysed separately. The result is summarised with a symbol:

☑ : good overhang

☒ : unusable palindromic sequence

Overhangs are unpaired nucleotides at the end of a double-stranded linear DNA molecule. Overhangs can be on either strand; 5' or 3' overhangs. After DNA ligation with another DNA with a complementary overhang, these remain in the sequence as fusion sites ("scars").

Overhang sets

Use the [GoldenHinges](#) Python package to generate a set of mutually compatible overhangs that can be used for DNA assembly.