

2019ADS2 Week10 T test

by Wanlu Liu, wanluliu@intl.zju.edu.cn

2019-11-18

1. Introduction

This R Markdown file contains a tutorial of how to do different form of t test manually and in R. For those problems that need to workout manually, please include the formula and process of how you get the final number.

2. one sample t test

Let's say a gene called SOX17, from the published dataset, we found that its average expression in different lines (more than 30) of human embryonic stem cells is 8.9 (unit in RPKM). From your experiment, you used a new human embryonic stem cell line that is generated in ZJE, and you did RNA-seq of three biological replicates to check its expression in this cell line and found out its expression is 15, 30, 50 (RPKM). Is there enough evidence to show our embryonic stem cell's SOX17 expression is very different from others under significance level of 0.05?

Please write out the formula you used to calculate the statistics.

1. step 1 : State the hypotheses and identify the claim.
2. Step 2: what distribution to use?
3. Step 3: Find the critical value.
4. Step 4: Compute the test value.
5. Step 5: Make the decision.
6. Step 6: Summarize the results.

Please write out the R code calculate the statistics.

2. two sample t test

2.1 paired two sample t test

We followed a set of 5 patients with acute myeloid leukemia. We want to investigate whether the oncogene AML1 expression is repressed after a new treatment. Thus we tested their AML1 expression before and after the therapy. The gene expression level for AML1 before the treatment is:

$$x_1, x_2, x_3, x_4, x_5 = c(102, 340, 234, 332, 129)$$

. And the gene expression level for AML1 after the treatment is:

$$y_1, y_2, y_3, y_4, y_5 = c(74, 56, 70, 104, 11)$$

. Is there enough evidence to support the claim that the new treatment significantly reduce the AML1 expression level in acute myeloid leukemia patients under significance level of 0.05?

1. step 1 : State the hypotheses and identify the claim.
2. Step 2: what distribution to use?
3. Step 3: Find the critical value.
4. Step 4: Compute the test value.
5. Step 5: Make the decision.
6. Step 6: Summarize the results.

Please write out the R code calculate the statistics.

2.2 unpaired two sample t test

There are two types of human embryonic stem cells (naive vs primed). We have the RNAseq data for naive hESC (4 biological replicate replicates) and primed hESCs (4 biological replicates). In each RNAseq dataset, there are 23368 genes identified. We want to find out those genes that are significantly differential expressed (either up regulated or down regulated) under significance level of 0.05. Hint, use unpaired t test to find out genes with p-value less than 0.05.

```
geneexp=read.csv("week10_t_test_problemset_testdata.csv")
head(geneexp)
```

```
##          gname naive_hESC_r1 naive_hESC_r2 naive_hESC_r3 naive_hESC_r4
## 1 1/2-SBSRNA4      3.657      3.808      7.239      4.607
## 2      A1BG        0.038      0.035      0.146      0.028
## 3    A1BG-AS1      0.032      0.348      0.361      0.299
## 4      A1CF        0.004      0.000      0.006      0.003
## 5     A2LD1        0.490      0.404      0.192      0.137
## 6      A2M        0.087      0.067      0.089      0.063
##  primed_hESC_r1 primed_hESC_r2 primed_hESC_r3 primed_hESC_r4
## 1      4.429      8.190      3.364      6.431
## 2      0.120      0.096      0.000      0.034
## 3      0.331      0.356      0.201      0.527
## 4      0.026      0.036      0.009      0.006
## 5      0.264      0.315      0.120      0.221
## 6      0.549      0.801      0.521      0.728
```

```
tail(geneexp)
```

```
##          gname naive_hESC_r1 naive_hESC_r2 naive_hESC_r3 naive_hESC_r4
## 23363    ZXDC      5.472      7.170      6.914      6.781
## 23364   ZYG11A     38.996     40.783     43.214     34.863
## 23365   ZYG11B      9.271      9.741      7.855      9.625
## 23366     ZYX     11.731      7.882      7.685      8.729
## 23367   ZZEF1      9.343      9.973      7.814      7.089
## 23368    ZZZ3     20.996     14.505      9.251     11.654
##  primed_hESC_r1 primed_hESC_r2 primed_hESC_r3 primed_hESC_r4
## 23363      5.866      6.931      4.513      5.958
## 23364      4.714      7.006      3.506      4.975
## 23365      8.533      9.096      6.318      7.359
## 23366      3.450      2.543      1.304      3.481
## 23367      2.608      3.334      1.208      2.705
## 23368     18.732     16.479     16.563     17.609
```

```
dim(geneexp)
```

```
## [1] 23368      9
```

2.3 One step further - Multiple testing correction (Advanced thinking, optional)

2.3.1 Why Multiple Testing Matters?

Genomics usually have Lots of Data which means there will be lots of Hypothesis Tests in one experiment. For example, a typical RNAseq experiment might result in performing 20000 separate hypothesis tests (like what we did before). If we use a standard p-value cut-off of 0.05, we'd expect 1000 (20000×0.05) genes to be deemed 'significant' by chance (not reasonable). Thus we usually will perform multiple testing correction after we calculate p-value for genomics. You can refer to this coursera online course video if you are interested in <https://www.coursera.org/lecture/statistical-genomics/multiple-testing-8-25-NsJfs>

R Markdown

This is an R Markdown document. Markdown is a simple formatting syntax for authoring HTML, PDF, and MS Word documents. For more details on using R Markdown see <http://rmarkdown.rstudio.com>.