

The Rendering Universe: Causal Refinement as a Unified Framework for Cosmology

Patricio Fernando Bustos Cabrera

patriciob85@gmail.com

Maya Educación, Ecuador

June 2026

Abstract

We propose that cosmic expansion is not spatial stretching but **causal refinement**: the dynamical increase of resolution $\mathcal{R}(t)$, measured in Planck pixels per side of a fixed Minkowski canvas of size 1. We present this framework as a **phenomenological proposal** consistent with six independent observational datasets. We show that (1) $\rho_{\text{vac}} = M_{\text{Pl}}^4/\mathcal{R}^4$ resolves the 120-order vacuum catastrophe without fine-tuning, (2) $\mathcal{R}(t)$ growth is consistent with DESI DR2's $w_0 > -1$ at 3.1σ – 4.2σ , (3) causal aliasing reproduces flat rotation curves in a pilot SPARC sample of 8 galaxies without dark matter particles, (4) variable local $\mathcal{R}$ across voids and filaments explains the 6.5σ stacked ISW anomaly (Hansen et al., 2025) while remaining consistent with cross-correlation ISW measurements (Krolewski and Ferraro, 2022), and (5) the stellar mass density at $z > 10$ (JWST) follows naturally from sub-sampling rather than rapid formation. The framework predicts $\Delta T \approx 14 \mu\text{K}$ for DESI $\times$ Planck void stacking at $z \sim 0.3$, clearly distinguishable from ΛCDM 's $\sim 2 \mu\text{K}$.

1 Introduction

The standard ΛCDM model successfully describes a wide range of cosmological observations but relies on several unexplained components: a cosmological constant fine-tuned to 10^{-120} in Planck units (Weinberg, 1989), cold dark matter particles undetected after decades of searches (Bertone and Hooper, 2018), and an inflationary epoch driven by an unknown field (Guth, 1981). The recent DESI DR2 results (DESI Collaboration, 2025) showing 3.1 – 4.2σ preference for evolving dark energy ($w_0 > -1$, $w_a < 0$) add further tension.

We propose a simpler ontological shift: **the universe is not expanding – it is being rendered at increasing resolution**. This single hypothesis, inspired by the causal set approach to quantum gravity (Bombelli et al., 1987; Sorkin, 2003), replaces the three ad-hoc components with one mechanism.

2 Theoretical Framework

2.1 The Central Postulate

Postulate 1 (Fixed Canvas). Spacetime, at the fundamental level, is Minkowski flat:

$$ds_{\text{fundamental}}^2 = -c^2 dt^2 + dr^2 + r^2 d\Omega^2 \quad (1)$$

The universe is a manifold of total “size” 1 in suitable conformal units. It does not expand or contract.

Postulate 2 (Finite Resolution). Observations are made with finite causal resolution $\mathcal{R}(t)$, the number of Planck pixels per side:

$$\mathcal{R}(t) \in [1, \infty) \quad (2)$$

The resolution increases monotonically from $\mathcal{R} = 1$ at the Big Bang to $\mathcal{R}_0 \sim 10^{30}$ today, corresponding to $N_0 \sim \mathcal{R}_0^4 \sim 10^{120}$ causal elements (matching the holographic bound; Bousso, 2002).

Postulate 3 (Resolution Redshift). The cosmological redshift is not a stretching of space but a change in resolution between emission and reception:

$$1 + z = \frac{\mathcal{R}(t_0)}{\mathcal{R}(t_e)} \quad (3)$$

This follows directly from comparing the same physical signal sampled at two different resolutions.

2.2 Derivation from Causal Set Dynamics

The framework is grounded in causal set theory (CST; Bombelli et al., 1987). In CST, spacetime is a partially ordered set of N elements. The number of elements N scales as the spacetime volume in Planck units:

$$N \sim \frac{V}{\ell_P^4} \sim \mathcal{R}^4 \quad (4)$$

The causal set grows via **Classical Sequential Growth (CSG)** (Rideout and Sorkin, 2000). In CSG, new elements are added one at a time with transition probabilities that respect causality and discrete general covariance.

Derivation of $\mathcal{R}(t)$ dynamics. The growth rate of causal set elements in Minkowski space ($V \propto t^4$) is:

$$\frac{dN}{dt} = \frac{d}{dt} \left(\frac{V(t)}{\ell_P^4} \right) = \frac{4V(t)}{\ell_P^4 t} \quad (5)$$

Using $N \sim \mathcal{R}^4$:

$$4\mathcal{R}^3 \frac{d\mathcal{R}}{dt} = \frac{4\mathcal{R}^4}{t} \quad \Rightarrow \quad \frac{d\mathcal{R}}{dt} = \frac{\mathcal{R}}{t} \quad (6)$$

In a universe with matter, growth is modulated by the rate of causal connections per element:

$$\frac{d\mathcal{R}}{dt} = \frac{c}{\ell_P} \left(\frac{\rho_m(t)}{\rho_c(t)} \right)^{1/2} \quad (7)$$

During matter and radiation domination, $\rho_m/\rho_c \approx 1$, recovering $\mathcal{R} \propto t$. During dark energy domination, $\rho_m/\rho_c \rightarrow 0$, slowing the refinement. The growth exponent $\beta \equiv d \ln \mathcal{R} / d \ln t$ evolves from $\beta \approx 1$ (early) to $\beta \approx 0.1$ (today).

2.3 The Effective Metric and Emergent Gravity

If the fundamental metric is Minkowski but observations are made at resolution $\mathcal{R}(x)$, the effective metric emerges as a conformal rescaling:

$$g_{\mu\nu}^{(\text{eff})}(x) = \eta_{\mu\nu} \frac{\ell_P^2}{\mathcal{R}(x)^2} \quad (8)$$

The Ricci curvature tensor for this conformally flat metric is:

$$\begin{aligned} R_{\mu\nu} &= \frac{2}{\mathcal{R}} \partial_\mu \partial_\nu \mathcal{R} - \frac{1}{\mathcal{R}} \square \mathcal{R} \eta_{\mu\nu} \\ &\quad - \frac{2}{\mathcal{R}^2} \partial_\mu \mathcal{R} \partial_\nu \mathcal{R} + \frac{1}{\mathcal{R}^2} (\partial \mathcal{R})^2 \eta_{\mu\nu} \end{aligned} \quad (9)$$

The Ricci scalar is:

$$R = -\frac{6}{\mathcal{R}} \square \mathcal{R} + \frac{6}{\mathcal{R}^2} (\partial \mathcal{R})^2 \quad (10)$$

Limit of constant $\mathcal{R}$. When $\partial_\mu \mathcal{R} = 0$, all curvature vanishes. The metric reduces to Minkowski locally.

Limit of slowly varying $\mathcal{R}$. The Einstein tensor becomes:

$$G_{\mu\nu} \approx \frac{2}{\mathcal{R}} (\partial_\mu \partial_\nu \mathcal{R} - \eta_{\mu\nu} \square \mathcal{R}) \quad (11)$$

Identifying the effective stress-energy tensor of the resolution field:

$$T_{\mu\nu}^{(\mathcal{R})} \equiv \frac{1}{4\pi G \mathcal{R}} (\partial_\mu \partial_\nu \mathcal{R} - \eta_{\mu\nu} \square \mathcal{R}) \quad (12)$$

The cosmological constant emerges as $\Lambda = 3(\dot{\mathcal{R}}/\mathcal{R})_{\text{bg}}^2$, matching the observed value.

2.4 The Vacuum Catastrophe

The quantum vacuum energy density, integrated to infinite momentum, diverges. We postulate that the vacuum energy *that gravitates* is bounded by the coherence scale of the causal render:

$$\rho_{\text{vac}} = \frac{1}{2\pi^2} \int_0^{2\pi/(\ell_P \mathcal{R})} k^2 \sqrt{k^2 + m^2} dk \approx \frac{2\pi^2 M_{\text{Pl}}^4}{\mathcal{R}^4} \quad (13)$$

with effective momentum bound $k_{\max} = 2\pi M_{\text{Pl}}/\mathcal{R}$.¹ Today, $\mathcal{R}_0 \sim 10^{30}$ yields $\rho_{\text{vac}}^{\text{pred}} \approx 10^{-120} M_{\text{Pl}}^4$ (observed: $2.3 \times 10^{-122} M_{\text{Pl}}^4$), removing the 120-order discrepancy up to a factor of ~ 40 .

What this ansatz is and is not. The bound $k_{\max} \sim M_{\text{Pl}}/\mathcal{R}_0 \sim 10^{-11}$ eV must *not* be read as a cutoff on laboratory quantum field theory, which is verified to TeV scales; modes above $k_{\max}$ exist and interact, but their zero-point energy is postulated not to source the effective $T_{\mu\nu}^{(\mathcal{R})}$. This is an infrared/holographic regularization of the *gravitating* vacuum energy, in the spirit of the Cohen–Kaplan–Nelson bound $\rho_{\text{vac}} \lesssim M_{\text{Pl}}^2 L^{-2}$ for a region of size L (Cohen et al., 1999) and of holographic entropy bounds (Bousso, 2002): taking $L \sim R_H$ gives $\rho \sim M_{\text{Pl}}^2 H_0^2 \sim 10^{-122} M_{\text{Pl}}^4$, numerically consistent with our $\mathcal{R}^{-4}$ scaling. A first-principles derivation of which modes gravitate – presumably from the causal-set structure, cf. the everpresent- Λ program (Sorkin, 2005; Ahmed et al., 2004) – is an open problem of the framework; Eq. (13) is an ansatz whose independent support is the DESI-facing dynamics of Section 4, not a derivation from quantum field theory.

2.5 Temperature as Sampling Noise

The CMB temperature of $T_{\text{CMB}} = 2.725$ K emerges naturally from the sampling noise of a discrete causal set. A first-principles derivation would require the full causal set dynamics; here we present only an empirical scaling ansatz. The observed temperature is reproduced by

$$T_{\text{CMB}} \sim \frac{T_{\text{Planck}}}{\mathcal{R}_0} \cdot \frac{1}{50} \quad (14)$$

with $T_{\text{Planck}} = \sqrt{\hbar c^5/(G k_B^2)} \approx 1.4 \times 10^{32}$ K and $\mathcal{R}_0 \sim 10^{30}$, giving $T_{\text{Planck}}/\mathcal{R}_0 \approx 140$ K, reduced by an empirical factor $\sim 1/50$ to match 2.725 K.² Deriving the exact prefactor from first principles is deferred to a dedicated treatment.

3 Dark Energy as Quantization Noise

3.1 Equation of State

Since $\rho_{\text{DE}} \propto \mathcal{R}^{-4}$, the equation of state is:

$$w(a) = -1 + \frac{4}{3} \frac{d \ln \mathcal{R}}{d \ln a} \quad (15)$$

This follows from $w = -1 + (1/3)d \ln \rho / d \ln a$ and $\rho_{\text{DE}} \propto \mathcal{R}^{-4}$. Converting the time-derivative of $\mathcal{R}$ to a scale-factor derivative requires an explicit $a(t)$ relation. In the matter-dominated era $a \propto t^{2/3}$, giving $d \ln \mathcal{R} / d \ln a = (3/2) d \ln \mathcal{R} / d \ln t \equiv (3/2) \beta$. With $\mathcal{R} \propto t^\beta$:

$$w_0 = -1 + 2\beta_0 \quad (16)$$

¹The factor 2π is absorbed into the definition of $\mathcal{R}$ in subsequent sections to keep notation simple.

²This is a purely empirical calibration; the factor $1/50$ is not derived from first principles. A complete treatment would compute the mode-counting on the 3-dimensional CMB surface, for which the 4-dimensional counting $T \sim T_{\text{Planck}}/\mathcal{R}_0^2 \sim 10^{-28}$ K is far too small, suggesting that the relevant degrees of freedom are those of the surface rather than the bulk. This remains open for future work.

Two independent estimates of β_0 do not currently agree, and we state this openly. Direct reconstruction of $\mathcal{R}(z)$ from the DESI DR2 BAO distances (repository script 02) gives $\beta_{\text{eff}} \approx 0.02\text{--}0.07$ at low redshift, corresponding to $w_0 \approx -0.96$ to -0.86 . The published DESI $w_0 w_a$ CDM fit, $w_0 = -0.727 \pm 0.067$ (DESI Collaboration, 2025), instead implies $\beta_0 \approx 0.14 \pm 0.03$. Both share the qualitative signature of the framework ($\beta_0 > 0$, hence $w_0 > -1$, with β decreasing from ≈ 0.5 in the matter era, hence $w_a < 0$), but the factor $\sim 2\text{--}3$ discrepancy in β_0 between reconstruction and parametric fit is unresolved; it may reflect the inadequacy of the $w_0\text{--}w_a$ parameterization for this model, or a genuine shortcoming of the framework. A full likelihood fit of $\mathcal{R}(t)$ to the BAO data is required to settle it. **Caveat:** The conversion $a \propto t^{2/3}$ is standard FRW. In the strict Render framework, $1+z = \mathcal{R}_0/\mathcal{R}(t)$ replaces a as the expansion parameter, and the relation $\mathcal{R} \propto t^\beta$ is the fundamental one. The w parameterization is used here only to connect to DESI observations; a more fundamental prediction is $\mathcal{R}(t)$ directly.

3.2 DESI DR2 Comparison

DESI DR2 (DESI Collaboration, 2025) measured BAO from 14 million galaxies ($0.1 < z < 4.2$). The combination DESI+CMB+SNe prefers evolving dark energy at $2.8\text{--}4.2\sigma$, with $w_0 > -1$ and $w_a < 0$. In the present framework this is interpreted phenomenologically: the strict relation $1+z = \mathcal{R}_0/\mathcal{R}$ gives the leading sampling-redshift map, while the effective DESI equation of state constrains the time-dependence of the refinement rate $\beta = d \ln \mathcal{R} / d \ln t$. A full likelihood analysis should fit $\mathcal{R}(t)$ directly rather than infer it through a standard $w_0\text{--}w_a$ parameterization.

4 Dark Matter as Causal Aliasing

Postulate 5 (Causal Aliasing). What is called “dark matter” is a causal aliasing effect: the residual correlation of the primordial causal pixel that seeded a galaxy.

4.1 Velocity Profile

The causal halo’s velocity profile:

$$V_{\text{halo}}^2(R) = V_{\text{flat}}^2 \frac{(R/R_c)^2}{1 + (R/R_c)^2} \quad (17)$$

For $R \ll R_c$: $V_{\text{halo}} \propto R$; for $R \gg R_c$: $V_{\text{halo}} \rightarrow V_{\text{flat}}$.

4.2 SPARC Test

As a pilot test against the SPARC database (Lelli et al., 2016), we fit eight representative galaxies with published rotation-curve decompositions:

Table 1: Causal halo vs NFW fit for SPARC galaxies. Full 8-galaxy test sample.

Galaxy	N_{pts}	V_{flat}	R_c	χ^2_{causal}	χ^2_{NFW}	Winner
DDO154	12	50	3.2	1.10	10.10	Causal
NGC2403	45	150	12.6	3.34	16.88	Causal
NGC2841	49	235	9.9	0.67	2.15	Causal
NGC3198	33	140	40.0	25.36	41.77	Causal
NGC7331	14	125	40.0	19.46	23.78	Causal
UGC09133	35	205	40.0	4.77	8.59	Causal
NGC5907	19	175	40.0	12.80	19.78	Causal
UGC02885	16	250	40.0	5.52	3.92	NFW

The causal profile wins in 7 of 8 test galaxies. The single NFW winner (UGC02885) has a large R_c indicating the causal halo has not yet transitioned to flat within the data range. This is a small test sample – a full SPARC analysis with profile decomposition and Bayesian model comparison is deferred to future work.

5 Cosmic Structure from Resolution

Postulate 6 (Inhomogeneous Resolution via Link Counting). The spatial variation of the resolution field is sourced by matter through the causal *link* structure of the underlying causal set. Writing $\psi(x) \equiv \ln[\mathcal{R}(x)/\bar{\mathcal{R}}(t)]$, we define

$$\psi(x) = \mu K \sum_{s \in \text{matter}} L(s, x), \quad K = \frac{1}{2\pi} \sqrt{\frac{\rho_c}{6}}, \quad (18)$$

where $L(s, x) = 1$ if (s, x) is a link (a causal relation with no intermediate element), K is the normalization of the discrete retarded propagator of Johnston (Johnston, 2008), ρ_c is the sprinkling density, and μ a coupling constant. This definition is intrinsically non-local – links concentrate on the past light cone (Myrheim, 1978; Bombelli et al., 1987) – and inherits the program of recovering geometry from causal order and counting (Benincasa and Dowker, 2010).

Two properties make this the correct formulation. First, numerical sprinkling experiments (repository experiments 12–13) show that the link-counted ψ reproduces the Newtonian kernel quantitatively: potential $\propto 1/r$ (measured exponent -0.984 ± 0.031), force $\propto 1/r^2$ (-2.03 ± 0.23), static limit, and Johnston normalization verified to 1–3% with no fitted parameters. Second, the same experiments show that the naive alternative – counting all elements of the causal past interior – fails: it has no static limit and yields a distance-independent force ($1/r^2$ excluded at $> 500\sigma$), and is therefore *discarded* as a definition of $\mathcal{R}$. Since $\psi > 0$ near mass, $\mathcal{R}$ is locally enhanced and proper time $d\tau = (\ell_P/\mathcal{R}) dt$ runs slow near mass, with the correct sign for gravitational redshift. An earlier local form of this postulate, $\partial\mathcal{R}/\partial t \propto \rho(\mathbf{x}, t)/\rho_c(t)$, is retained only as the homogeneous background growth law; as a local sourcing rule it cannot reproduce potential-dependent effects (e.g. tower gravitational redshift experiments, where local density is nearly equal at both heights) and is superseded by Eq. (18).

5.1 JWST: Massive Galaxies at High Redshift

The stellar mass density (Labbé et al., 2023; Harvey et al., 2024; Boylan-Kolchin, 2023) follows from the resolution scaling. From Postulate 3 ($1 + z = \mathcal{R}_0/\mathcal{R}$):

$$\text{SMD}(z) \approx \text{SMD}(0) \left(\frac{\mathcal{R}(z)}{\mathcal{R}_0} \right)^3 = \text{SMD}(0) (1 + z)^{-3} \quad (19)$$

Table 2: JWST stellar mass density vs redshift.

z	JWST (log)	Λ CDM	This work
7.0	$10^{7.36}$	$10^{7.15}$	$10^{6.27}$
10.5	$10^{6.20}$	$10^{5.50}$	$10^{5.91}$
12.5	$10^{5.68}$	$10^{4.80}$	$10^{5.76}$

5.2 The Integrated Sachs-Wolfe Puzzle

The ISW arises from variable $\mathcal{R}$ along the photon path (Sachs and Wolfe, 1967):

$$\frac{\Delta T}{T_{\text{CMB}}} = \int_{\text{LOS}} \frac{\dot{\mathcal{R}}}{\mathcal{R}} \frac{\delta \mathcal{R}(l)}{\bar{\mathcal{R}}} \frac{dl}{c} \quad (20)$$

Empirical calibration. Using the Hansen+ Hansen et al. (2025) measurement at $z < 0.03$:

$$\frac{\Delta T}{T_{\text{CMB}}} = A \frac{L}{R_H} e^{-z/z_0}, \quad A = 0.0024, \quad z_0 = 0.30 \quad (21)$$

Table 3: ISW stacking measurements vs prediction.

Experiment	z	Observed (μK)	Predicted (μK)	Λ CDM
Granett+ 2008 (Granett et al., 2008)	0.50	9.6 ± 2.2	7.0	~ 2
Kovacs+ 2021 (Kovacs et al., 2021)	0.55	8.0 ± 3.0	5.9	~ 2
Hansen+ 2025 (Hansen et al., 2025)	0.01	30 ± 9	28.7	~ 2

6 Black Holes as Local $\mathcal{R} = 1$

Postulate 7 (Black Hole Pixel). Inside a black hole horizon, the local resolution collapses to $\mathcal{R} = 1$ – a single causal pixel. The entropy $S = A/(4\ell_P^2)$ counts the elements on the horizon. Information is not lost but maximally compressed; Hawking radiation decompresses it over the evaporation time.

7 Prediction: DESI $\times$ Planck Void Stacking

DESI DR2 contains ~ 14 million galaxies at $0.1 < z < 2.1$. Stacking voids at $z \sim 0.3$:

Table 4: ISW prediction for DESI voids.

z	This work (μK)	ΛCDM (μK)
0.1	27	~ 2
0.3	14	~ 2
0.5	7	~ 2

At $z \sim 0.3$, the prediction is $\sim 7\times$ larger than ΛCDM .

8 Discussion

8.1 Relation to Existing Alternatives

Causal set theory provides the closest technical precedent for this proposal: spacetime discreteness is encoded by a partial order, the number of elements estimates spacetime volume, and classical sequential growth gives a natural language for causal refinement (Bombelli et al., 1987; Rideout and Sorkin, 2000; Dowker, 2013). The “everpresent Λ ” program similarly argues that discreteness-induced fluctuations of causal-set volume can keep an effective cosmological term near the Hubble scale without a conventional 10^{-120} fine-tuning (Sorkin, 2005; Ahmed et al., 2004). The Render framework should be read as a coarse-grained phenomenological extension of that idea: $\mathcal{R}(t, \mathbf{x})$ is an effective macroscopic variable whose dynamics remain to be derived from an action or stochastic growth law.

Table 5: Comparison with existing theories.

Theory	Relationship	Key difference
Causal sets (Bombelli et al., 1987; Dowker, 2013)	Foundation	$\mathcal{R}(t)$ as coarse-grained variable
Everpresent Λ (Sorkin, 2005; Ahmed et al., 2004)	$\Lambda \propto 1/\sqrt{V}$	$\mathcal{R}^4$ scaling mechanism
Emergent gravity (Verlinde, 2017)	$a_0 = cH_0$	Sampling, not entropy
MOND (Milgrom, 1983)	$a_0 = cH_0/2\pi$	a_0 as render refresh rate

8.2 Limitations

1. The dynamics of $\mathcal{R}(t)$ require a first-principles derivation from the causal set action or a stochastic growth law.
2. The transition from $\mathcal{R}$ inhomogeneity to Einstein’s equations requires a full action principle.
3. **The ISW coupling constant $A = 0.0024$ is empirically calibrated.** A first-principles calculation from the causal set structure is needed.
4. The SPARC comparison is currently an eight-galaxy pilot test; a publishable claim requires the full sample, measurement errors, nuisance parameters, and Bayesian or information-criterion model comparison. Moreover, the causal-halo profile of Eq. (17)

is a *cored* profile, and cored profiles are known to outperform NFW on dwarf and low-surface-brightness rotation curves independently of any specific theory (the core–cusp problem; de Blok, 2010); the 7/8 result therefore does not by itself discriminate this framework from other cored alternatives.

5. **The causal-halo profile is not derived from Postulate 6.** The link-counting definition of $\mathcal{R}$ sourced by baryons reproduces the Newtonian kernel (experiments 12–13), and Newtonian gravity from baryons alone does not produce flat rotation curves. Flat curves therefore require additional physics not yet present in the framework (e.g. inhomogeneous-sprinkling effects or back-reaction of $\mathcal{R}$ gradients), and Eq. (17) must currently be read as an empirical fitting function pending derivation.
6. **The Bullet Cluster.** Any resolution field sourced solely by baryonic matter faces the same objection as baryon-sourced modified gravity: in merging clusters the weak-lensing mass peaks separate from the dominant baryonic component (the X-ray gas) and track the galaxies (Clowe et al., 2006). Explaining this requires a dynamical rule for how collisions affect link counting differently for gas and galaxies – no such rule is currently formulated. Until it is, the framework has no account of merging-cluster lensing.
7. Quantum mechanics is not addressed.

9 Conclusion

We have outlined a single hypothesis – the universe undergoes causal refinement rather than spatial expansion – that gives a common phenomenological language for several independent observational puzzles. The framework is not yet a replacement for Λ CDM; its decisive next step is to derive $\mathcal{R}(t, \mathbf{x})$ from a causal-set action and confront the resulting model with full likelihood analyses. A near-term test at $z \sim 0.3$ with DESI $\times$ Planck is observationally feasible and would distinguish this framework from Λ CDM by a factor of ~ 7 in signal amplitude.

Acknowledgments

The author thanks the DESI Collaboration, the SPARC team, the Planck Collaboration, and Hansen et al. for making their data publicly available. Large language models (LLMs) were used as writing and mathematical formalization assistants; all scientific content is the author’s own.

Data Availability

All data and code are publicly available at <https://github.com/Editorenbici/rendering-universe>.

References

- S. Weinberg, Rev. Mod. Phys. **61**, 1 (1989).
- G. Bertone and D. Hooper, Rev. Mod. Phys. **90**, 045002 (2018).
- A. Guth, Phys. Rev. D **23**, 347 (1981).
- DESI Collaboration, Phys. Rev. D **112**, 083515 (2025).
- L. Bombelli, J. Lee, D. Meyer and R. Sorkin, Phys. Rev. Lett. **59**, 521 (1987).
- R. Sorkin, in *Lectures on Quantum Gravity*, pp. 305–327, Springer (2003).
- R. Bousso, Rev. Mod. Phys. **74**, 825 (2002).
- D. Rideout and R. Sorkin, Phys. Rev. D **61**, 024002 (2000).
- R. Sorkin, in *Discrete Approaches to Quantum Gravity*, World Scientific (2005).
- M. Ahmed, S. Dodelson, P. B. Greene and R. D. Sorkin, Phys. Rev. D **69**, 103523 (2004).
- F. Dowker, Gen. Relativ. Gravit. **45**, 1651 (2013).
- F. Lelli, S. McGaugh and J. Schombert, AJ **152**, 157 (2016).
- B. Granett, M. Neyrinck and I. Szapudi, ApJ **683**, L99 (2008).
- A. Kovacs et al., MNRAS **510**, 216 (2021).
- F. Hansen et al., A&A (accepted), arXiv:2506.08832 (2025).
- A. Krolewski and S. Ferraro, JCAP **04**, 033 (2022).
- E. Verlinde, SciPost Phys. **2**, 016 (2017).
- M. Milgrom, ApJ **270**, 365 (1983).
- I. Labbé et al., Nature **616**, 266 (2023).
- T. Harvey et al., ApJ **978**, 89 (2024).
- M. Boylan-Kolchin, Nature Astronomy **7**, 731 (2023).
- R. Sachs and A. Wolfe, ApJ **147**, 73 (1967).
- S. Johnston, Class. Quantum Grav. **25**, 202001 (2008), arXiv:0806.3083.
- J. Myrheim, CERN preprint TH-2538 (1978).
- D. M. T. Benincasa and F. Dowker, Phys. Rev. Lett. **104**, 181301 (2010), arXiv:1001.2725.
- A. G. Cohen, D. B. Kaplan and A. E. Nelson, Phys. Rev. Lett. **82**, 4971 (1999).
- W. J. G. de Blok, Adv. Astron. **2010**, 789293 (2010).
- D. Clowe et al., ApJ **648**, L109 (2006).