

The Rendering Universe: Causal Refinement as a Unified Framework for Cosmology

Patricio Fernando Bustos Cabrera

patriciob85@gmail.com

Maya Educación, Ecuador

July 2026

Status note (2026-07-06). This framework paper predates the project's 2026-07-05 refoundation. Since then: the original Exp 17 ISW measurement was invalidated by a coordinate-frame bug (found and documented by this project) and re-measured with verified astrometry; the refutation stands, firmer (Exp 17b: $\Delta T = +0.37 \pm 0.93 \mu\text{K}$ over 1454 voids; the framework predictions are excluded at 20.7σ and 67σ). The repository file `FUNDAMENTOS.md` supersedes this document wherever they disagree. A technical companion (counting laws of links) and a methodological companion (the falsification arc) are in preparation.

Abstract

We propose that cosmic expansion is not spatial stretching but **causal refinement**: the dynamical increase of resolution $\mathcal{R}(t)$, measured in Planck pixels per side of a fixed Minkowski canvas of size 1. We present this as a **phenomenological effective framework** whose macro-dynamics are consistent with several independent datasets. We show that (1) $\rho_{\text{vac}} = M_{\text{Pl}}^4/\mathcal{R}^4$ resolves the 120-order vacuum catastrophe up to a factor ~ 40 (as an IR/holographic regularization of gravitating vacuum energy), (2) a direct $\mathcal{R} \propto t^\beta$ likelihood against the official DESI DR2 BAO data (13 points, full covariance) gives $\beta_0 = 0.055^{+0.045}_{-0.050}$, resolving an earlier estimator tension, though BAO alone prefer this over ΛCDM by only $\Delta\chi^2 \approx 1$, (3) a pilot SPARC comparison shows a two-parameter cored profile fits better than NFW in 7/8 galaxies (though cored profiles outperform NFW independently of theory and the profile is not yet derived from the link-counting Postulate 6), (4) the variable- $\mathcal{R}$ account of the stacked ISW anomaly made a sharp, pre-registered prediction – $\Delta T \approx -19 \mu\text{K}$ for the stack of the full DESIVAST DR1 void catalog against Planck – which the measurement **refuted**: $\Delta T = +0.37 \pm 0.93 \mu\text{K}$ over 1454 voids with verified astrometry, consistent with ΛCDM (repository experiment 17b, superseding the bug-invalidated experiment 17; falsifier 1 of Section 8.3 fired, and the ISW sector is retracted as calibrated), and (5) JWST high- z stellar mass densities are compatible with resolution sub-sampling only at $z \gtrsim 10$. A first-principles derivation of $\mathcal{R}(t, \mathbf{x})$ from causal-set dynamics remains the outstanding step; the surviving quantitative content is the BAO sector (2) and the numerically validated link-counting gravity of Postulate 6.

Claims overview

This short table is the referee shortcut: live claims, refuted claims, auxiliary tools, ansatz, and pending checks with their current state and file provenance.

Table 1: Live and closed claims for RENDER UNIVERSE v2.0 (2026-07-03).

Claim	Status	Evidence / File
$\epsilon_{\text{link}} = 3\sqrt{6}/(\sqrt{\rho} T^2)$	MEDIDO+DERIVADO	Exp 24 (bit-exact repro) + Exp
$v_{\text{link}} = \pi\sqrt{6}\sqrt{\rho} T^2$	MEDIDO+DERIVADO	Exp 18b: $p = 2.089 \pm 0.025$, am
Kernel $K(x, y)$ selector of covering relations	LIVE	Sec. 5.1; notes/kernel_paper_sec
$[\mathcal{R}] = 1$ dimensionless	LIVE	Sticky convention in notes/sym
Exp 22 interval metrics separation	LIVE	outputs/exp22_formal_run_2026-
Postulado 3' disformal metric	ANSATZ	notes/action_disforme_postulad
Factor 1/2 in attrition (Exp 17)	ABIERTO	Pending pencil check
ISW distinctive signature	REFUTADO-FIRME	Exp 17b: $+0.37 \pm 0.93 \mu\text{K}$, excl
DM aliasing (Bullet Cluster / SPARC 8)	ABIERTO / DÉBIL	notes/claims_table_2026-07-03.m

1 Introduction and context

The standard ΛCDM model successfully describes a wide range of cosmological observations but relies on several unexplained components: a cosmological constant fine-tuned to 10^{-120} in Planck units (Weinberg, 1989), cold dark matter particles undetected after decades of searches (Bertone and Hooper, 2018), and an inflationary epoch driven by an unknown field (Guth, 1981). The recent DESI DR2 results (DESI Collaboration, 2025) showing $3.1\text{--}4.2\sigma$ preference for evolving dark energy ($w_0 > -1$, $w_a < 0$) add further tension.

We propose a simpler ontological shift: **the universe is not expanding – it is being rendered at increasing resolution**. This single hypothesis, inspired by the causal set approach to quantum gravity (Bombelli et al., 1987; Sorkin, 2003), replaces the three ad-hoc components with one mechanism.

2 Theoretical Framework

2.1 The Central Postulate

Postulate 1 (Fixed Canvas). Spacetime, at the fundamental level, is Minkowski flat:

$$ds_{\text{fundamental}}^2 = -c^2 dt^2 + dr^2 + r^2 d\Omega^2 \quad (1)$$

The universe is a manifold of total “size” 1 in suitable conformal units. It does not expand or contract.

Postulate 2 (Finite Resolution). Observations are made with finite causal resolution $\mathcal{R}(t)$, the number of Planck pixels per side:

$$\mathcal{R}(t) \in [1, \infty) \quad (2)$$

The resolution increases monotonically from $\mathcal{R} = 1$ at the Big Bang to $\mathcal{R}_0 \sim 10^{30}$ today, corresponding to $N_0 \sim \mathcal{R}_0^4 \sim 10^{120}$ causal elements (matching the holographic bound; Bousso, 2002).

Postulate 3 (Resolution Redshift). The cosmological redshift is not a stretching of space but a change in resolution between emission and reception:

$$1 + z = \frac{\mathcal{R}(t_0)}{\mathcal{R}(t_e)} \quad (3)$$

This follows directly from comparing the same physical signal sampled at two different resolutions.

Postulate 3' (Effective Disformal Coupling). In the weak-field, slow-motion regime, matter couples to an effective metric induced by the resolution scalar $\psi = \nabla \ln \mathcal{R}(t)$:

$$d\tilde{s}^2 = -e^{-2\psi} c^2 dt^2 + e^{+2\psi} d\mathbf{x}^2, \quad (4)$$

with $\psi \rightarrow 0$ recovering Minkowski. The scalar field equation on flat background is

$$\square_\eta \psi + \rho e^{-\psi} = 0, \quad (5)$$

where we use units $4\pi G/c^2 = 1$ for brevity. In the linearized limit $|\psi| \ll 1$ this reduces to Poisson's equation with the identification $\Phi_N = -c^2\psi$. This postulate is auxiliary: it provides a controlled effective-field language for post-Newtonian corrections and screening, without claiming that ψ is a dynamically independent field. Precedents include the generic disformal coupling of Bekenstein (1993), Sakstein (2014), Bettoni–Liberati (2013), and Zumalacárregui (2010); the specific form $\psi = \nabla \ln \mathcal{R}(t)$ is the project-specific ansatz motivated by a fixed canvas of unit size.

2.2 Derivation from Causal Set Dynamics

The framework is grounded in causal set theory (CST; Bombelli et al., 1987). In CST, spacetime is a partially ordered set of N elements. The number of elements N scales as the spacetime volume in Planck units:

$$N \sim \frac{V}{\ell_P^4} \sim \mathcal{R}^4 \quad (6)$$

The causal set grows via **Classical Sequential Growth (CSG)** (Rideout and Sorkin, 2000). In CSG, new elements are added one at a time with transition probabilities that respect causality and discrete general covariance.

Derivation of $\mathcal{R}(t)$ dynamics. The growth rate of causal set elements in Minkowski space ($V \propto t^4$) is:

$$\frac{dN}{dt} = \frac{d}{dt} \left(\frac{V(t)}{\ell_P^4} \right) = \frac{4V(t)}{\ell_P^4 t} \quad (7)$$

Using $N \sim \mathcal{R}^4$:

$$4\mathcal{R}^3 \frac{d\mathcal{R}}{dt} = \frac{4\mathcal{R}^4}{t} \Rightarrow \frac{d\mathcal{R}}{dt} = \frac{\mathcal{R}}{t} \quad (8)$$

In a universe with matter, growth is modulated by the rate of causal connections per element:

$$\frac{d\mathcal{R}}{dt} = \frac{c}{\ell_P} \left(\frac{\rho_m(t)}{\rho_c(t)} \right)^{1/2} \quad (9)$$

During matter and radiation domination, $\rho_m/\rho_c \approx 1$, recovering $\mathcal{R} \propto t$. During dark energy domination, $\rho_m/\rho_c \rightarrow 0$, slowing the refinement. The growth exponent $\beta \equiv d \ln \mathcal{R} / d \ln t$ evolves from $\beta \approx 1$ (early) to $\beta_0 \approx 0.05\text{--}0.15$ today; the current spread in β_0 reflects a real tension between two estimators, discussed openly in Section 3.

2.3 The Effective Metric and Emergent Gravity

If the fundamental metric is Minkowski but observations are made at resolution $\mathcal{R}(x)$, the effective metric was originally taken to be a conformal rescaling:

$$g_{\mu\nu}^{(\text{eff})}(x) = \eta_{\mu\nu} \frac{\ell_P^2}{\mathcal{R}(x)^2} \quad (10)$$

Postulate 3' (Effective Metric, amended 2026-07-04). The render maps resolution to the effective metric *anisotropically*: clocks respond as $1/\mathcal{R}$ and rulers as $\mathcal{R}$. In the weak field, with $\psi \equiv \ln(\mathcal{R}/\bar{\mathcal{R}})$,

$$ds^2 = -e^{-2\psi} c^2 dt^2 + e^{+2\psi} d\mathbf{x}^2, \quad (11)$$

which yields $\gamma_{\text{PPN}} = 1$ by construction: gravitational light deflection, lensing and the Shapiro delay are recovered at linear order, and the Cassini bound $\gamma - 1 = (2.1 \pm 2.3) \times 10^{-5}$ is satisfied. The conformal form of Eq. (10) is retained *only* for the homogeneous cosmological background, where the two coincide up to a redefinition of the scale factor; as a local metric it gives $\gamma_{\text{PPN}} = -1$ – null geodesics are conformally invariant, so light would not bend at all – and is observationally excluded at the $\sim 10^5$ level. The amendment must apply universally (photons, gravitons and matter couple to the same $g_{\mu\nu}$), which keeps it consistent with the GW170817 bound $|c_{\text{GW}}/c - 1| < 10^{-15}$. The $\mathcal{O}(\psi^2)$ sector (PPN β , perihelion precession) and the derivation of the anisotropic response from causal-set counting remain open problems; a covariant completion is deferred to future work.

The curvature formulas below are derived for the conformal metric of Eq. (10) and therefore apply to the **homogeneous cosmological background only**; the local weak-field sector is governed by Postulate 3', Eq. (11). The Ricci curvature tensor for the conformally flat background metric is:

$$\begin{aligned} R_{\mu\nu} = & \frac{2}{\mathcal{R}} \partial_\mu \partial_\nu \mathcal{R} - \frac{1}{\mathcal{R}} \square \mathcal{R} \eta_{\mu\nu} \\ & - \frac{2}{\mathcal{R}^2} \partial_\mu \mathcal{R} \partial_\nu \mathcal{R} + \frac{1}{\mathcal{R}^2} (\partial \mathcal{R})^2 \eta_{\mu\nu} \end{aligned} \quad (12)$$

The Ricci scalar is:

$$R = -\frac{6}{\mathcal{R}} \square \mathcal{R} + \frac{6}{\mathcal{R}^2} (\partial \mathcal{R})^2 \quad (13)$$

Limit of constant $\mathcal{R}$. When $\partial_\mu \mathcal{R} = 0$, all curvature vanishes. The metric reduces to Minkowski locally.

Limit of slowly varying $\mathcal{R}$. The Einstein tensor becomes:

$$G_{\mu\nu} \approx \frac{2}{\mathcal{R}} (\partial_\mu \partial_\nu \mathcal{R} - \eta_{\mu\nu} \square \mathcal{R}) \quad (14)$$

Identifying the effective stress-energy tensor of the resolution field:

$$T_{\mu\nu}^{(\mathcal{R})} \equiv \frac{1}{4\pi G \mathcal{R}} (\partial_\mu \partial_\nu \mathcal{R} - \eta_{\mu\nu} \square \mathcal{R}) \quad (15)$$

The cosmological constant emerges as $\Lambda = 3(\dot{\mathcal{R}}/\mathcal{R})_{\text{bg}}^2$, matching the observed value.

2.4 The Vacuum Catastrophe

The quantum vacuum energy density, integrated to infinite momentum, diverges. We postulate that the vacuum energy *that gravitates* is bounded by the coherence scale of the causal render:

$$\rho_{\text{vac}} = \frac{1}{2\pi^2} \int_0^{2\pi/(\ell_P \mathcal{R})} k^2 \sqrt{k^2 + m^2} dk \approx \frac{2\pi^2 M_{\text{Pl}}^4}{\mathcal{R}^4} \quad (16)$$

with effective momentum bound $k_{\text{max}} = 2\pi M_{\text{Pl}}/\mathcal{R}$.¹ Today, $\mathcal{R}_0 \sim 10^{30}$ yields $\rho_{\text{vac}}^{\text{pred}} \approx 10^{-120} M_{\text{Pl}}^4$ (observed: $2.3 \times 10^{-122} M_{\text{Pl}}^4$), removing the 120-order discrepancy up to a factor of ~ 40 .

What this ansatz is and is not. The bound $k_{\text{max}} \sim M_{\text{Pl}}/\mathcal{R}_0 \sim 10^{-11}$ eV must *not* be read as a cutoff on laboratory quantum field theory, which is verified to TeV scales; modes above k_{max} exist and interact, but their zero-point energy is postulated not to source the effective $T_{\mu\nu}^{(\mathcal{R})}$. This is an infrared/holographic regularization of the *gravitating* vacuum energy, in the spirit of the Cohen–Kaplan–Nelson bound $\rho_{\text{vac}} \lesssim M_{\text{Pl}}^2 L^{-2}$ for a region of size L (Cohen et al., 1999) and of holographic entropy bounds (Bousso, 2002): taking $L \sim R_H$ gives $\rho \sim M_{\text{Pl}}^2 H_0^2 \sim 10^{-122} M_{\text{Pl}}^4$, numerically consistent with our $\mathcal{R}^{-4}$ scaling. A first-principles derivation of which modes gravitate – presumably from the causal-set structure, cf. the everpresent- Λ program (Sorkin, 2005; Ahmed et al., 2004) – is an open problem of the framework; Eq. (16) is an ansatz whose independent support is the DESI-facing dynamics of Section 4, not a derivation from quantum field theory.

2.5 Temperature as Sampling Noise

The CMB temperature of $T_{\text{CMB}} = 2.725$ K emerges naturally from the sampling noise of a discrete causal set. A first-principles derivation would require the full causal set dynamics; here we present only an empirical scaling ansatz. The observed temperature is reproduced by

$$T_{\text{CMB}} \sim \frac{T_{\text{Planck}}}{\mathcal{R}_0} \cdot \frac{1}{50} \quad (17)$$

with $T_{\text{Planck}} = \sqrt{\hbar c^5/(G k_B^2)} \approx 1.4 \times 10^{32}$ K and $\mathcal{R}_0 \sim 10^{30}$, giving $T_{\text{Planck}}/\mathcal{R}_0 \approx 140$ K, reduced by an empirical factor $\sim 1/50$ to match 2.725 K.² Deriving the exact prefactor from first principles is deferred to a dedicated treatment.

¹The factor 2π is absorbed into the definition of $\mathcal{R}$ in subsequent sections to keep notation simple.

²This is a purely empirical calibration; the factor $1/50$ is not derived from first principles. A complete

3 Dark Energy as Quantization Noise

3.1 Equation of State

Since $\rho_{\text{DE}} \propto \mathcal{R}^{-4}$, the equation of state is:

$$w(a) = -1 + \frac{4}{3} \frac{d \ln \mathcal{R}}{d \ln a} \quad (18)$$

This follows from $w = -1 + (1/3)d \ln \rho / d \ln a$ and $\rho_{\text{DE}} \propto \mathcal{R}^{-4}$. Converting the time-derivative of $\mathcal{R}$ to a scale-factor derivative requires an explicit $a(t)$ relation. In the matter-dominated era $a \propto t^{2/3}$, giving $d \ln \mathcal{R} / d \ln a = (3/2) d \ln \mathcal{R} / d \ln t \equiv (3/2) \beta$. With $\mathcal{R} \propto t^\beta$:

$$w_0 = -1 + 2\beta_0 \quad (19)$$

An earlier version of this section reported a factor ~ 2 – 3 tension between two estimates of β_0 : per-bin reconstruction of $\mathcal{R}(z)$ from the BAO distances gave $\beta_{\text{eff}} \approx 0.02$ – 0.07 , while mapping the published DESI $w_0 w_a$ CDM fit ($w_0 = -0.727 \pm 0.067$; DESI Collaboration, 2025) through $w_0 = -1 + 2\beta_0$ suggested $\beta_0 \approx 0.14$. We have now performed the direct likelihood fit that we had identified as the arbiter (repository experiment 15): the self-consistent model $\mathcal{R} \propto t^\beta$, $\rho_{\text{DE}} \propto \mathcal{R}^{-4}$, fit to the official DESI DR2 release of 13 BAO measurements (D_V, D_M, D_H over r_d) with the full published covariance, profiling over Ω_m and $H_0 r_d$. The result is $\beta_0 = 0.055^{+0.045}_{-0.050}$ (68%, BAO alone; $0.040^{+0.040}_{-0.040}$ with a Planck-like Ω_m prior), with excellent goodness of fit ($\chi^2/\text{dof} = 0.92$). The reconstruction band lies inside the 68% interval, while $\beta_0 = 0.14$ is excluded at 95% once the Ω_m prior is applied: the apparent tension was an artifact of translating a different model ($w_0 w_a$ CDM) fit to a different data combination (BAO+CMB+SNe) through the matter-era conversion. Two honest caveats follow. First, BAO alone prefer $\beta > 0$ over Λ CDM ($\beta = 0$, nested) by only $\Delta\chi^2 \approx 1$: the framework is *consistent with*, but not yet *preferred by*, these data. Second, the exact present-day conversion is $w_0 = -1 + (4/3)\beta_0/(H_0 t_0)$, giving $w_0^{\text{eff}} \approx -0.92$ at the best fit; the combined BAO+SNe likelihood required by falsifier 4 (Section 8.3) remains the outstanding test. **Caveat:** The conversion $a \propto t^{2/3}$ is standard FRW. In the strict Render framework, $1+z = \mathcal{R}_0/\mathcal{R}(t)$ replaces a as the expansion parameter, and the relation $\mathcal{R} \propto t^\beta$ is the fundamental one. The w parameterization is used here only to connect to DESI observations; a more fundamental prediction is $\mathcal{R}(t)$ directly.

3.2 DESI DR2 Comparison

DESI DR2 (DESI Collaboration, 2025) measured BAO from 14 million galaxies ($0.1 < z < 4.2$). The combination DESI+CMB+SNe prefers evolving dark energy at 2.8 – 4.2σ , with $w_0 > -1$ and $w_a < 0$. In the present framework this is interpreted phenomenologically: the strict relation $1+z = \mathcal{R}_0/\mathcal{R}$ gives the leading sampling-redshift map, while the effective DESI equation of state constrains the time-dependence of the refinement rate $\beta = d \ln \mathcal{R} / d \ln t$. A full likelihood analysis should fit $\mathcal{R}(t)$ directly rather than infer it through a standard w_0 – w_a parameterization.

treatment would compute the mode-counting on the 3-dimensional CMB surface, for which the 4-dimensional counting $T \sim T_{\text{Planck}}/\mathcal{R}_0^2 \sim 10^{-28}$ K is far too small, suggesting that the relevant degrees of freedom are those of the surface rather than the bulk. This remains open for future work.

4 Dark Matter as Causal Aliasing

Postulate 5 (Causal Aliasing). What is called “dark matter” is a causal aliasing effect: the residual correlation of the primordial causal pixel that seeded a galaxy.

4.1 Velocity Profile

The causal halo’s velocity profile:

$$V_{\text{halo}}^2(R) = V_{\text{flat}}^2 \frac{(R/R_c)^2}{1 + (R/R_c)^2} \quad (20)$$

For $R \ll R_c$: $V_{\text{halo}} \propto R$; for $R \gg R_c$: $V_{\text{halo}} \rightarrow V_{\text{flat}}$.

4.2 SPARC Test

As a pilot test against the SPARC database (Lelli et al., 2016), we fit eight representative galaxies with published rotation-curve decompositions:

Table 2: Causal halo vs NFW fit for SPARC galaxies. Full 8-galaxy test sample.

Galaxy	N_{pts}	V_{flat}	R_c	χ_{causal}^2	χ_{NFW}^2	Winner
DDO154	12	50	3.2	1.10	10.10	Causal
NGC2403	45	150	12.6	3.34	16.88	Causal
NGC2841	49	235	9.9	0.67	2.15	Causal
NGC3198	33	140	40.0	25.36	41.77	Causal
NGC7331	14	125	40.0	19.46	23.78	Causal
UGC09133	35	205	40.0	4.77	8.59	Causal
NGC5907	19	175	40.0	12.80	19.78	Causal
UGC02885	16	250	40.0	5.52	3.92	NFW

The causal profile wins in 7 of 8 test galaxies. The single NFW winner (UGC02885) has a large R_c indicating the causal halo has not yet transitioned to flat within the data range. This is a small test sample – a full SPARC analysis with profile decomposition and Bayesian model comparison is deferred to future work.

5 Cosmic Structure from Resolution

Postulate 6 (Inhomogeneous Resolution via Link Counting). The spatial variation of the resolution field is sourced by matter through the causal *link* structure of the underlying causal set. Writing $\psi(x) \equiv \ln[\mathcal{R}(x)/\bar{\mathcal{R}}(t)]$, we define

$$\psi(x) = \mu K \sum_{s \in \text{matter}} L(s, x), \quad K = \frac{1}{2\pi} \sqrt{\frac{\rho_c}{6}}, \quad (21)$$

where $L(s, x) = 1$ if (s, x) is a link (a causal relation with no intermediate element), K is the normalization of the discrete retarded propagator of Johnston (Johnston, 2008), ρ_c is the sprinkling density, and μ a coupling constant. This definition is intrinsically non-local – links concentrate on the past light cone (Myrheim, 1978; Bombelli et al., 1987) – and inherits the program of recovering geometry from causal order and counting (Benincasa and Dowker, 2010).

An earlier local form of this postulate, $\partial\mathcal{R}/\partial t \propto \rho(\mathbf{x}, t)/\rho_c(t)$, is retained only as the homogeneous background growth law; as a local sourcing rule it cannot reproduce potential-dependent effects (e.g. tower gravitational redshift experiments, where local density is nearly equal at both heights) and is superseded by Eq. (21).

5.1 Link Kernel and Newtonian Limit

The link-counted potential of Eq. (21) can be written as an integral over the causal past with an irreducible kernel selector. We define

$$K(x, y) = \begin{cases} \mathbf{1}_{\mathcal{L}}(y, x) \exp\left(-\rho \mathcal{V}[I(y, x)]\right), & y \prec x, \\ 0, & \text{otherwise,} \end{cases} \quad (22)$$

where $\mathbf{1}_{\mathcal{L}}(y, x) = 1$ iff (y, x) is a link, i.e. $y \prec x$ and no element z satisfies $y \prec z \prec x$; $\mathcal{V}[I(y, x)]$ is the Alexandrov interval volume; and the Poisson factor gives the probability that no intermediate element lies in the interval. The screened potential is then

$$\psi(x) = \mu K \int K(x, y) \rho_{\text{matter}}(y) d^4y, \quad (23)$$

where K is the Johnston normalization constant (Johnston, 2008). In the weak-field static limit, projecting Eq. (23) onto a time slice gives $\psi(r) \propto 1/r$: numerical experiments at fixed resolution reproduce the Newtonian exponent -1.03 ± 0.03 and force law -2.03 ± 0.23 with no fitted parameters (Johnston, 2008; Benincasa and Dowker, 2010). The irreducible selector is the operative content: removing non-link elements from the kernel leaves only the covering relations, which concentrate on the light cone and produce the familiar $1/r$ tail, whereas summing over the full causal past yields a distance-independent force and was discarded in the pre-registered comparison.

5.2 JWST: Massive Galaxies at High Redshift

The stellar mass density (Labbé et al., 2023; Harvey et al., 2024; Boylan-Kolchin, 2023) follows from the resolution scaling. From Postulate 3 ($1 + z = \mathcal{R}_0/\mathcal{R}$):

$$\text{SMD}(z) \approx \text{SMD}(0) \left(\frac{\mathcal{R}(z)}{\mathcal{R}_0} \right)^3 = \text{SMD}(0) (1 + z)^{-3} \quad (24)$$

Table 3: JWST stellar mass density vs redshift.

z	JWST (log)	Λ CDM	This work
7.0	$10^{7.36}$	$10^{7.15}$	$10^{6.27}$
10.5	$10^{6.20}$	$10^{5.50}$	$10^{5.91}$
12.5	$10^{5.68}$	$10^{4.80}$	$10^{5.76}$

We note the crossover openly: the $(1+z)^{-3}$ sub-sampling scaling outperforms the Λ CDM expectation only at $z \gtrsim 10$ (0.3 dex vs 0.9 dex deviation at $z = 12.5$), while at $z \approx 7$ it *underpredicts* the observed SMD by an order of magnitude, worse than Λ CDM. The natural reading is that Eq. (24) is at best the high-redshift limiting behavior, with standard hierarchical growth dominating by $z \sim 7$; a model combining both regimes is required before this can be claimed as evidence, and we list it as such rather than as a confirmed success.

5.3 The Integrated Sachs-Wolfe Puzzle

The ISW arises from variable $\mathcal{R}$ along the photon path (Sachs and Wolfe, 1967). The cross-correlation ISW signal is also consistent with this picture (Krolewski and Ferraro, 2022):

$$\frac{\Delta T}{T_{\text{CMB}}} = \int_{\text{LOS}} \frac{\dot{\mathcal{R}}}{\mathcal{R}} \frac{\delta \mathcal{R}(l)}{\bar{\mathcal{R}}} \frac{dl}{c} \quad (25)$$

Empirical calibration. Using the Hansen+ Hansen et al. (2025) measurement at $z < 0.03$:

$$\frac{\Delta T}{T_{\text{CMB}}} = A \frac{L}{R_H} e^{-z/z_0}, \quad A = 0.0024, \quad z_0 = 0.30 \quad (26)$$

Table 4: ISW stacking measurements vs prediction.

Experiment	z	Observed (μK)	Predicted (μK)	Λ CDM
Granett+ 2008 (Granett et al., 2008)	0.50	9.6 ± 2.2	7.0	~ 2
Kovacs+ 2021 (Kovacs et al., 2021)	0.55	8.0 ± 3.0	5.9	~ 2
Hansen+ 2025 (Hansen et al., 2025)	0.01	30 ± 9	28.7	~ 2

6 Black Holes as Local $\mathcal{R} = 1$

Postulate 7 (Black Hole Pixel). Inside a black hole horizon, the local resolution collapses to $\mathcal{R} = 1$ – a single causal pixel. The entropy $S = A/(4\ell_P^2)$ counts the elements on the horizon. Information is not lost but maximally compressed; Hawking radiation decompresses it over the evaporation time.

7 The DESI $\times$ Planck Void Stacking Test: Prediction Refuted

The original version of this section predicted $\Delta T \approx 14 \mu\text{K}$ (in magnitude) for void stacking at $z \sim 0.3$, $\sim 7\times$ larger than Λ CDM’s $\sim 2 \mu\text{K}$. **We performed this test** (repositi-

tory experiment 17): the full DESIVAST DR1 VoidFinder catalog (1,489 non-edge voids, $0.055 < z < 0.234$, median $R_{\text{eff}} = 22$ Mpc) was stacked against the Planck SMICA map with a compensated top-hat filter at each void’s rescaled angular radius. The analysis was pre-registered and frozen (with author and independent audit) before unblinding; the null distribution (1,000 random-position sets) and a $-10 \mu\text{K}$ injection test validated the pipeline before any real void position was examined.

Result:

$$\Delta T_{\text{stack}} = +1.24 \pm 1.46 \mu\text{K} \quad (460 \text{ voids with valid patches}), \quad (27)$$

consistent with zero and with ΛCDM . The frozen framework predictions $-18.5 \mu\text{K}$ (empirical calibration) and $-61.7 \mu\text{K}$ (cumulative-deficit mechanism of experiment 16) – are excluded at $\sim 13\sigma$ and $\sim 41\sigma$ respectively. Robustness splits (NGC only; R_{eff} above the median) are also null. Falsifier 1 of Section 8.3 has therefore **fired**: the variable- $\mathcal{R}$ account of the stacked ISW anomaly is refuted as calibrated. Because the coupling A was calibrated against extreme-supervoid measurements (Granett et al., 2008; Hansen et al., 2025), this null result on an unselected catalog also supports the interpretation that those anomalies reflect a-posteriori selection effects rather than a universal coupling; consequently the $\beta_{\text{ISW}}\text{--}\beta_{\text{BAO}}$ consistency reported in experiment 16 loses its evidential basis. Any revival of the ISW sector must be accompanied by a new pre-registered prediction (e.g. for an independent supervoid-selected sample) rather than a reinterpretation of this outcome.

8 Discussion

8.1 Relation to Existing Alternatives

Causal set theory provides the closest technical precedent for this proposal: spacetime discreteness is encoded by a partial order, the number of elements estimates spacetime volume, and classical sequential growth gives a natural language for causal refinement (Bombelli et al., 1987; Rideout and Sorkin, 2000; Dowker, 2013). The “everpresent Λ ” program similarly argues that discreteness-induced fluctuations of causal-set volume can keep an effective cosmological term near the Hubble scale without a conventional 10^{-120} fine-tuning (Sorkin, 2005; Ahmed et al., 2004). The Render framework should be read as a coarse-grained phenomenological extension of that idea: $\mathcal{R}(t, \mathbf{x})$ is an effective macroscopic variable whose dynamics remain to be derived from an action or stochastic growth law.

The initial condition $\mathcal{R} = 1$ has a precise counterpart in cyclic causal-set cosmology: a *post* – a single element through which every causal path passes – separates successive cosmic cycles in CSG models, and Sorkin’s “cosmic renormalization” shows that the effective dynamical parameters are renormalized at each passage through a post, providing a mechanism for parameter naturalness without fine-tuning (Sorkin, 2000; Dowker and Zalel, 2017); observables for such cyclic models have recently been constructed (Dowker and Zalel, 2023). A speculative extension of the present framework would identify the $\mathcal{R} \rightarrow \infty$ asymptotics of one cycle with the $\mathcal{R} = 1$ post of the next, in structural analogy with the conformal crossover of Penrose’s Conformal Cyclic Cosmology (Gurzadyan and Penrose, 2010). We note, however, that CCC’s proposed CMB signatures (low-variance concentric circles) were

not confirmed by independent reanalyses (Moss et al., 2011), and we do *not* adopt the cyclic identification here: the framework as formulated requires only a single monotonic epoch, and none of its falsifiers (Section 8.3) depends on it.

Table 5: Comparison with existing theories.

Theory	Relationship	Key difference
Causal sets (Bombelli et al., 1987; Dowker, 2013)	Foundation	$\mathcal{R}(t)$ as coarse-grained variable
Everpresent Λ (Sorkin, 2005; Ahmed et al., 2004)	$\Lambda \propto 1/\sqrt{V}$	$\mathcal{R}^4$ scaling mechanism
Emergent gravity (Verlinde, 2017)	$a_0 = cH_0$	Sampling, not entropy
MOND (Milgrom, 1983)	$a_0 = cH_0/2\pi$	a_0 as render refresh rate

8.2 Limitations

1. The dynamics of $\mathcal{R}(t)$ require a first-principles derivation from the causal set action or a stochastic growth law.
2. The transition from $\mathcal{R}$ inhomogeneity to Einstein’s equations requires a full action principle.
3. **The ISW coupling derivation is void following the experiment-17 refutation.** The analysis below is retained for the record, but its premise – that $A = 0.0024$ measured a universal coupling – did not survive the pre-registered stacking test (Section 7), which found no anomalous signal in an unselected void catalog. The $\beta_{\text{ISW}}\text{--}\beta_{\text{BAO}}$ consistency therefore no longer constitutes evidence. Original text follows. Repository experiment 16 shows that modeling a void as a region of *cumulative refinement deficit*, $\delta\mathcal{R}/\bar{\mathcal{R}} = \gamma\beta|\delta_0|I(z)$ with $I(z) = \int D dt/t$, yields $A = \gamma\beta^2|\delta_0|I(0)/(H_0t_0)$ with no new parameters; reproducing the calibrated $A = 0.0024$ requires $\beta_{\text{ISW}} = 0.045$, inside the 68% BAO interval of Section 3 ($\beta_{\text{BAO}} = 0.055^{+0.045}_{-0.050}$) – two independent datasets pointing to the same refinement rate. Three caveats: (i) the redshift dependence is *not* reproduced (the mechanism predicts mild growth with z , while the data decay as e^{-z/z_0} ; $z_0 = 0.30$ remains empirical, plausibly encoding void-catalog selection and $\delta_0(z)$ evolution); (ii) the mechanism is the cumulative cosmological deficit, *not* the static link potential of Postulate 6, which underpredicts the effect by four orders of magnitude – the theoretical consistency between the two sectors is an open requirement; (iii) the sourcing exponent $\gamma \in [1/2, 1]$ is not yet fixed by the theory.
4. The SPARC comparison is currently an eight-galaxy pilot test; a publishable claim requires the full sample, measurement errors, nuisance parameters, and Bayesian or information-criterion model comparison. Moreover, the causal-halo profile of Eq. (20) is a *cored* profile, and cored profiles are known to outperform NFW on dwarf and low-surface-brightness rotation curves independently of any specific theory (the core–cusp problem; de Blok, 2010); the 7/8 result therefore does not by itself discriminate this framework from other cored alternatives.

5. **The causal-halo profile is not derived from Postulate 6.** The link-counting definition of $\mathcal{R}$ sourced by baryons reproduces the Newtonian kernel (experiments 12–13), and Newtonian gravity from baryons alone does not produce flat rotation curves. Flat curves therefore require additional physics not yet present in the framework (e.g. inhomogeneous-sprinkling effects or back-reaction of $\mathcal{R}$ gradients), and Eq. (20) must currently be read as an empirical fitting function pending derivation.
6. **The Bullet Cluster.** Any resolution field sourced solely by baryonic matter faces the same objection as baryon-sourced modified gravity: in merging clusters the weak-lensing mass peaks separate from the dominant baryonic component (the X-ray gas) and track the galaxies (Clowe et al., 2006). Explaining this requires a dynamical rule for how collisions affect link counting differently for gas and galaxies – no such rule is currently formulated. Until it is, the framework has no account of merging-cluster lensing.
7. Quantum mechanics is not addressed.

8.3 What Would Falsify This Framework

A framework whose central variable parameterizes the whole cosmic history risks becoming unfalsifiable by reinterpretation. To guard against this, we state in advance the observations that would refute the framework *as formulated in this paper*, without appeal to new parameters or post-hoc modifications:

1. **DESI×Planck void stacking at $z \sim 0.3$ yields $\Delta T \approx 2 \mu\text{K}$.** The flagship prediction of Section 7 was $\approx 14 \mu\text{K}$ in magnitude, a factor ~ 7 above ΛCDM . A stacking measurement consistent with ΛCDM refutes the variable- $\mathcal{R}$ account of the ISW anomaly. **STATUS (2026-07-02): FIRED.** The pre-registered measurement (Section 7; repository experiment 17) gave $+1.24 \pm 1.46 \mu\text{K}$, consistent with ΛCDM ; the framework predictions were excluded at $\gtrsim 13\sigma$. The ISW sector is retracted as calibrated. We record the outcome here rather than deleting the falsifier, as the falsification discipline requires.
2. **A confirmed phantom equation of state, $w_0 < -1$.** Postulate 2 requires $\mathcal{R}(t)$ to grow monotonically, hence $\beta_0 > 0$ and $w_0 = -1 + 2\beta_0 \geq -1$ with no adjustable freedom. A robust cosmological convergence on $w_0 < -1$ (or a confirmed phantom crossing) falsifies the dark-energy mechanism outright. This is the cleanest falsifier: it depends on no fitted parameter of the framework.
3. **Direct detection of a dark matter particle** with the relic abundance required by ΛCDM . Postulate 5 attributes the missing-mass phenomenology to causal aliasing rather than particles; a confirmed particle detection removes its motivation and the framework’s dark-matter sector with it.
4. **An irreconcilable β_0 .** Section 3 reports a factor ~ 2 – 3 tension between the BAO reconstruction of β_0 (≈ 0.02 – 0.07) and the value implied by the DESI parametric fit (≈ 0.14). We commit to the direct $\mathcal{R}(t)$ likelihood analysis as the arbiter: if no single

monotonic $\mathcal{R}(t)$ fits the combined BAO+SNe data, the quantitative dark-energy sector fails even though the qualitative quadrant ($w_0 > -1$, $w_a < 0$) matches.

5. **Energy-dependent photon dispersion.** The framework as formulated assumes the underlying sprinkling preserves local Lorentz invariance (Bombelli et al., 1987), so photon propagation speed must be independent of energy and of local $\mathcal{R}$ to current experimental precision. Fermi-LAT gamma-ray-burst bounds already constrain linear Lorentz violation beyond the Planck scale; if any future refinement of the render dynamics predicts energy-dependent arrival times, those bounds apply immediately. Conversely, a confirmed detection of Lorentz-violating dispersion would require a fundamental revision, since the fixed Minkowski canvas (Postulate 1) provides no mechanism for it.

Each rescue of the framework from one of these outcomes would have to generate new testable predictions of its own; rescues that merely reinterpret a failure are not admissible. We adopted this discipline internally as well: the link-counting formulation of Postulate 6 replaced an earlier local form precisely because pre-registered numerical experiments (repository experiments 12–13) refuted the original mechanism, and the negative results are published in the repository alongside the positive ones.

9 Conclusion

We have outlined a single hypothesis – the universe undergoes causal refinement rather than spatial expansion – and subjected it to its own falsifiability program. The outcome to date is mixed and is reported without adjustment. Refuted: the variable- $\mathcal{R}$ account of the stacked ISW anomaly, by our own pre-registered DESI×Planck measurement (Section 7), which came out consistent with Λ CDM. Surviving: the direct $\mathcal{R} \propto t^\beta$ likelihood against DESI DR2 BAO ($\beta_0 = 0.055^{+0.045}_{-0.050}$, consistent with but not preferred over Λ CDM), and the numerically validated link-counting formulation of Postulate 6, which reproduces the Newtonian kernel from causal structure with no fitted parameters. Open: the first-principles derivation of $\mathcal{R}(t, \mathbf{x})$ from a causal-set action, the dark-matter sector, and the reconciliation of the sampling and refinement roles of $\mathcal{R}$. The framework is not a replacement for Λ CDM. We submit that its chief contribution, beyond the surviving sectors, is methodological: a speculative cosmology can and should ship with its own falsifiers, and record their outcomes when they fire.

Acknowledgments

The author thanks the DESI Collaboration, the SPARC team, the Planck Collaboration, and Hansen et al. for making their data publicly available. Large language models (LLMs) were used as writing and mathematical formalization assistants; all scientific content is the author’s own.

Data Availability

All data and code are publicly available at <https://github.com/Editorenbici/rendering-universe>.

References

- S. Weinberg, Rev. Mod. Phys. **61**, 1 (1989).
- G. Bertone and D. Hooper, Rev. Mod. Phys. **90**, 045002 (2018).
- A. Guth, Phys. Rev. D **23**, 347 (1981).
- DESI Collaboration, Phys. Rev. D **112**, 083515 (2025).
- L. Bombelli, J. Lee, D. Meyer and R. Sorkin, Phys. Rev. Lett. **59**, 521 (1987).
- R. Sorkin, in *Lectures on Quantum Gravity*, pp. 305–327, Springer (2003).
- R. Bousso, Rev. Mod. Phys. **74**, 825 (2002).
- D. Rideout and R. Sorkin, Phys. Rev. D **61**, 024002 (2000).
- R. Sorkin, in *Discrete Approaches to Quantum Gravity*, World Scientific (2005).
- M. Ahmed, S. Dodelson, P. B. Greene and R. D. Sorkin, Phys. Rev. D **69**, 103523 (2004).
- F. Dowker, Gen. Relativ. Gravit. **45**, 1651 (2013).
- F. Lelli, S. McGaugh and J. Schombert, AJ **152**, 157 (2016).
- B. Granett, M. Neyrinck and I. Szapudi, ApJ **683**, L99 (2008).
- A. Kovacs et al., MNRAS **510**, 216 (2021).
- F. Hansen et al., A&A (accepted), arXiv:2506.08832 (2025).
- A. Krolewski and S. Ferraro, JCAP **04**, 033 (2022).
- E. Verlinde, SciPost Phys. **2**, 016 (2017).
- M. Milgrom, ApJ **270**, 365 (1983).
- I. Labbé et al., Nature **616**, 266 (2023).
- T. Harvey et al., ApJ **978**, 89 (2024).
- M. Boylan-Kolchin, Nature Astronomy **7**, 731 (2023).
- R. Sachs and A. Wolfe, ApJ **147**, 73 (1967).
- S. Johnston, Class. Quantum Grav. **25**, 202001 (2008), arXiv:0806.3083.

- J. Myrheim, CERN preprint TH-2538 (1978).
- D. M. T. Benincasa and F. Dowker, Phys. Rev. Lett. **104**, 181301 (2010), arXiv:1001.2725.
- A. G. Cohen, D. B. Kaplan and A. E. Nelson, Phys. Rev. Lett. **82**, 4971 (1999).
- W. J. G. de Blok, Adv. Astron. **2010**, 789293 (2010).
- D. Clowe et al., ApJ **648**, L109 (2006).
- R. D. Sorkin, Int. J. Theor. Phys. **39**, 1731 (2000), gr-qc/0003043.
- F. Dowker and S. Zalel, C. R. Physique **18**, 246 (2017), arXiv:1703.07556.
- F. Dowker and S. Zalel, Class. Quantum Grav. **40**, 155015 (2023), arXiv:2212.01149.
- V. G. Gurzadyan and R. Penrose, arXiv:1011.3706 (2010).
- A. Moss, D. Scott and J. P. Zibin, JCAP **04**, 033 (2011), arXiv:1012.1305.