

Teoria dei sistemi.

State Space Analysis

Luigi Palopoli

December 23, 2016

Table of contents

Discrete Time Linear System Solution

Connection between the State Space Representation and the Z-Transform

The Role of the Eigenvalues in the Discrete Time Transfer Function

The Role of the Eigenvalues for Discrete Time Systems

Matrix Powers for diagonal Matrices

Matrix Powers for Diagonal Matrices with Complex Eigenvalues

Exponential Matrix of Defective Matrices

Matrix Powers of Defective Matrices with Complex Eigenvalues

Modes Analysis

Continuous Time Systems

The discrete time case

- ▶ Consider the evolution

$$\begin{aligned}x(t+1) &= Ax(t) + Bu(t) \\ y(t) &= Cx(t) + Du(t).\end{aligned}$$

The discrete time case

- Consider the evolution

$$\begin{aligned}x(t+1) &= Ax(t) + Bu(t) \\ y(t) &= Cx(t) + Du(t).\end{aligned}$$

- By iterative solution we find

$$x(1) = Ax(0) + Bu(0),$$

$$x(2) = Ax(1) + Bu(1) = A^2x(0) + ABu(0) + Bu(1),$$

$$\vdots$$

$$x(k) = Ax(k-1) + Bu(k-1) = A^kx(0) + A^{k-1}Bu(0) + \cdots + ABu(k-2) + Bu(k-1)$$

The discrete time case

- ▶ Consider the evolution

$$\begin{aligned}x(t+1) &= Ax(t) + Bu(t) \\ y(t) &= Cx(t) + Du(t).\end{aligned}$$

- ▶ By iterative solution we find

$$x(1) = Ax(0) + Bu(0),$$

$$x(2) = Ax(1) + Bu(1) = A^2x(0) + ABu(0) + Bu(1),$$

$$\vdots$$

$$x(k) = Ax(k-1) + Bu(k-1) = A^kx(0) + A^{k-1}Bu(0) + \dots + ABu(k-2) + Bu(k-1)$$

- ▶ ...which can be written as

$$x(k) = A^kx(0) + \sum_{i=0}^{k-1} A^{k-1-i}Bu(i), \tag{1}$$

$$y(k) = CA^kx(0) + C \sum_{i=0}^{k-1} A^{k-1-i}Bu(i) + Du(k).$$

Observations

Free and forced evolution

As for the continuous-time systems, we can identify a free evolution (depending on the initial condition $x(0)$) and a forced evolution (resulting from the application of the input function).

Observations

Free and forced evolution

As for the continuous-time systems, we can identify a free evolution (depending on the initial condition $x(0)$) and a forced evolution (resulting from the application of the input function).

Role of A^k

- ▶ Since A^k is a constant once k is fixed, it turns out that the states are combined through a time invariant linear transformation.
- ▶ The combination of forced and unforced response is an application of the *superposition principle* between the states and the inputs.

Connection with Z-transform

- Consider the autonomous system:

$$x(k+1) = Ax(k),$$

whose Z-Transform are given by

$$\mathcal{Z}(x(k+1)) = zX(z) - zx(0), \text{ and } \mathcal{Z}(Ax(k)) = AX(z),$$

that has been obtained by applying the *time-shifting rule*.

Connection with Z-transform

- Consider the autonomous system:

$$x(k+1) = Ax(k),$$

whose Z-Transform are given by

$$\mathcal{Z}(x(k+1)) = zX(z) - zx(0), \text{ and } \mathcal{Z}(Ax(k)) = AX(z),$$

that has been obtained by applying the *time-shifting rule*.

- Hence,

$$\begin{aligned} zX(z) - zx(0) &= AX(z) \Rightarrow (zI - A)X(z) = zx(0) \\ \Rightarrow X(z) &= (zI - A)^{-1}zx(0). \end{aligned}$$

Inverse Z-transform

- ▶ The sequence $x(k)$ can be recovered via the inverse Z-transform.

$$x(k) = \mathcal{Z}^{-1} (z(zI - A)^{-1}) x(0),$$

Proposition

The matrix A^k can be computed through the inverse of Z-Transform:

$$A^k = \mathcal{Z}^{-1} (z(zI - A)^{-1}).$$

Example

Harmonic Oscillator

Consider the following *harmonic oscillator*

$$x(k+1) = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} x(k) = Ax.$$

Compute the evolution.

- ▶ Since

$$zI - A = \begin{bmatrix} z & 1 \\ -1 & z \end{bmatrix},$$

- ▶ we have that the eigenvalues are $\pm j$ and hence

$$(zI - A)^{-1} = \frac{1}{z^2 + 1} \begin{bmatrix} z & -1 \\ 1 & z \end{bmatrix}$$

$$\begin{aligned} A^k &= \mathcal{Z}^{-1} (z(zI - A)^{-1}) \\ &= \mathcal{Z}^{-1} \left(\begin{bmatrix} \frac{z^2}{z^2+1} & -\frac{z}{z^2+1} \\ \frac{z}{z^2+1} & \frac{z^2}{z^2+1} \end{bmatrix} \right) \end{aligned}$$

Example

► ...finally...

$$A^k = \begin{bmatrix} \cos(\frac{\pi}{2}k) & -\sin(\frac{\pi}{2}k) \\ \sin(\frac{\pi}{2}k) & \cos(\frac{\pi}{2}k) \end{bmatrix}$$

Discrete convolution Summation

- ▶ The computation of the forced response requires the extension to matrices of the *discrete convolution sum*

Discrete convolution Summation

- ▶ The computation of the forced response requires the extension to matrices of the *discrete convolution sum*
- ▶ Consider a SISO system and assume that $u(\bar{k}) = 0 \ \forall \bar{k} < 0$, we can write

$$\sum_{i=0}^{k-1} A^{k-1-i} B u(i) = A^{k-1} * B u(k),$$

Discrete convolution Summation

- ▶ The computation of the forced response requires the extension to matrices of the *discrete convolution sum*
- ▶ Consider a SISO system and assume that $u(\bar{k}) = 0 \ \forall \bar{k} < 0$, we can write

$$\sum_{i=0}^{k-1} A^{k-1-i} Bu(i) = A^{k-1} * Bu(k),$$

- ▶ As a consequence,

$$\mathcal{Z} \left(A^{k-1} * Bu(k) \right) = (zI - A)^{-1} BU(z).$$

Discrete convolution Summation

- ▶ Applying the superposition principle, we get

$$\mathcal{Z}(y(k)) = Cz(zI - A)^{-1}x(0) + C(zI - A)^{-1}BU(z) + DU(z).$$

Discrete convolution Summation

- ▶ Applying the superposition principle, we get

$$\mathcal{Z}(y(k)) = Cz(zI - A)^{-1}x(0) + C(zI - A)^{-1}BU(z) + DU(z).$$

- ▶ If we consider $x(0) = 0$ we find the expression for the transfer function.

$$\frac{Y(z)}{U(z)} = C(zI - A)^{-1}B + D.$$

Example

Second order system

For the linear system

$$\begin{aligned}x(k+1) &= \begin{bmatrix} 0.5 & 1 \\ 0 & -0.5 \end{bmatrix} x(k) + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u(k), \\ y(k) &= [1 \quad -1] x(k),\end{aligned}$$

compute the transfer function.

Example

Second order system

For the linear system

$$\begin{aligned}x(k+1) &= \begin{bmatrix} 0.5 & 1 \\ 0 & -0.5 \end{bmatrix} x(k) + \begin{bmatrix} 0 \\ 1 \end{bmatrix} u(k), \\ y(k) &= [1 \quad -1] x(k),\end{aligned}$$

compute the transfer function.

- The result is in the following function:

$$G(z) = [1 \quad -1] (zI - A)^{-1} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \frac{-z + 1.5}{z^2 - 0.25}$$

The Role of the Eigenvalues in the discrete transfer function

- ▶ We know that for the transfer function $G(z)$ the poles play a prominent role

The Role of the Eigenvalues in the discrete transfer function

- ▶ We know that for the transfer function $G(z)$ the poles play a prominent role
- ▶ Let us see how this role is transferred to the eigenvalues for the state space.

The Role of the Eigenvalues in the discrete transfer function

- ▶ We know that for the transfer function $G(z)$ the poles play a prominent role
- ▶ Let us see how this role is transferred to the eigenvalues for the state space.
- ▶ Hence, assuming $x(0) = 0$ as in the transfer function analysis case, we have

$$Y(z) = C(zI - A)^{-1}BU(z) + DU(z),$$

where D can be neglected since it does not involve the system characteristics.

The Role of the Eigenvalues in the discrete transfer function

- ▶ We know that for the transfer function $G(z)$ the poles play a prominent role
- ▶ Let us see how this role is transferred to the eigenvalues for the state space.
- ▶ Hence, assuming $x(0) = 0$ as in the transfer function analysis case, we have

$$Y(z) = C(zI - A)^{-1}BU(z) + DU(z),$$

where D can be neglected since it does not involve the system characteristics.

- ▶ We can restrict to $G(z) = C(zI - A)^{-1}BU(z)$.

The Role of the Eigenvalues in the discrete transfer function

- ▶ the (i, j) element of $(zI - A)^{-1}$ can be computed using the Cramer's rule

$$(-1)^{i+j} \frac{\det \text{Adj}_{i,j}}{\mathcal{P}(A)},$$

where $\mathcal{P}(A)$ is the *characteristic polynomial* of A , defined as

$$\mathcal{P}(A) = \det(zI - A).$$

- ▶ $\det \text{Adj}_{i,j}$ has degree less than n and there is still the possibility of having a *cancellation* between the roots of $\mathcal{P}(A)$ and the roots of $\det \text{Adj}_{i,j}$.

The Role of the Eigenvalues for Discrete Time Systems

Let us start with some properties of the matrix power and its eigenvalues that will be useful:

- ▶ $A_1^k A_2^k = A_2^k A_1^k = (A_1 A_2)^k, \forall k$, iff $A_1 A_2 = A_2 A_1$;

The Role of the Eigenvalues for Discrete Time Systems

Let us start with some properties of the matrix power and its eigenvalues that will be useful:

- ▶ $A_1^k A_2^k = A_2^k A_1^k = (A_1 A_2)^k, \forall k$, iff $A_1 A_2 = A_2 A_1$;
- ▶ If A is invertible, then $(A^k)^{-1} = (A^{-1})^k = A^{-k}$;

The Role of the Eigenvalues for Discrete Time Systems

Let us start with some properties of the matrix power and its eigenvalues that will be useful:

- ▶ $A_1^k A_2^k = A_2^k A_1^k = (A_1 A_2)^k, \forall k$, iff $A_1 A_2 = A_2 A_1$;
- ▶ If A is invertible, then $(A^k)^{-1} = (A^{-1})^k = A^{-k}$;
- ▶ If $v \in \mathbb{R}^n$ is an *eigenvector* of a matrix A associated with the eigenvalue λ , i.e., $Av = \lambda v$, then it immediately follows that v is also an eigenvector of the matrix A^k associated with the eigenvalue λ^k , i.e., $A^k v = \lambda^k v$;

The Role of the Eigenvalues for Discrete Time Systems

Let us start with some properties of the matrix power and its eigenvalues that will be useful:

- ▶ $A_1^k A_2^k = A_2^k A_1^k = (A_1 A_2)^k, \forall k$, iff $A_1 A_2 = A_2 A_1$;
- ▶ If A is invertible, then $(A^k)^{-1} = (A^{-1})^k = A^{-k}$;
- ▶ If $v \in \mathbb{R}^n$ is an *eigenvector* of a matrix A associated with the eigenvalue λ , i.e., $Av = \lambda v$, then it immediately follows that v is also an eigenvector of the matrix A^k associated with the eigenvalue λ^k , i.e., $A^k v = \lambda^k v$;
- ▶ For any given A and for any transformation matrix T , we have that $(T^{-1}AT)^k = T^{-1}A^k T$;

The Role of the Eigenvalues for Discrete Time Systems

Let us start with some properties of the matrix power and its eigenvalues that will be useful:

- ▶ $A_1^k A_2^k = A_2^k A_1^k = (A_1 A_2)^k, \forall k$, iff $A_1 A_2 = A_2 A_1$;
- ▶ If A is invertible, then $(A^k)^{-1} = (A^{-1})^k = A^{-k}$;
- ▶ If $v \in \mathbb{R}^n$ is an *eigenvector* of a matrix A associated with the eigenvalue λ , i.e., $Av = \lambda v$, then it immediately follows that v is also an eigenvector of the matrix A^k associated with the eigenvalue λ^k , i.e., $A^k v = \lambda^k v$;
- ▶ For any given A and for any transformation matrix T , we have that $(T^{-1}AT)^k = T^{-1}A^k T$;
- ▶ $\det(A^k) = [\det(A)]^k$. From this property, from $\det(AB) = \det(A)\det(B)$ it follows that $\det(AA^{-1}) = \det(A)\det(A^{-1}) = 1$, which is of course $\det(I) = 1$. Hence, the *eigenvalues does not change for similarity transformation*.

Matrix Powers for Diagonal Matrices

- ▶ If a matrix is diagonalisable, the matrix power can be easily obtained. Indeed, let

$$T^{-1}AT = \Lambda = \begin{bmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{bmatrix},$$

Matrix Powers for Diagonal Matrices

- ▶ If a matrix is diagonalisable, the matrix power can be easily obtained. Indeed, let

$$T^{-1}AT = \Lambda = \begin{bmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n \end{bmatrix},$$

- ▶ We have that

$$A^k = T\Lambda^k T^{-1} = T \begin{bmatrix} \lambda_1^k & 0 & \dots & 0 \\ 0 & \lambda_2^k & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \lambda_n^k \end{bmatrix} T^{-1}.$$

Matrix Powers for Diagonal Matrices with Complex Eigenvalues

- ▶ If the matrix A has complex eigenvalues, it is possible to transform a diagonalisable matrix on the complex set into a *block diagonal matrix*, whose block dimension is at most 2.

Matrix Powers for Diagonal Matrices with Complex Eigenvalues

- ▶ If the matrix A has complex eigenvalues, it is possible to transform a diagonalisable matrix on the complex set into a *block diagonal matrix*, whose block dimension is at most 2.
- ▶ The diagonal blocks are associated to each complex and conjugated pair.

Matrix Powers for Diagonal Matrices with Complex Eigenvalues

- ▶ If the matrix A has complex eigenvalues, it is possible to transform a diagonalisable matrix on the complex set into a *block diagonal matrix*, whose block dimension is at most 2.
- ▶ The diagonal blocks are associated to each complex and conjugated pair.
- ▶ As for the continuous time case, since $AV = V\Lambda$ we have $AVH = V\Lambda H = VHH^{-1}\Lambda H$, we have

$$A[v_r + jv_c | v_r - jv_c]H = A[v_r | v_c] = [v_r | v_c] \begin{bmatrix} \sigma & \omega \\ -\omega & \sigma \end{bmatrix},$$

- ▶ The matrix can be further transformed using the modulus $\rho = \sqrt{\sigma^2 + \omega^2}$ and phase $\theta = \arctan\left(\frac{\omega}{\sigma}\right)$, i.e.,

$$A[v_r | v_c] = [v_r | v_c]\rho \begin{bmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{bmatrix}.$$

Matrix Powers for Diagonal Matrices with Complex Eigenvalues

- ...which leads us to...

$$A^k[v_r|v_c] = [v_r|v_c]\rho^k \begin{bmatrix} \cos(k\theta) & \sin(k\theta) \\ -\sin(k\theta) & \cos(k\theta) \end{bmatrix}.$$

Matrix Powers for Diagonal Matrices with Complex Eigenvalues

- ...which leads us to...

$$A^k[v_r|v_c] = [v_r|v_c]\rho^k \begin{bmatrix} \cos(k\theta) & \sin(k\theta) \\ -\sin(k\theta) & \cos(k\theta) \end{bmatrix}.$$

A 4 X 4 example

let A be a matrix defined in $\mathbb{R}^{4 \times 4}$, with λ_1 and λ_4 real and $\lambda_{2,3} = \sigma \pm j\omega$, we have

$$A^k = T\Lambda^k T^{-1} = T \begin{bmatrix} \lambda_1^k & 0 & 0 & 0 \\ 0 & \rho^k \cos(k\theta) & \rho^k \sin(k\theta) & 0 \\ 0 & -\rho^k \sin(k\theta) & \rho^k \cos(k\theta) & 0 \\ 0 & 0 & 0 & \lambda_4^k \end{bmatrix} T^{-1}.$$

Defective matrices

- ▶ We can still use the Jordan form

Defective matrices

- ▶ We can still use the Jordan form
- ▶ Indeed for a block diagonal matrix we have:

$$\begin{bmatrix} A_1 & 0 & \dots & 0 \\ 0 & A_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & A_n \end{bmatrix}^k = \begin{bmatrix} A_1^k & 0 & \dots & 0 \\ 0 & A_2^k & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & A_n^k \end{bmatrix}.$$

- ▶ Hence, the name of the game becomes to understand how to compute the power of *Jordan miniblocks*.

Defective matrices

- We can see that

$$J_{i,l}^k = \begin{bmatrix} \lambda_i & 1 & 0 & \dots & 0 & 0 \\ 0 & \lambda_i & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \lambda_i & 1 \\ 0 & 0 & 0 & \dots & 0 & \lambda_i \end{bmatrix}^l = (\Lambda_i + \bar{J}_{i,l})^k = \sum_{q=0}^k C_q^k \lambda_i^{k-q} \bar{J}_{i,l}^q,$$

where

$$\begin{cases} C_q^k = \binom{k}{q} = \frac{k!}{q!(k-q)!} & \text{if } k \geq q, \\ C_q^k = 0 & \text{if } k < q, \end{cases},$$

that is a polynomial function of k of degree q , i.e., the *binomial coefficient*.

Defective matrices

- Therefore, for a miniblock of dimension $p = m_{i,k}$, we have

$$J_{i,l}^k = \begin{bmatrix} \lambda_i^k & C_1^k \lambda_i^{k-1} & C_2^k \lambda_i^{k-2} & \dots & C_{q-2}^k \lambda_i^{k-q+2} & C_{q-1}^k \lambda_i^{k-q+1} \\ 0 & \lambda_i^k & C_1^k \lambda_i^{k-1} & \dots & C_{q-3}^k \lambda_i^{k-q+3} & C_{q-2}^k \lambda_i^{k-q+2} \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \lambda_i^k & C_1^k \lambda_i^{k-1} \\ 0 & 0 & 0 & \dots & 0 & \lambda_i^k \end{bmatrix}.$$

An example

A 3 X 3 Matrix

Let us consider the matrix

$$A = \begin{bmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 1 \\ 0 & 0 & \lambda \end{bmatrix}.$$

An example

A 3 X 3 Matrix

Let us consider the matrix

$$A = \begin{bmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 1 \\ 0 & 0 & \lambda \end{bmatrix}.$$

- By applying the formulas, it follows that

$$A^k = \begin{bmatrix} \lambda^k & k\lambda^{k-1} & \frac{k(k-1)}{2}\lambda^{k-2} \\ 0 & \lambda^k & k\lambda^{k-1} \\ 0 & 0 & \lambda^k \end{bmatrix}.$$

Matrix Powers of Defective Matrices with Complex Eigenvalues

- ▶ If we have generalised eigenvectors associated with complex eigenvalues, we can repeat the same reasoning of the continuous time case.

Matrix Powers of Defective Matrices with Complex Eigenvalues

- ▶ If we have generalised eigenvectors associated with complex eigenvalues, we can repeat the same reasoning of the continuous time case.
- ▶ For the case $p = 2$ we have

$$J_r = H^{-1} J H = \begin{bmatrix} \sigma & \omega & 1 & 0 \\ -\omega & \sigma & 0 & 1 \\ 0 & 0 & \sigma & \omega \\ 0 & 0 & -\omega & \sigma \end{bmatrix} = \begin{bmatrix} W & I \\ 0 & W \end{bmatrix}.$$

- ▶ For the matrix power, we recall that

$$J_r^k = \begin{bmatrix} W^k & kW^{k-1} \\ 0 & W^k \end{bmatrix} = \begin{bmatrix} W^k & C_1^k W^{k-1} \\ 0 & W^k \end{bmatrix},$$

Matrix Powers of Defective Matrices with Complex Eigenvalues

Which corresponds to

$$J_r^k = \begin{bmatrix} \rho^k \begin{bmatrix} \cos(k\theta) & \sin(k\theta) \\ -\sin(k\theta) & \cos(k\theta) \end{bmatrix} & 0 \\ 0 & \rho^k \begin{bmatrix} \cos(k\theta) & \sin(k\theta) \\ -\sin(k\theta) & \cos(k\theta) \end{bmatrix} \end{bmatrix} = k\rho^{k-1} \begin{bmatrix} \begin{bmatrix} \cos((k-1)\theta) & \sin((k-1)\theta) \\ -\sin((k-1)\theta) & \cos((k-1)\theta) \end{bmatrix} & 0 \\ 0 & \begin{bmatrix} \cos(k\theta) & \sin(k\theta) \\ -\sin(k\theta) & \cos(k\theta) \end{bmatrix} \end{bmatrix}.$$

Matrix Powers of Defective Matrices with Complex Eigenvalues

Which corresponds to

$$J_r^k = \begin{bmatrix} \rho^k \begin{bmatrix} \cos(k\theta) & \sin(k\theta) \\ -\sin(k\theta) & \cos(k\theta) \end{bmatrix} & 0 \\ 0 & \rho^k \begin{bmatrix} \cos(k\theta) & \sin(k\theta) \\ -\sin(k\theta) & \cos(k\theta) \end{bmatrix} \end{bmatrix} k\rho^{k-1} \begin{bmatrix} \cos((k-1)\theta) & \sin((k-1)\theta) \\ -\sin((k-1)\theta) & \cos((k-1)\theta) \end{bmatrix}.$$

We can generalise this to $p > 2$

Mode Analysis

- ▶ The unforced response of a linear system is just a linear composition of *only* the functions that can be found in the real Jordan form matrix.
- ▶ Such functions are called the *modes* of the system.

Continuous Time Systems

For CT systems, such modes are given by:

- ▶ Simple exponential functions $e^{\lambda t}$ given by Jordan miniblocks of dimension one with real eigenvalue λ
 - ▶ converge to 0 if $\lambda < 0$, constant if $\lambda = 0$, diverges if $\lambda > 0$
- ▶ Composite exponential functions $t^k e^{\lambda t}$ given by Jordan miniblocks of dimension $m > 1$ with real eigenvalue λ .
 - ▶ converge to 0 if $\lambda < 0$, diverge polynomially if $\lambda = 0$, diverge exponentially if $\lambda > 0$
- ▶ Oscillating functions of the type $e^{\sigma t} \cos(\omega t)$ and $e^{\sigma t} \sin(\omega t)$ given by two miniblocks of dimension 1 associated to complex and conjugated eigenvalues $\lambda = \sigma \pm j\omega$.
 - ▶ converge to 0 if $\sigma < 0$, persistently oscillate if $\lambda = 0$, diverge exponentially if $\sigma > 0$

Continuous Time Systems

For CT systems, such modes are given by:

- ▶ Oscillating functions of the type $t^k e^{\sigma t} \cos(\omega t)$ and $t^k e^{\sigma t} \sin(\omega t)$ given by Jordan miniblocks of dimension $m > 1$ (and hence $k = 0, \dots, m - 1$) with complex and conjugated eigenvalues $\lambda = \sigma \pm j\omega$.
 - ▶ converge $\sigma < 0$, diverge polynomially if $\sigma = 0$ and $k > 0$ or diverge exponentially if $\sigma > 0$

Discrete Time Systems

For DT systems, such modes are given by:

- ▶ Powers λ^k given by Jordan miniblocks of dimension one with real eigenvalue λ .
 - ▶ converge $|\lambda| < 1$, constant if $|\lambda| = 1$ or diverge exponentially if $|\lambda| > 1$. We have an oscillation if $\lambda < 0$

Discrete Time Systems

For DT systems, such modes are given by:

- ▶ Powers λ^k given by Jordan miniblocks of dimension one with real eigenvalue λ .
 - ▶ converge $|\lambda| < 1$, constant if $|\lambda| = 1$ or diverge exponentially if $|\lambda| > 1$. We have an oscillation if $\lambda < 0$
- ▶ Series of functions $C_q^k \lambda^{k-q}$ given by Jordan miniblocks of dimension $m > 1$ (and hence $k = 0, \dots, m-1$) with real eigenvalue λ .
 - ▶ Converge if $|\lambda| < 1$, diverge polynomially if $\lambda = 1$, diverge exponentially if $|\lambda| > 1$. We still have an oscillation if $\lambda < 0$.

Discrete Time Systems

For DT systems, such modes are given by:

- ▶ Powers λ^k given by Jordan miniblocks of dimension one with real eigenvalue λ .
 - ▶ converge $|\lambda| < 1$, constant if $|\lambda| = 1$ or diverge exponentially if $|\lambda| > 1$. We have an oscillation if $\lambda < 0$
- ▶ Series of functions $C_q^k \lambda^{k-q}$ given by Jordan miniblocks of dimension $m > 1$ (and hence $k = 0, \dots, m-1$) with real eigenvalue λ .
 - ▶ Converge if $|\lambda| < 1$, diverge polynomially if $\lambda = 1$, diverge exponentially if $|\lambda| > 1$. We still have an oscillation if $\lambda < 0$.
- ▶ Oscillating sequence of the type $\rho^k \cos(k\theta)$ and $\rho^k \sin(k\theta)$ given by two miniblocks of dimension 1 associated to complex and conjugated eigenvalues $\lambda = \sigma \pm j\omega = \rho e^{\pm j\theta}$.
 - ▶ Converge if $\rho < 1$, oscillate persistently if $\rho = 1$, diverge exponentially if $\rho > 1$

Discrete Time Systems

For DT systems, such modes are given by:

- ▶ Oscillating series of the type $C_q^k \rho^{k-q} \cos((k-q)\theta)$ and $C_q^k \rho^{k-q} \sin((k-q)\theta)$ given by Jordan miniblocks of dimension $m > 1$ (and hence $k = 0, \dots, m-1$) with complex and conjugated eigenvalues $\lambda = \sigma \pm j\omega = \rho e^{\pm j\theta}$.
 - ▶ Converge if $\rho < 1$, diverge polynomially if $\rho = 1$ and $k > 0$ or diverge exponentially if $\rho > 1$.