
CIS 471/571 (Fall 2020): Introduction to Artificial Intelligence

Assignment Project Exam Help

Lectu <https://eduassistpro.github.io/> Nets
Add WeChat edu_assist_pro

Thanh H. Nguyen

Source: <http://ai.berkeley.edu/home.html>



Reminder:

- Written assignment 3:
 - Deadline: Nov 10, 2020

Assignment Project Exam Help

- Programming proje <https://eduassistpro.github.io/>
 - Deadline: Nov 10, 2020

Add WeChat edu_assist_pro

Probabilistic Models

- Models describe how (a portion of) the world works
- Models are always simplifications
 - May not account for every variable
 - May not account for all interactions
 - “All models are wrong; but some are useful”
 - George E. P. Box
- What do we do with probabilistic models?
 - We (or our agents) need to reason about unknown variables, given evidence
 - Example: explanation (diagnostic reasoning)
 - Example: prediction (causal reasoning)

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



Probability Recap

- Conditional probability $P(x|y) = \frac{P(x, y)}{P(y)}$
- Product rule $P(x, y) = P(x|y)P(y)$

- Chain rule

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



Independence

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



Independence

- Two variables are *independent* if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

Assignment Project Exam Help

- This says that their joint distribution is the product of two simpler distributions

<https://eduassistpro.github.io/>


- Another form: $\forall x, y : P(x|y) = P(x)$

Add WeChat edu_assist_pro

- We write: $X \perp\!\!\!\perp Y$

- Independence is a simplifying *modeling assumption*

- What could we assume for {Weather, Traffic, Cavity, Toothache}?



Example: Independence?

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

$P(T)$

T	P
hot	0.5
cold	0.5

$P_1(T, W)$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

$P(W)$

W	P
sun	0.6
rain	0.4

$P_2(T, W)$

T	W	P
hot	sun	0.3
hot	rain	0.2
cold	sun	0.3
cold	rain	0.2

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



Example: Independence

- N fair, independent coin flips:

$$P(X_1)$$

H	0.5
T	0.5

$$P(X_2)$$



H	0.5
T	0.5

Assignment Project Exam Help

$$P(X_n)$$

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



Conditional Independence

- Unconditional (absolute) independence very rare
- *Conditional independence* is our most basic and robust form of knowledge about uncertain environments.

<https://eduassistpro.github.io/>

- X is conditionally independent of Y given Z

if and only if:

$$\forall x, y, z : P(x, y|z) = P(x|z)P(y|z)$$

or, equivalently, if and only if

$$\forall x, y, z : P(x|z, y) = P(x|z)$$



Conditional Independence

- $P(\text{Toothache}, \text{Cavity}, \text{Catch})$
- If I have a cavity, the probability that the probe catches in it doesn't depend on whether I have a toothache:
 - $P(+\text{catch} \mid +\text{toothache}, +\text{cavity}) = P(+\text{catch} \mid +\text{cavity})$
- The same independence holds if $P(+\text{catch} \mid +\text{toothache}, -\text{cavity}) = P(+\text{catch} \mid -\text{cavity})$
- Catch is *conditionally independent* of Toothache given Cavity:
 - $P(\text{Catch} \mid \text{Toothache}, \text{Cavity}) = P(\text{Catch} \mid \text{Cavity})$
- **Equivalent statements:**
 - $P(\text{Toothache} \mid \text{Catch}, \text{Cavity}) = P(\text{Toothache} \mid \text{Cavity})$
 - $P(\text{Toothache}, \text{Catch} \mid \text{Cavity}) = P(\text{Toothache} \mid \text{Cavity}) P(\text{Catch} \mid \text{Cavity})$
 - One can be derived from the other easily

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



Conditional Independence

- What about this domain:

- Traffic
- Umbrella
- Raining

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



Conditional Independence


- What about this domain:

- Fire
- Smoke
- Alarm

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



Conditional Independence and the Chain Rule

- Chain rule: $P(X_1, X_2, \dots, X_n) = P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots$

- Trivial decomposition: **Assignment Project Exam Help**


<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

- With assumption of conditional independence

$$P(\text{Traffic}, \text{Rain}, \text{Umbrella}) = \\ P(\text{Rain})P(\text{Traffic}|\text{Rain})P(\text{Umbrella}|\text{Rain})$$

- Bayes'nets / graphical models help us express conditional independence assumptions



Ghostbusters Chain Rule

- Each sensor depends only on where the ghost is

$$P(T,B,G) = P(G) P(T|G) P(B|G)$$

- That means, the two sensors are conditionally independent, given the ghost position

- T: Top square is red
- B: Bottom square is red
- G: Ghost is in the top

- Givens:

$$P(+g) = 0.5$$


$$P(-g) = 0.5$$

$$P(+t | +g) = 0.8$$

$$P(+t | -g) = 0.4$$

$$P(+b | +g) = 0.4$$

$$P(+b | -g) = 0.8$$



T	B	G	P(T,B,G)
			0.16
			0.16
			0.24
+t	-b	-g	0.04
-t	+b	+g	0.04
-t	+b	-g	0.24
-t	-b	+g	0.06
-t	-b	-g	0.06



Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

Bayes' Nets: Big Picture

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro




Bayes' Nets: Big Picture

- Two problems with using full joint distribution tables as our probabilistic models:
 - Unless there are only a few variables, the joint is WAY too big to represent explicitly
 - Hard to learn (estimate) anything empirically about more than a few variables at a time

Assignment Project Exam Help

<https://eduassistpro.github.io/>

- Bayes' nets:** a technique for describing complex distributions (models) using simple, local distributions (conditional probabilities)
 - More properly called **graphical models**
 - We describe how variables locally interact
 - Local interactions chain together to give global, indirect interactions
 - For about 10 min, we'll be vague about how these interactions are specified



Example Bayes' Net: Insurance

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



Example Bayes' Net: Car

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



Graphical Model Notation

- Nodes: variables (with domains)
 - Can be assigned (observed) or unassigned (unobserved)
- Arcs: interactions
 - Similar to CSP constraints
 - Indicate “direct influence” between variables
 - Formally: encode conditional independence (more later)

- For now: imagine that arrows mean direct causation (in general, they don't!)

Assignment Project Exam Help



<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



Example: Coin Flips


- N independent coin flips



Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro





- No interactions between variables: **absolute independence**



Example: Traffic

- Variables:
 - R: It rains
 - T: There is traffic

- Model 1: independence




- Why is an agent using model 2 better?

Assignment Project Exam Help

<https://eduassistpro.github.io/causes/traffic>

Add WeChat edu_assist_pro



Example: Traffic II

- Let's build a causal graphical model!

- Variables

- T: Traffic
- R: It rains
- L: Low pressure
- D: Roof drips
- B: Ballgame
- C: Cavity

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



Example: Alarm Network

- Variables
 - B: Burglary
 - A: Alarm goes off
 - M: Mary calls
 - J: John calls
 - E: Earthquake!

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



Bayes' Net Semantics

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro





Bayes' Net Semantics

- A set of nodes, one per variable X

- A directed, acyclic graph

- A conditional distribution

- A collection of distributions over X , one for each combination of parents' values


$$P(X|a_1 \dots a_n)$$

- CPT: conditional probability table
 - Description of a noisy “causal” process

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



A Bayes net = Topology (graph) + Local Conditional Probabilities





Probabilities in BNs


- Bayes' nets **implicitly** encode joint distributions
 - As a product of local conditional distributions
 - To see what probability a BN gives to a full assignment, multiply all the

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

- Example:



$$P(+cavity, +catch, -toothache)$$





Probabilities in BNs

- Why are we guaranteed that setting

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$

results in a proper joint distribution?

Assignment Project Exam Help

- Chain rule (valid for all distrib <https://eduassistpro.github.io/>



- Assume conditional independences: Add WeChat edu_assist_pro

→ Consequence:
$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$


- Not every BN can represent every joint distribution
 - The topology enforces certain conditional independencies



Example: Coin Flips



...



$P(X_1)$


h	0.5
t	0.5

$P(X_2)$

h	0.5
t	0.5

<https://eduassistpro.github.io/>

Add WeChat [edu_assist_pro](#)




$$P(h, h, t, h) =$$

Only distributions whose variables are absolutely independent can be represented by a Bayes' net with no arcs.



Example: Traffic


$$P(R)$$

+r	1/4
-r	3/4

$$P(T|R)$$

+r	<table><tr><td>+t</td><td>3/4</td></tr><tr><td>-t</td><td>1/4</td></tr></table>	+t	3/4	-t	1/4
+t	3/4				
-t	1/4				
-r	<table><tr><td>+t</td><td>1/2</td></tr><tr><td>-t</td><td>1/2</td></tr></table>	+t	1/2	-t	1/2
+t	1/2				
-t	1/2				

$$P(+r, -t) =$$

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro




Example: Alarm Network

B	P(B)
+b	0.001
-b	0.999




E	P(E)
+e	0.002
-e	0.998



Assignment Project Exam Help



<https://eduassistpro.github.io/>



Add WeChat edu_assist_pro

A	J	P(J A)
+a	+j	0.9
+a	-j	0.1
-a	+j	0.05
-a	-j	0.95


A	M	P(M A)
+a	+m	0.7
+a	-m	0.3
-a	+m	0.01
-a	-m	0.99

	E	A	P(A B,E)
	+e	+a	0.95
+b	+e	-a	0.05
+b	-e	+a	0.94
+b	-e	-a	0.06
-b	+e	+a	0.29
-b	+e	-a	0.71
-b	-e	+a	0.001
-b	-e	-a	0.999



Example: Alarm Network


B	P(B)
+b	0.001
-b	0.999



E	P(E)
+e	0.002
-e	0.998

A	J	P(J A)
+a	+j	0.9
+a	-j	0.1
-a	+j	0.05
-a	-j	0.95

		P(M A)
		.7
		.3
-a	+m	0.01
-a	-m	0.99



B	E	A	P(A B,E)
+b	+e	+a	0.95
+b	+e	-a	0.05
+b	-e	+a	0.94
+b	-e	-a	0.06
-b	+e	+a	0.29
-b	+e	-a	0.71
-b	-e	+a	0.001
-b	-e	-a	0.999

$$P(+b, -e, +a, -j, +m) =$$


Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

Example: Traffic

- Causal direction



$P(R)$

+r	1/4
-r	3/4

$P(T|R)$

+r	+t	3/4
	-t	1/4
-r	+t	1/2
	-t	1/2

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

+r	+t	3/16
+r	-t	1/16
-r	+t	6/16
-r	-t	6/16




Example: Reverse Traffic

- Reverse causality?

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



$P(T)$

+t	9/16
-t	7/16

$P(R|T)$

+t	+r	1/3
	-r	2/3

-t	+r	1/7
	-r	6/7

+r	+t	3/16
+r	-t	1/16
-r	+t	6/16
-r	-t	6/16



Size of a Bayes' Net

- How big is a joint distribution over N Boolean variables?

$$2^N$$

- Both give you the power to calculate

$$P(X_1, X_2, \dots, X_n)$$

Assignment Project Exam Help
ENX Huge space savings!

- How big is an N-node net if have up to k parents?

$$O(N * 2^{k+1})$$

<https://eduassistpro.github.io/> to elicit local CPTs

Add WeChat [edu_assist_pro](https://eduassistpro.github.io/) to answer queries (coming)



Causality?

- When Bayes' nets reflect the true causal patterns:

- Often simpler (nodes have fewer parents)
- Often easier to think about
- Often easier to elicit from experts

Assignment Project Exam Help

- BNs need not actually be causal <https://eduassistpro.github.io/>

- Sometimes no causal net exists over the domain (espec variables are missing)
- E.g. consider the variables *Traffic* and *Drips*
- End up with arrows that reflect correlation, not causation

Add WeChat edu_assist_pro

- What do the arrows really mean?

- Topology may happen to encode causal structure
- **Topology really encodes conditional independence**

$$P(x_i | x_1, \dots, x_{i-1}) = P(x_i | \text{parents}(X_i))$$



Bayes' Nets

- So far: how a Bayes' net encodes a joint distribution
- Next: how to answer queries about that distribution
 - Today:
 - First assembled BNs using an intuitive notion of conditional independence as causality
 - Then saw that key property is conditional independence
 - Main goal: answer queries about conditional independence and influence
- After that: how to answer numerical queries (inference)

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

