

COMP 250

Assignment Project Exam Help


INTRODUC TER SCIENCE

<https://eduassistpro.github.io/>

Week 9-1 : I
Add WeChat edu_assist_pro

Giulia Alberini, Fall 2020

WHAT ARE WE GOING TO DO IN THIS VIDEO?



- Inductive/Recursive definitions
 - Inductive/Recursive definitions
 - Mathematical Induction
- <https://eduassistpro.github.io/>
Add WeChat edu_assist_pro

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

PROOFS

For all $n \geq 1$,

$$1 + 2 + 3 + \dots + (n-1) + n = \frac{n(n+1)}{2}$$

<https://eduassistpro.github.io/>

How can we prove such a statement?

Add WeChat edu_assist_pro

- By “proof”, we mean a formal logical argument that convincingly demonstrate the truth of a given proposition.
- Note that “convincingly” is itself not well defined.

EXAMPLE

$$1 + 2 + \dots + (n - 1) + n$$

Rewrite by considering $n/2$ pairs :

Assignment Project Exam Help

$$1 + 2 + \dots + \frac{n}{2} + \left(\frac{n}{2} + 1 \right) + \dots + (n - 1) + n$$

Add WeChat edu_assist_pro

If n is even, then adding up the $n/2$ pairs gives

$$n/2 * (n + 1)$$

- What if n is odd?

EXAMPLE

- What if n is odd? Then, $n-1$ is even. So,

Assignment Project Exam Help
 $1 + 2 + \dots + (n-1) + n$

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro
 $= \left(\frac{n-1}{2} + 1 \right)$

$$= \frac{n+1}{2} * n$$

which is the same formula as before.

RECURSIVE (INDUCTIVE) DEFINITION

- Some set of elements can be define recursively/inductively.

- A recursive/inductive definition consists of the following:

- A *base clause*

Which one or more basic/initial eleme

- One or more *inductive clauses*

Rules on how to generate “new” elements of the set from “old” ones.

- A *final clause*

which simply states that no other element is part of the set.

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

EXAMPLE – NATURAL NUMBERS

The set of natural numbers can be defined as follows:

- *Base clause:*

0 is a natural number

Assignment Project Exam Help

<https://eduassistpro.github.io/>

- *Inductive clause:*

If n is a natural number, then $n + 1$ is also

Add WeChat edu_assist_pro
mber.

- *Final clause:* Nothing else is a natural number.

MATHEMATICAL INDUCTION

Consider a statement of the form:

“For all $n \geq n_0$, $P(n)$ is true”

where n_0 is some constant and $P(n)$ has value true or false for each n .

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

- If n is an element of an inductively defined set, then the statement above can be proven using a technique called *mathematical induction*.

(WEAK) MATHEMATICAL INDUCTION

To prove a property by mathematical induction, we proceed as follows:

- *Base case*

Show that the property holds for the basic/initial elements of the set.

Assignment Project Exam Help

<https://eduassistpro.github.io/>

- *Induction step*

Assume the property hold for some element n (Induction Hypothesis)

Show that the property also holds for any element generated from n using the inductive clauses.

Add WeChat edu_assist_pro

- *Conclusion*

The property holds for all elements.

EXAMPLE

“For all $n \geq n_0$, $P(n)$ is true”

Assignment Project Exam Help

For all $n \geq 1$,

$$1 + 2 + 3 + \dots + (n - 1) + n = \frac{n(n+1)}{2}$$

https://eduassistpro.github.io/
Add WeChat edu_assist_pro

This is a property of natural numbers. Since this is a set that can be defined inductively, we can use mathematical induction to prove such property!

PROOF BY MATHEMATICAL INDUCTION

We need to prove the following:

- Base case:

$P(n_0)$ is true, i.e. the property holds for the first element n_0 .
this case is 1.

Assignment Project Exam Help

<https://eduassistpro.github.io/>

- Induction step:

IH: Assume $P(k)$ is true, i.e. the property holds for element k .

Add WeChat edu_assist_pro


Prove that $P(k + 1)$ is true, i.e. the property holds for $k + 1$.

Base case:

$P(n_0)$ is true.

Induction step:

For any $k \geq n_0$, if $P(k)$ is true
then $P(k+1)$ is true.




<https://eduassistpro.github.io/>

Thus we have proved:

Add WeChat edu_assist_pro

For any $n \geq n_0$, $P(n)$ is true.



BACK TO THE PROOF

For all $n \geq 1$, $1 + 2 + 3 + \dots + (n - 1) + n = \frac{n(n+1)}{2}$

Assignment Project Exam Help

- Base case: $n = 1$, to prov

<https://eduassistpro.github.io/>

Add WeChat $\frac{1 * (1 + 1)}{2}$ edu_assist_pro

$$1 = \frac{2}{2} = 1$$



Statement: For all $n \geq 1$, $1 + 2 + 3 + \dots + (n - 1) + n = \frac{n(n+1)}{2}$

BACK TO THE PROOF

- Induction step:

IH: Assume that it holds for k , that is

Assignment Project Exam Help
 $k(k + 1)$

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

Statement: For all $n \geq 1$, $1 + 2 + 3 + \dots + (n - 1) + n = \frac{n(n+1)}{2}$

BACK TO THE PROOF

- Induction step:

IH: Assume that it holds for k , that is

Assignment Project Exam Help
 $k(k+1)$

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro
 $1 + 2 + \dots + k$

Prove it for $k + 1$:

$$= \frac{k(k+1)}{2} + (k + 1), \text{ by IH}$$

$$= (k + 1) * \left(\frac{k}{2} + 1\right) = \frac{(k+1)(k+2)}{2}$$



EXAMPLE 2


- Prove the following statement:

For all $n \geq 3$, $2n + 1 < 2^n$.

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



EXAMPLE 2

Statement: For all $n \geq 3$, $2n + 1 < 2^n$.

Assignment Project Exam Help

- Note: $P(n)$ is false for
But that has nothing to do

<https://eduassistpro.github.io/>
e.

Add WeChat edu_assist_pro

EXAMPLE 2

Statement: For all $n \geq 3$, $2n + 1 < 2^n$.

Assignment Project Exam Help

Proof: (by mathematical in

■ Base case ($n = 3$):

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

$$2 * 3 + 1 = 7$$



Statement: For all $n \geq 3$, $2n + 1 < 2^n$.

EXAMPLE 2

- Induction step:

IH: Assume $2 * k + 1 < 2^k$.

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

Statement: For all $n \geq 3$, $2n + 1 < 2^n$.

EXAMPLE 2

■ Induction step:

IH: Assume $2 * k + 1 < 2^k$.

Prove it for $k + 1$:

Assignment Project Exam Help

<https://eduassistpro.github.io/>

()

Add WeChat edu_assist_pro

$= 2 * k +$

Statement: For all $n \geq 3$, $2n + 1 < 2^n$.

EXAMPLE 2

■ Induction step:

IH: Assume $2 * k + 1 < 2^k$.

Prove it for $k + 1$:

<https://eduassistpro.github.io/>
Add WeChat edu_assist_pro

$< 2^k + 2$, by IH

$< 2^k + 2^k$, for $k \geq 3$

$= 2^{k+1}$




EXAMPLE 3

Statement: For all $n \geq 5$, $n^2 < 2^n$.

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



EXAMPLE 3

Statement: For all $n \geq 5$, $n^2 < 2^n$.

Assignment Project Exam Help

Proof: (by mathematical in

- Base case ($n = 5$):

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



$$5^2 = 25 <$$

EXAMPLE 3

Statement: For all $n \geq 5$, $n^2 < 2^n$.

- Induction step.

Assignment Project Exam Help

What should we assume? <https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

What do we need to prove?

EXAMPLE 3

Statement: For all $n \geq 5$, $n^2 < 2^n$.

■ Induction step.

Assignment Project Exam Help

What should we assume? <https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

What do we need to prove? $(k + 1)^2 < 2^{(k+1)}$

EXAMPLE 3

Statement: For all $n \geq 5$, $n^2 < 2^n$.

- Induction step.

IH: $k^2 < 2^k$ for a $k \geq 5$

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

EXAMPLE 3

Statement: For all $n \geq 5$, $n^2 < 2^n$.

- Induction step.

IH: $k^2 < 2^k$ for a $k \geq 5$

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

$< 2^k + 2^k$, by Example 2

$= 2^{k+1}$



(STRONG) MATHEMATICAL INDUCTION

- Sometimes one would like to assume the induction hypothesis not only for the previous element, but also for smaller elements. This leads to a logically equivalent proof method called *strong (or complete) mathematical induction*.
- To prove a property by strong induction, proceed as follows:
 - Induction step
Assume the property holds for all elements less than arbitrary k . (Induction Hypothesis)
Show that the property also holds for the k element which was generated using the inductive clauses.
 - Conclusion
The property holds for all elements.

FIBONACCI NUMBERS

- The Fibonacci sequence is one of the most common example of a recursively-defined set.

Assignment Project Exam Help

- Consider the following s <https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144,

Let f_n denote the n th Fibonacci number. How can we define the sequence above?

FIBONACCI NUMBERS – INDUCTIVE DEFINITION

- Fibonacci sequence: 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, ...

Assignment Project Exam Help

- Base clause:

$f_0 = f_1 = 1$ are Fibonacci <https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

- Inductive clause:

If f_{n-1} and f_{n-2} are Fibonacci numbers, then $f_n = f_{n-1} + f_{n-2}$ is a Fibonacci number.

EXAMPLE 4

Statement: For all $n \geq 0$, $f_n \leq \left(\frac{7}{4}\right)^n$

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

EXAMPLE 4

Statement: For all $n \geq 0$, $f_n \leq \left(\frac{7}{4}\right)^n$

Proof: (by strong mathematical induction)

■ Induction step

IH: Let $k \geq 0$, and assume that for any i such that $0 \leq i < k$ then

$$f_i \leq \left(\frac{7}{4}\right)^i$$

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

EXAMPLE 4

Statement: For all $n \geq 0$, $f_n \leq \left(\frac{7}{4}\right)^n$

Proof: (by strong mathematical induction)

■ Induction step

IH: Let $k \geq 0$, and assume that for any n such that $0 \leq n < k$ then

$$f_n \leq \left(\frac{7}{4}\right)^n$$

To show: $f_k \leq \left(\frac{7}{4}\right)^k$

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

EXAMPLE 4

There are 3 possible cases:

1. $k = 0$

$f_0 = 1$ and $\left(\frac{7}{4}\right)^0 = 1$, so t

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

2. $k = 1$

$f_1 = 1$ and $\left(\frac{7}{4}\right)^1 > 1$, so the claim holds.

EXAMPLE 4

There are 3 possible cases:

3. $k > 1$

$$\begin{aligned} f_k &= f_{k-1} + f_{k-2} \\ &= \binom{-}{-}^{k-1} \binom{-}{-}^{k-2}, \text{ by IH} \\ &= \left(\frac{7}{4}\right)^k \binom{-}{-}^{k-2} \binom{7}{4}^{k-2} \left(\frac{11}{4}\right) \\ &= \left(\frac{7}{4}\right)^{k-2} \left(\frac{44}{16}\right) \\ &< \left(\frac{7}{4}\right)^{k-2} \left(\frac{49}{16}\right) = \left(\frac{7}{4}\right)^{k-2} \left(\frac{7}{4}\right)^2 \\ &= \left(\frac{7}{4}\right)^k \end{aligned}$$

RECOMMENDED EXERCISES

1. Prove that for all $n \geq 0$, $\sum_{i=0}^n 2^i = 2^{n+1} - 1$

2. Prove that for all $n \geq 0$, $\sum_{i=0}^n i^3 = \left(\frac{n(n+1)}{2}\right)^2$

3. Consider the following re

n ('+') on natural numbers:

■ Base clause:

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

■ Inductive clause:

$$(n + 1) + m = (n + m) + 1$$

Prove that addition is associative, i.e. for all natural numbers $(a + b) + c = a + (b + c)$

Hint: use mathematical induction on a



Coming Soon

Assignment Project Exam Help

In the next

■ Recursive <https://eduassistpro.github.io/>

Add WeChat edu_assist_pro