

COMP 250

Assignment Project Exam Help


INTRODUC TER SCIENCE

<https://eduassistpro.github.io/>

Week 9-1 : I
Add WeChat edu_assist_pro

Giulia Alberini, Fall 2020

WHAT ARE WE GOING TO DO IN THIS VIDEO?



- Inductive/Recursive definitions
 - Inductive/Recursive definitions
 - Mathematical Induction
- <https://eduassistpro.github.io/>
Add WeChat edu_assist_pro

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

PROOFS

For all $n \geq 1$,

$$1 + 2 + 3 + \dots + (n-1) + n = \frac{n(n+1)}{2}$$

<https://eduassistpro.github.io/>

How can we prove such a statement?

Add WeChat edu_assist_pro

- By “proof”, we mean a formal logical argument that convincingly demonstrate the truth of a given proposition.
- Note that “convincingly” is itself not well defined.

EXAMPLE

$$1 + 2 + \dots + (n - 1) + n$$

Rewrite by considering $n/2$ pairs :

Assignment Project Exam Help

$$1 + 2 + \dots + \frac{n}{2} + \left(\frac{n}{2} + 1 \right) + \dots + (n - 1) + n$$

Add WeChat edu_assist_pro

If n is even, then adding up the $n/2$ pairs gives

$$n/2 * (n + 1)$$

- What if n is odd?

EXAMPLE

- What if n is odd? Then, $n-1$ is even. So,

Assignment Project Exam Help
 $1 + 2 + \dots + (n-1) + n$

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro
 $= \left(\frac{n-1}{2} + 1 \right)$

$$= \frac{n+1}{2} * n$$

which is the same formula as before.

RECURSIVE (INDUCTIVE) DEFINITION

- Some set of elements can be define recursively/inductively.

Assignment Project Exam Help

- A recursive/inductive definition consists of the following:

- A *base clause*

<https://eduassistpro.github.io/>

Which one or more basic/initial eleme

Add WeChat edu_assist_pro

- One or more *inductive clauses*

Rules on how to generate “new” elements of the set from “old” ones.

- A *final clause*

which simply states that no other element is part of the set.

EXAMPLE – NATURAL NUMBERS

The set of natural numbers can be defined as follows:

- *Base clause:*

0 is a natural number

Assignment Project Exam Help

<https://eduassistpro.github.io/>

- *Inductive clause:*

If n is a natural number, then $n + 1$ is also

Add WeChat edu_assist_pro
mber.

- *Final clause:* Nothing else is a natural number.

MATHEMATICAL INDUCTION

Consider a statement of the form:

“For all $n \geq n_0$, $P(n)$ is true”

where n_0 is some constant and $P(n)$ has value true or false for each n .

<https://eduassistpro.github.io/>
Add WeChat edu_assist_pro

- If n is an element of an inductively defined set, then the statement above can be proven using a technique called *mathematical induction*.

(WEAK) MATHEMATICAL INDUCTION

To prove a property by mathematical induction, we proceed as follows:

- *Base case*

Show that the property holds for the basic/initial elements of the set.

Assignment Project Exam Help

<https://eduassistpro.github.io/>

- *Induction step*

Assume the property hold for some element n (Induction Hypothesis)

Show that the property also holds for any element generated from n using the inductive clauses.

Add WeChat edu_assist_pro

- *Conclusion*

The property holds for all elements.

EXAMPLE

“For all $n \geq n_0$, $P(n)$ is true”

Assignment Project Exam Help

For all $n \geq 1$,

$$1 + 2 + 3 + \dots + (n - 1) + n = \frac{n(n+1)}{2}$$

https://eduassistpro.github.io/
Add WeChat edu_assist_pro

This is a property of natural numbers. Since this is a set that can be defined inductively, we can use mathematical induction to prove such property!

PROOF BY MATHEMATICAL INDUCTION

We need to prove the following:

- Base case:

$P(n_0)$ is true, i.e. the property holds for the first element n_0 .
this case is 1.

Assignment Project Exam Help

<https://eduassistpro.github.io/>

- Induction step:

IH: Assume $P(k)$ is true, i.e. the property holds for element k .

Add WeChat edu_assist_pro


Prove that $P(k + 1)$ is true, i.e. the property holds for $k + 1$.

Base case:

$P(n_0)$ is true.

Induction step:

For any $k \geq n_0$, if $P(k)$ is true
then $P(k+1)$ is true.




<https://eduassistpro.github.io/>

Thus we have proved:

Add WeChat edu_assist_pro

For any $n \geq n_0$, $P(n)$ is true.



BACK TO THE PROOF

For all $n \geq 1$, $1 + 2 + 3 + \dots + (n - 1) + n = \frac{n(n+1)}{2}$

Assignment Project Exam Help

- Base case: $n = 1$, to prov

<https://eduassistpro.github.io/>

Add WeChat $\frac{1 * (1 + 1)}{2}$ edu_assist_pro

$$1 = \frac{2}{2} = 1$$



Statement: For all $n \geq 1$, $1 + 2 + 3 + \dots + (n - 1) + n = \frac{n(n+1)}{2}$

BACK TO THE PROOF

- Induction step:

IH: Assume that it holds for k , that is

Assignment Project Exam Help
 $k(k + 1)$

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

Statement: For all $n \geq 1$, $1 + 2 + 3 + \dots + (n - 1) + n = \frac{n(n+1)}{2}$

BACK TO THE PROOF

- Induction step:

IH: Assume that it holds for k , that is

Assignment Project Exam Help
 $k(k+1)$

<https://eduassistpro.github.io/>

Prove it for $k + 1$:

Add WeChat edu_assist_pro
 $1 + 2 + \dots + k$

$$= \frac{k(k+1)}{2} + (k + 1), \text{ by IH}$$

$$= (k + 1) * \left(\frac{k}{2} + 1\right) = \frac{(k+1)(k+2)}{2}$$



EXAMPLE 2


- Prove the following statement:

For all $n \geq 3$, $2n + 1 < 2^n$.

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



EXAMPLE 2

Statement: For all $n \geq 3$, $2n + 1 < 2^n$.

Assignment Project Exam Help

- Note: $P(n)$ is false for
But that has nothing to do

<https://eduassistpro.github.io/>
e.

Add WeChat edu_assist_pro

EXAMPLE 2

Statement: For all $n \geq 3$, $2n + 1 < 2^n$.

Assignment Project Exam Help

Proof: (by mathematical in

■ Base case ($n = 3$):

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

$$2 * 3 + 1 = 7$$



Statement: For all $n \geq 3$, $2n + 1 < 2^n$.

EXAMPLE 2

- Induction step:

IH: Assume $2 * k + 1 < 2^k$.

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

Statement: For all $n \geq 3$, $2n + 1 < 2^n$.

EXAMPLE 2

■ Induction step:

IH: Assume $2 * k + 1 < 2^k$.

Prove it for $k + 1$:

Assignment Project Exam Help

<https://eduassistpro.github.io/>

()

Add WeChat edu_assist_pro

$= 2 * k +$

Statement: For all $n \geq 3$, $2n + 1 < 2^n$.

EXAMPLE 2

■ Induction step:

IH: Assume $2 * k + 1 < 2^k$.

Prove it for $k + 1$:

<https://eduassistpro.github.io/>
Add WeChat edu_assist_pro

$< 2^k + 2$, by IH

$< 2^k + 2^k$, for $k \geq 3$

$= 2^{k+1}$




EXAMPLE 3

Statement: For all $n \geq 5$, $n^2 < 2^n$.

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



EXAMPLE 3

Statement: For all $n \geq 5$, $n^2 < 2^n$.

Assignment Project Exam Help

Proof: (by mathematical in

- Base case ($n = 5$):

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro



$$5^2 = 25 <$$

EXAMPLE 3

Statement: For all $n \geq 5$, $n^2 < 2^n$.

- Induction step.

Assignment Project Exam Help

What should we assume? <https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

What do we need to prove?

EXAMPLE 3

Statement: For all $n \geq 5$, $n^2 < 2^n$.

■ Induction step.

Assignment Project Exam Help

What should we assume? <https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

What do we need to prove? $(k + 1)^2 < 2^{(k+1)}$

EXAMPLE 3

Statement: For all $n \geq 5$, $n^2 < 2^n$.

- Induction step.

IH: $k^2 < 2^k$ for a $k \geq 5$

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

EXAMPLE 3

Statement: For all $n \geq 5$, $n^2 < 2^n$.

■ Induction step.

IH: $k^2 < 2^k$ for a $k \geq 5$

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

$< 2^k + 2^k$, by Example 2

$= 2^{k+1}$



(STRONG) MATHEMATICAL INDUCTION

- Sometimes one would like to assume the induction hypothesis not only for the previous element, but also for smaller elements. This leads to a logically equivalent proof method called *strong (or complete) mathematical induction*.
- To prove a property by strong induction, proceed as follows:
 - Induction step
Assume the property holds for all elements less than arbitrary k . (Induction Hypothesis)
Show that the property also holds for the k element which was generated using the inductive clauses.
 - Conclusion
The property holds for all elements.

FIBONACCI NUMBERS

- The Fibonacci sequence is one of the most common example of a recursively-defined set.

Assignment Project Exam Help

- Consider the following s <https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144,

Let f_n denote the n th Fibonacci number. How can we define the sequence above?

FIBONACCI NUMBERS – INDUCTIVE DEFINITION

- Fibonacci sequence: 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, ...

Assignment Project Exam Help

- Base clause:

$f_0 = f_1 = 1$ are Fibonacci <https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

- Inductive clause:

If f_{n-1} and f_{n-2} are Fibonacci numbers, then $f_n = f_{n-1} + f_{n-2}$ is a Fibonacci number.

EXAMPLE 4

Statement: For all $n \geq 0$, $f_n \leq \left(\frac{7}{4}\right)^n$

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

EXAMPLE 4

Statement: For all $n \geq 0$, $f_n \leq \left(\frac{7}{4}\right)^n$

Proof: (by strong mathematical induction)

■ Induction step

IH: Let $k \geq 0$, and assume that for any i such that $0 \leq i < k$ then

$$f_i \leq \left(\frac{7}{4}\right)^i$$

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

EXAMPLE 4

Statement: For all $n \geq 0$, $f_n \leq \left(\frac{7}{4}\right)^n$

Proof: (by strong mathematical induction)

■ Induction step

IH: Let $k \geq 0$, and assume that for any n such that $0 \leq n < k$ then

$$f_n \leq \left(\frac{7}{4}\right)^n$$

To show: $f_k \leq \left(\frac{7}{4}\right)^k$

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

EXAMPLE 4

There are 3 possible cases:

1. $k = 0$

$f_0 = 1$ and $\left(\frac{7}{4}\right)^0 = 1$, so t

Assignment Project Exam Help

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

2. $k = 1$

$f_1 = 1$ and $\left(\frac{7}{4}\right)^1 > 1$, so the claim holds.

EXAMPLE 4

There are 3 possible cases:

3. $k > 1$

$f_k = f_{k-1} + f_{k-2}$
Assignment Project Exam Help
 $\binom{-}{-}^{k-1} \binom{-}{-}^{k-2}$, by IH
<https://eduassistpro.github.io/>
Add WeChat edu_assist_pro
 $= \left(\frac{7}{4}\right)^k \binom{-}{-}^{k-2} \binom{11}{4}$
 $= \left(\frac{7}{4}\right)^{k-2} \left(\frac{44}{16}\right)$
 $< \left(\frac{7}{4}\right)^{k-2} \left(\frac{49}{16}\right) = \left(\frac{7}{4}\right)^{k-2} \left(\frac{7}{4}\right)^2$
 $= \left(\frac{7}{4}\right)^k$

RECOMMENDED EXERCISES

1. Prove that for all $n \geq 0$, $\sum_{i=0}^n 2^i = 2^{n+1} - 1$

2. Prove that for all $n \geq 0$, $\sum_{i=0}^n i^3 = \left(\frac{n(n+1)}{2}\right)^2$

3. Consider the following re

n ('+') on natural numbers:

- Base clause:

<https://eduassistpro.github.io/>

Add WeChat edu_assist_pro

- Inductive clause:

$$(n + 1) + m = (n + m) + 1$$

Prove that addition is associative, i.e. for all natural numbers $(a + b) + c = a + (b + c)$

Hint: use mathematical induction on a

An orange paint roller with a red handle, positioned horizontally. The roller is partially filled with orange paint, and there are orange paint splatters and drips around it. The background features faint, concentric circular lines.

Coming Soon

Assignment Project Exam Help

In the next

■ Recursive <https://eduassistpro.github.io/>

Add WeChat edu_assist_pro