

COMP2610 / COMP6261 - Information Theory

Lecture 4: Bayesian Inference

Assignment Project Exam Help

<https://eduassistpro.github.io>



Australian
National
University

Add WeChat edu_assist_pro

July 31, 2018

Assignment Project Exam Help

- Examples of joint, marginal and conditional distributions

- Wh <https://eduassistpro.github.io>

- What, if anything, does $p(X = x|Y = y)$ tell you about $p(Y = y|X = x)$?
Add WeChat: edu_assist_pro

Review Exercise

Suppose we have binary random variables X, Y such that

$$p(X = 1) = 0.6$$

$$p(Y = 1 | X = 0) = 0.7$$

$$p(Y = 1 | X = 1) = 0.8$$

Assignment Project Exam Help

Then,

<https://eduassistpro.github.io>

$$p(X = 1 | Y = 1) =$$

Add WeChat edu_assist_pr

Review Exercise

Suppose we have binary random variables X, Y such that

$$p(X = 1) = 0.6$$

$$p(Y = 1|X = 0) = 0.7$$

$$p(Y = 1|X = 1) = 0.8$$

Assignment Project Exam Help

Then,

<https://eduassistpro.github.io>

$$p(X = 1|Y = 1) = \frac{p(Y = 1|X = 1)p(X = 1)}{p(Y = 1)}$$

Add WeChat edu_assist_pro

Review Exercise

Suppose we have binary random variables X, Y such that

$$p(X = 1) = 0.6$$

$$p(Y = 1|X = 0) = 0.7$$

$$p(Y = 1|X = 1) = 0.8$$

Assignment Project Exam Help

Then,

<https://eduassistpro.github.io>

$$p(X = 1|Y = 1) = \frac{p(Y = 1|X = 1)p(X = 1)}{p(Y = 1)}$$

Add WeChat edu_assist_pro

$$= \frac{p(Y = 1|X = 1)p(X = 1) + (p(Y = 1|X = 0)p(X = 0))}{p(Y = 1)}$$

Review Exercise

Suppose we have binary random variables X, Y such that

$$p(X = 1) = 0.6$$

$$p(Y = 1|X = 0) = 0.7$$

$$p(Y = 1|X = 1) = 0.8$$

Assignment Project Exam Help

Then,

<https://eduassistpro.github.io>

$$p(X = 1|Y = 1) = \frac{p(Y = 1|X = 1)p(X = 1)}{p(Y = 1)}$$

Add WeChat edu_assist_pro

$$= \frac{p(Y = 1|X = 1)p(X = 1)}{p(Y = 1|X = 1)p(X = 1) + (p(Y = 1|X = 0)p(X = 0))}$$

$$= \frac{(0.8)(0.6)}{(0.8)(0.6) + (0.7)(0.4)}$$

$$\approx 0.63$$

This time

Assignment Project Exam Help

- More examples on Bayes' theorem:
 - ▶ Eating hamburgers

▶ <https://eduassistpro.github.io>

▶ Add WeChat edu_assist_pro

- Are there notions of probability beyond frequ

Outline

1 Bayes' Rule: Examples

- Eating Hamburgers
- Detecting Terrorists
- Th

2 Mom

3 The meaning of Probability

4 Wrapping Up

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

1 Bayes' Rule: Examples

- Eating Hamburgers
- Detecting Terrorists
- The Monty Hall Problem

Assignment Project Exam Help

2 Mom

<https://eduassistpro.github.io>

3 The meaning of Probability

Add WeChat edu_assist_pr

4 Wrapping Up

Bayesian Inference:

Example 1 (Barber, BRML, 2011)

Assignment Project Exam Help

- 90% of people with McD syndrome are frequent hamburger eaters

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Bayesian Inference:

Example 1 (Barber, BRML, 2011)

Assignment Project Exam Help

- 90% of people with McD syndrome are frequent hamburger eaters

- Pro <https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Bayesian Inference:

Example 1 (Barber, BRML, 2011)

Assignment Project Exam Help

- 90% of people with McD syndrome are frequent hamburger eaters

- Pro <https://eduassistpro.github.io>

- Add WeChat edu_assist_pr
Proportion of hamburger eaters is about 50

Bayesian Inference:

Example 1 (Barber, BRML, 2011)

Assignment Project Exam Help

- 90% of people with McD syndrome are frequent hamburger eaters

- Pro <https://eduassistpro.github.io>

- Add WeChat edu_assist_pr
Proportion of hamburger eaters is about 50

What is the probability that a hamburger eater will have McD syndrome?

Bayesian Inference:

Example 1: Formalization

Assignment Project Exam Help

Let $McD \in \{0, 1\}$ be the variable denoting having the McD syndrome and $H \in \{0,$

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Bayesian Inference:

Example 1: Formalization

Assignment Project Exam Help

Let $McD \in \{0, 1\}$ be the variable denoting having the McD syndrome and $H \in \{0,$

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Bayesian Inference:

Example 1: Formalization

Assignment Project Exam Help

Let $McD \in \{0, 1\}$ be the variable denoting having the McD syndrome and $H \in \{0,$

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Bayesian Inference:

Example 1: Formalization

Assignment Project Exam Help

Let $McD \in \{0, 1\}$ be the variable denoting having the McD syndrome and $H \in \{0,$

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Bayesian Inference:

Example 1: Formalization

Assignment Project Exam Help

Let $McD \in \{0, 1\}$ be the variable denoting having the McD syndrome and $H \in \{0, 1\}$ be the variable denoting having the Huntington's disease.

<https://eduassistpro.github.io>

We need to compute $p(McD = 1 | H = 1)$,
the probability of having the McD syndrome given that the patient has Huntington's disease.

Add WeChat [edu_assist_pro](https://eduassistpro.github.io)

Bayesian Inference:

Example 1: Formalization

Assignment Project Exam Help

Let $McD \in \{0, 1\}$ be the variable denoting having the McD syndrome and $H \in \{0, 1\}$ be the variable denoting having the Huntington's disease.

<https://eduassistpro.github.io>

We need to compute $p(McD = 1 | H = 1)$,
the probability of having the McD syndrome given having the Huntington's disease.

Any ballpark estimates of this probability?

Bayesian Inference:

Example 1: Solution

Assignment Project Exam Help

<https://eduassistpro.github.io/>¹⁾
= . ×

Add WeChat edu_assist_pr

Bayesian Inference:

Example 1: Solution

Assignment Project Exam Help

$\frac{1}{n} \sum_{i=1}^n \log \pi(y_i | \theta)$
 $= \frac{1}{n} \sum_{i=1}^n \log \pi(y_i | \theta)$

Repeat the above computation if the proportion of h
rather small. (say in France) 0.001.

Add WeChat edu_assist_pr

Example 2: Detecting Terrorists:

From understandinguncertainty.org

Assignment Project Exam Help

- Scanner detects true terrorists with 95% accuracy

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 2: Detecting Terrorists:

From understandinguncertainty.org

Assignment Project Exam Help

- Scanner detects true terrorists with 95% accuracy

- Sca

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 2: Detecting Terrorists:

From understandinguncertainty.org

Assignment Project Exam Help

- Scanner detects true terrorists with 95% accuracy

- Sca

<https://eduassistpro.github.io>

- There is 1 terrorist on your plane with 100 passe

Add WeChat edu_assist_pr

Example 2: Detecting Terrorists:

From understandinguncertainty.org

Assignment Project Exam Help

- Scanner detects true terrorists with 95% accuracy

- Scanner

<https://eduassistpro.github.io>

- There is 1 terrorist on your plane with 100 passengers

Add WeChat edu_assist_pro

- The shiftily looking man sitting next to you tests positive

Example 2: Detecting Terrorists:

From understandinguncertainty.org

Assignment Project Exam Help

- Scanner detects true terrorists with 95% accuracy

- Sca

<https://eduassistpro.github.io>

- There is 1 terrorist on your plane with 100 passe

Add WeChat edu_assist_pr

- The shifty looking man sitting next to you tests p

What are the chances of this man being a terrorist?

Example 2: Detecting Terrorists:

Simple Solution Using “Natural Frequencies” (David Spiegelhalter)

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat Figure reproduced from [an interview with](#) edu_assist_pro

Example 2: Detecting Terrorists:

Simple Solution Using “Natural Frequencies” (David Spiegelhalter)

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat Figure reproduced from [inters and in](#) edu_assist_pro

The chances of the man being a terrori $\approx \frac{1}{6}$

Example 2: Detecting Terrorists:

Simple Solution Using “Natural Frequencies” (David Spiegelhalter)

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat Figure reproduced from [inters.aid.in](#) edu_assist_pr

The chances of the man being a terrori $\approx \frac{1}{6}$

- Relation to disease example
- Consequences when catching criminals

Example 2: Detecting Terrorists:

Formalization with Actual Probabilities

Assignment Project Exam Help

Let $T \in \{0, 1\}$ denote the variable regarding whether the person is a terrorist a

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 2: Detecting Terrorists:

Formalization with Actual Probabilities

Assignment Project Exam Help

Let $T \in \{0, 1\}$ denote the variable regarding whether the person is a terrorist a

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 2: Detecting Terrorists:

Formalization with Actual Probabilities

Assignment Project Exam Help

Let $T \in \{0, 1\}$ denote the variable regarding whether the person is a terrorist a

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 2: Detecting Terrorists:

Formalization with Actual Probabilities

Assignment Project Exam Help

Let $T \in \{0, 1\}$ denote the variable regarding whether the person is a terrorist a

<https://eduassistpro.github.io>

$$p(T = 1) = 0.01$$

Add WeChat edu_assist_pr

Example 2: Detecting Terrorists:

Formalization with Actual Probabilities

Assignment Project Exam Help

Let $T \in \{0, 1\}$ denote the variable regarding whether the person is a terrorist a

<https://eduassistpro.github.io>

$$p(T = 1) = 0.01$$

We want to compute $p(T = 1 | S = 1)$, the terrorist given that he has tested positive.

Example 2: Detecting Terrorists:

Solution with Bayes' Rule

Assignment Project Exam Help

$p(T =$

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 2: Detecting Terrorists:

Solution with Bayes' Rule

Assignment Project Exam Help

$p(T =$

<https://eduassistpro.github.io>

$\frac{(0.95)(0.01) + (0.05)}$

Add WeChat edu_assist_pr

Example 2: Detecting Terrorists:

Solution with Bayes' Rule

Assignment Project Exam Help

$$p(T = 1) = \frac{(0.95)(0.01) + (0.05)(0.01)}{(0.95)(0.01) + (0.05)(0.01) + (0.99)(0.99)}$$

≈ 0.11

Add WeChat edu_assist_pr

Example 2: Detecting Terrorists:

Solution with Bayes' Rule

Assignment Project Exam Help

$$p(T = 1) = \frac{(0.95)(0.01) + (0.05)(0.01)}{(0.95)(0.01) + (0.05)(0.01)}$$

Add WeChat edu_assist_pr

The probability of the man being a terrorist is $\bar{6}$

Example 2: Detecting Terrorists:

Posterior Versus Prior Belief

While the man has a low probability of being a terrorist, our belief has

increased compared to our prior

$$\frac{p(T = 1 | S = 1)}{p(T = 1)} = \frac{0.16}{0.01}$$

i.e. our belief

Since terrorists are so rare, a factor of 16 does not result

(absolute) probability or belief

(Aside: They are indeed very rare. For an intriguing (and surprising) example of the implications of inability to take account of actual base rates (in the example above we made the numbers up), and the effect on people's subsequent decisions, see Gerd Gigerenzer, Dread Risk, September 11, and Fatal Traffic Accidents, *Psychological Science* 15(4), 286–287, (2004); Gerd Gigerenzer, Out of the Frying Pan into the Fire: Behavioural Reactions to Terrorist Attacks, *Risk Analysis* 26(2), 347–351 (2006). His calculation (which of course is based on some assumptions) is that in the year following 9/11, 6 times the number of people who were killed as passengers *additionally* died on roads (that is the increase in road deaths due to people choosing to drive instead of flying)! He calls the reaction to very low probability events with a bad outcome “dread risk”.)

Example 3: The Monty Hall Problem

Problem Statement

- Three boxes, one with a prize and the other two are empty

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 3: The Monty Hall Problem

Problem Statement

- Three boxes, one with a prize and the other two are empty
- Each box has equal probability of having the prize

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 3: The Monty Hall Problem

Problem Statement

- Three boxes, one with a prize and the other two are empty
- Each box has equal probability of having the prize

- Yo

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 3: The Monty Hall Problem

Problem Statement

- Three boxes, one with a prize and the other two are empty
- Each box has equal probability of having the prize
- You
- You select one of the boxes

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 3: The Monty Hall Problem

Problem Statement

- Three boxes, one with a prize and the other two are empty
- Each box has equal probability of having the prize
- You
- You select one of the boxes
- The host, who knows the location of the prize, takes out of the other two boxes

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pro

Example 3: The Monty Hall Problem

Problem Statement

- Three boxes, one with a prize and the other two are empty
- Each box has equal probability of having the prize
- You choose one of the boxes
- The host, who knows the location of the prize, removes one of the other two boxes

Should you switch to the other box? Would that increase your chances of winning the prize?

Example 3: The Monty Hall Problem:

Formalization

Let $C \in \{r, g, b\}$ denote the box that contains the prize where r, g, b refer to the identity of each box.

WLOG assume the following:

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 3: The Monty Hall Problem:

Formalization

Let $C \in \{r, g, b\}$ denote the box that contains the prize where r, g, b refer to the identity of each box.

WLOG assume the following:

- Yo

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 3: The Monty Hall Problem:

Formalization

Let $C \in \{r, g, b\}$ denote the box that contains the prize where r, g, b refer to the identity of each box.

WLOG assume the following:

- Yo
- De

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 3: The Monty Hall Problem:

Formalization

Let $C \in \{r, g, b\}$ denote the box that contains the prize where r, g, b refer to the identity of each box.

WLOG assume the following:

- Yo
- De

<https://eduassistpro.github.io>

Add WeChat [edu_assist_pro](#)

Example 3: The Monty Hall Problem:

Formalization

Let $C \in \{r, g, b\}$ denote the box that contains the prize where r, g, b refer to the identity of each box.

WLOG assume the following:

- You

- De

<https://eduassistpro.github.io>

Add WeChat [edu_assist_pro](https://eduassistpro.github.io)

$$p(C = r) = \frac{1}{3} \quad p(C = g) = \frac{1}{3} \quad p(C = b) = \frac{1}{3}$$
$$p(H = b | C = r) = \frac{1}{2} \quad p(H = b | C = g) = 0 \quad p(H = b | C = b) = 0$$

Example 3: The Monty Hall Problem:

Formalization

Let $C \in \{r, g, b\}$ denote the box that contains the prize where r, g, b refer to the identity of each box.

WLOG assume the following:

- You

- De

<https://eduassistpro.github.io>

$$\begin{aligned} p(C = r) &= \frac{1}{3} & p(C = g) &= \frac{1}{3} \\ p(H = b | C = r) &= \frac{1}{2} & p(H = b | C = g) &= 0 \end{aligned}$$

We want to compute $p(C = r | H = b)$ and $p(C = g | H = b)$ to decide if we should switch from our initial choice.

Example 3: The Monty Hall Problem:

Solution

We have that:

$$P(H = b) = \sum_{c \in \{r, g, b\}} P(H = b | C = c) P(C = c)$$

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 3: The Monty Hall Problem:

Solution

We have that:

$$P(H = b) = \sum_{c \in \{r, g, b\}} P(H = b | C = c) P(C = c)$$

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 3: The Monty Hall Problem:

Solution

We have that:

$$P(H = b) = \sum_{c \in \{r, g, b\}} P(H = b | C = c) P(C = c)$$

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Example 3: The Monty Hall Problem:

Solution

We have that:

$$P(H = b) = \sum_{c \in \{r, g, b\}} P(H = b | C = c) P(C = c)$$

Assignment Project Exam Help

<https://eduassistpro.github.io>

Therefo

Add WeChat edu_assist_pr

Example 3: The Monty Hall Problem:

Solution

We have that:

$$p(H = b) = \sum_{c \in \{r, g, b\}} p(H = b | C = c) p(C = c)$$

<https://eduassistpro.github.io>

Therefo

$$p(C = r | H = b) = \frac{p(H = b | C = r) p(C = r)}{p(H = b)}$$

Example 3: The Monty Hall Problem:

Solution

We have that:

$$p(H = b) = \sum_{c \in \{r, g, b\}} p(H = b | C = c) p(C = c)$$

<https://eduassistpro.github.io>

Therefo

$$p(C = r | H = b) = \frac{p(H = b | C = r) p(C = r)}{p(H = b)}$$

Example 3: The Monty Hall Problem:

Solution

We have that:

$$p(H = b) = \sum_{c \in \{r, g, b\}} p(H = b | C = c) p(C = c)$$

<https://eduassistpro.github.io>

Therefo

$$p(C = r | H = b) = \frac{p(H = b | C = r) p(C = r)}{p(H = b)} = \frac{1/3}{1/3}$$

Similarly, $p(C = g | H = b) = 2/3$.

Example 3: The Monty Hall Problem:

Solution

We have that:

$$p(H = b) = \sum_{c \in \{r, g, b\}} p(H = b | C = c) p(C = c)$$

<https://eduassistpro.github.io>

Therefo

$$p(C = r | H = b) = \frac{p(H = b | C = r) p(C = r)}{p(H = b)} = \frac{1/3}{1/3}$$

Similarly, $p(C = g | H = b) = 2/3$.

You should switch from your initial choice to the other box in order to increase your chances of winning the prize!

Example 3: The Monty Hall Problem:

Illustration of the Solution

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pro

Illustration of the solution when you have initially selected box r .

Example 3: The Monty Hall Problem:

Another Perspective

Assignment Project Exam Help

Switching is bad if, and only if, we initially picked the prize box (because if not, the other

We picked
host's action

Hence, with probability $2/3$ switching will reveal

<https://eduassistpro.github.io>

Add WeChat [edu_assist_pro](#)

Example 3: The Monty Hall Problem:

Variants to Ponder

Assignment Project Exam Help

Would switching be rational if

- The host only revealed a box when he knew we picked the right one?

- The
one?

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

- The host is himself unaware of the prize box, an random, which by chance does not have the prize?

1 Bayes' Rule: Examples

- Eating Hamburgers

- Detecting Spam

- The Monty Hall Problem

Assignment Project Exam Help

2 Mom

<https://eduassistpro.github.io>

3 The meaning of Probability

Add WeChat edu_assist_pr

4 Wrapping Up

The Expected Value of a Function of Two Discrete Random Variables

(Assuming you have met **Expectation** $E[X]$ and **Variance** $\text{Var}(X)$ before.)

The expected value of a function $g(X, Y)$ of two discrete random variables is defined

<https://eduassistpro.github.io> (1)

In particular, the expected value of X is given

$$E[X] = \sum_x \sum_y xp(X, Y) \quad (2)$$

It should be noted that if we have already calculated the marginal distribution of X , then it is simpler to calculate $E[X]$ using this.

Covariance and the Correlation Coefficient

The covariance between X and Y , $\text{Cov}(X, Y)$ is given by

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y) \quad (3)$$

Note that

The coefficient

<https://eduassistpro.github.io>

$$\rho(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)}\sqrt{\text{Var}(Y)}} \quad (4)$$

Add WeChat [edu_assist_pro](#)

Always in $[-1, 1]$.

Example

Discrete random variables X and Y have the following joint distribution:

	$Y = 0$	$Y = 1$
$X = 0$	$\frac{0}{1}$	$\frac{1}{3}$
$X = 1$	$\frac{0}{1}$	$\frac{0}{1}$

Calculate

- 1 $p(X)$
- 2 marginal distributions of X and Y
- 3 expected values and variances of X
- 4 coefficient of correlation between X and Y

Are X and Y independent?

Example

Assignment Project Exam Help

To calculate the probability of such an event, note that we sum over all the cells with

<https://eduassistpro.github.io>

$$+ p(X = 1, Y = -$$

Add WeChat edu_assist_pro

Example

Recall that

$$p(X = x) = \sum_y p(X = x, Y = y).$$

Assignment Project Exam Help

Hence,

<https://eduassistpro.github.io>

Add WeChat edu_assist_pro

Note that after obtaining $p(X = 0)$, we could calculate $p(X = 1)$ using the fact that

$$p(X = 1) = 1 - p(X = 0), \quad (5)$$

since X only takes the values 0 and 1.

Example

Similarly, Assignment Project Exam Help

1

<https://eduassistpro.github.io>

$$p(Y = 0) = \sum_{x=0} p(X = x, Y = 0) = \frac{1}{2} + 0 = \frac{1}{2}$$

Add WeChat edu_assist_pro

$$p(Y = 1) = 1 - p(Y = 0) = \frac{1}{2}$$

Example

We then calculate the expected values and variances of X and Y from these marginal distributions.

<https://eduassistpro.github.io>

$$E(X) = \sum_{y=-1}^1 y P(Y=y) = -\frac{1}{2} \times$$

Example

To calculate the variances of X and Y , $\text{Var}(X)$ and $\text{Var}(Y)$, we use the formula

$$\text{Var}(X) = E(X^2) - (E(X))^2$$

$$E(X^2) = \sum_{i=1}^2 x_i^2 p(X = x_i) = 1^2 \cdot \frac{1}{2} + 2^2 \cdot \frac{1}{2} = \frac{5}{2}$$

<https://eduassistpro.github.io>

$$E(Y^2) = \sum_{j=-1}^1 y_j^2 p(Y = y_j) = (-1)^2 \cdot \frac{1}{3} + 0^2 \cdot \frac{1}{3} + 1^2 \cdot \frac{1}{3} = \frac{2}{3}$$

Thus we get

$$\text{Var}(X) = \frac{5}{2} - \left(\frac{3}{2}\right)^2 = \frac{1}{2}$$

$$\text{Var}(Y) = \frac{2}{3} - (0)^2 = \frac{2}{3}$$

Example

To calculate the correlation coefficient, we first calculate the covariance between X and Y . We have

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y).$$

where

$$E(XY)$$

$$= 0(-1)0 + 0(0)\frac{1}{3} + 0(1)0 + \dots - \frac{1}{3} = 0$$

Thus we get

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y) = 0 - \frac{1}{3} \times 0 = 0.$$

From the definition of the correlation coefficient,

$$\rho(X, Y) = 0.$$

Example - is X and Y independent

Assignment Project Exam Help

We have t

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

1 Bayes' Rule: Examples

- Eating Hamburgers

- Detecting Spam

- The Monty Hall Problem

Assignment Project Exam Help

2 Mom

<https://eduassistpro.github.io>

3 The meaning of Probability

Add WeChat edu_assist_pr

4 Wrapping Up

The meaning of Probability

Frequentist : Frequencies of random repeatable experiments

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

The meaning of Probability

Frequentist : Frequencies of random repeatable experiments

- E.g. Prob. of biased coin landing "Heads"

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

The meaning of Probability

Frequentist : Frequencies of random repeatable experiments

- E.g. Prob. of biased coin landing "Heads"

Bayesian : Degrees of Belief

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

The meaning of Probability

Frequentist : Frequencies of random repeatable experiments

- E.g. Prob. of biased coin landing "Heads"

Bayesian : Degrees of Belief



d

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

The meaning of Probability

Frequentist : Frequencies of random repeatable experiments

- E.g. Prob. of biased coin landing "Heads"

Bayesian : Degrees of Belief



d

Cox Axio

Given $B(x)$, $B(\bar{x})$, $B(x, y)$, $B(x|y)$, $B(y)$:

- 1 Degrees of belief can be ordered
- 2 $B(x) = f[B(\bar{x})]$
- 3 $B(x, y) = g[B(x|y), B(y)]$

The meaning of Probability

Frequentist : Frequencies of random repeatable experiments

- E.g. Prob. of biased coin landing "Heads"

Bayesian : Degrees of Belief



d

Cox Axio

Given $B(x)$, $B(\bar{x})$, $B(x, y)$, $B(x|y)$, $B(y)$:

- 1 Degrees of belief can be ordered
- 2 $B(x) = f[B(\bar{x})]$
- 3 $B(x, y) = g[B(x|y), B(y)]$

If a set of Beliefs satisfy these axioms they can be mapped onto probabilities satisfying the rules of probability.

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Image from <http://normaldeviate.wordpress.com/2012/11/09/anti-xkcd/>

In practice one needs to make use of both interpretations. Wise to be open to both. This is a huge topic which we can not get into further here. Note that Mackay was firmly in the Bayesian camp. . .

1 Bayes' Rule: Examples

- Eating Hamburgers

- Detecting Spam Emails

- The Monty Hall Problem

Assignment Project Exam Help

2 Mom

<https://eduassistpro.github.io>

3 The meaning of Probability

Add WeChat edu_assist_pr

4 Wrapping Up

Summary

- Examples of application of Bayes' rule
- Formalization

- Intuition <https://eduassistpro.github.io/>

- Frequentist vs Bayesian probabilities

- Cox axioms

Next time

Assignment Project Exam Help

- Working through some useful probability distributions

<https://eduassistpro.github.io>

- More on Bayesian inference

Add WeChat edu_assist_pro