

COMP2610 / COMP6261 Information Theory

Lecture 2: First Steps and Basic Probability

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat  edu\_assist\_pro

July 24, 2018

## Outline

1 A General Communication System

2 The R

3 Basic

4 Relating Joint, Marginal and Conditional Prob

5 Wrapping Up

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

- 1 A General Communication System

# Assignment Project Exam Help

- 2 The Role of Uncertainty

- 3 Basic

<https://eduassistpro.github.io>

- 4 Relating Joint, Marginal and Conditional Prob

Add WeChat edu\_assist\_pr

- 5 Wrapping Up

# Assignment Project Exam Help

<https://eduassistpro.github.io>

**Source** : The information source that generates the message to be communicated

**Encoder** : Operates on the message to produce a signal for transmission

**Channel** : The medium used to transmit the signal

**Decoder** : Reconstructs the message from the signal

**Destination** : The entity that receives the message

# Communication over Noisy Channels

## Assignment Project Exam Help

A channel is some medium for transmitting messages

A noisy channel  
sender's


<https://eduassistpro.github.io>

### The Problem

"The fundamental problem of communication is that at the receiver's point either exactly or approximately a message is received."  
(Claude Shannon, 1948)

Add WeChat: edu\_assist\_pro

## Example: Telephone Network



Source : Aditya

Encoder : Telephone handset

Channel : Analogue telephone line

Decoder : Telephone handset

Destination : Mark

## Examples of Noisy Communication Channels

Other examples of noisy channels:

- A radio communication link

- VDSL NBN connection

- The internet

- Reproducing cells

- A magnetic hard disk drive

- ▶ Channel does not need to involve physical movement

What would the other components be for each of these channels?

1 A General Communication System

# Assignment Project Exam Help

2 The Role of Uncertainty

3 Basic

<https://eduassistpro.github.io>

4 Relating Joint, Marginal and Conditional Prob

Add WeChat edu\_assist\_pr

5 Wrapping Up



# Assignment Project Exam Help

We can not avoid uncertainty when

- 1 Dealing with noise (imperfections) in the channel

<https://eduassistpro.github.io>

- 2 “Compressing” the messages (compare a h  
of a manuscript with the typed text that capture  
transcript of a spoken utterance)

Add WeChat edu\_assist\_pr

# Assignment Project Exam Help

A noisy ch

Thus, rec

<https://eduassistpro.github.io>

How to model, quantify, and mitigate this uncertain

Add WeChat edu\_assist\_pr

# Message Compression – I

Cover and Thomas, Example 1.1.2

## Assignment Project Exam Help

- Suppose we'd like to relay the outcome of an 8 horse race to a friend
  - ▶ We wish to convey one of  $A, B, \dots, H$

- Suppose

<https://eduassistpro.github.io>

Add WeChat [edu\\_assist\\_pro](#)

$B \rightarrow 0$

$C \rightarrow 0$

$\vdots$

$H \rightarrow 111$

## Message Compression – II

Cover and Thomas, Example 1.1.2

- Now say the probabilities of the horses winning are

( $1/2, 1/4, 1/8, 1/16, 1/64, 1/64, 1/64, 1/64$ .)

- Encoding messages based on their probability of the being chosen will g

<https://eduassistpro.github.io>

C  $\rightarrow$  110

D  $\rightarrow$  111

E  $\rightarrow$  111

F  $\rightarrow$  111100

G  $\rightarrow$  111101

H  $\rightarrow$  111111

# What is “Information”?

Assignment Project Exam Help

For noise correction and message compression, we will need to quantify the **information** contained in a message

Roughly

- Un

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

- The receiver's **uncertainty** is reduced on seeing

# The Case for Probability

We run into the notion of **uncertainty** when trying to pin down:

1 How to deal with channel noise

2 Ho

3 Wh

<https://eduassistpro.github.io>

To make progress, we need to formalise uncertainty

We will do this using **probability theory**

We now commence our *review* of probability; this will be hard going if you have not met it before!

1 A General Communication System

# Assignment Project Exam Help

2 The Role of Uncertainty

3 Basic

<https://eduassistpro.github.io>

4 Relating Joint, Marginal and Conditional Prob

Add WeChat edu\_assist\_pr

5 Wrapping Up

## Probability: Example

Quantification and Manipulation of Uncertainty (Bishop, PRML, 2006)

# Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr



## Probability: Example

Quantification and Manipulation of Uncertainty (Bishop, PRML, 2006)

1 Pick a box at random

# Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

# Probability: Example

Quantification and Manipulation of Uncertainty (Bishop, PRML, 2006)

1 Pick a box at random

2 From the selected box, pick a fruit (apple or orange) at random

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

# Probability: Example

Quantification and Manipulation of Uncertainty (Bishop, PRML, 2006)

- 1 Pick a box at random
- 2 From the selected box, pick a fruit (apple or orange) at random
- 3 Ob

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Probability: Example

Quantification and Manipulation of Uncertainty (Bishop, PRML, 2006) — Cont'd

# Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

- Identity of the box is a random variable  $\in \{ , \}$
- Identity of the fruit is a random variable  $F \in \{a, o\}$

# Probability: Example

Quantification and Manipulation of Uncertainty (Bishop, PRML, 2006) — Cont'd

## Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pro

- Identity of the box is a random variable  $\in \{ , \}$
- Identity of the fruit is a random variable  $F \in \{a, o\}$

**Probability of an event:** Proportion of times it happens out of a large number of trials

# Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pro

Suppose we repeat the selection process many times, picking up the blue box 60% of the time and the red box 40% of the time

- $p(B = r) = 0.4, p(B = b) = 0.6$

## Probability: Basic Properties

By definition,  $0 \leq p(B = b) \leq 1$  and  $0 \leq p(B = r) \leq 1$

Assignment Project Exam Help

Outcomes are mutually exclusive:

<https://eduassistpro.github.io>

Outcomes are jointly exhaustive:

$$\begin{aligned} p(B = r \text{ or } B = b) &= p(B = r) + p(B = b) \\ &= p(B = r) + p(B = b) \\ &= 1 \end{aligned}$$

Add WeChat edu\_assist\_pro

# Probability

What Types of Questions Can We Answer?

## Assignment Project Exam Help

- What is the probability of picking the red box, and an apple within that box?

- What is the probability of picking the red box, and an apple within that box?

- Given that we selected a red box, what is the probability of picking an apple?

We can answer these and more complex questions by using the *rules of probability*.



## Joint Probability

What is the probability of selecting the red box **and** selecting an apple?

**Assignment Project Exam Help**

**<https://eduassistpro.github.io>**

**Add WeChat edu\_assist\_pr**

## Joint Probability

What is the probability of selecting the red box **and** selecting an apple?

**Assignment Project Exam Help**

Joint Probability of a Set of Events

The proportion of times these events happened **together** out of the total number of

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pro

## Joint Probability

What is the probability of selecting the red box **and** selecting an apple?

Joint Probability of a Set of Events

The proportion of times these events happened **together** out of the total number of trials

If we repeat the trials we say

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pro

$$\begin{aligned} p(B = r \text{ AND } F = a) &= p \\ &= \frac{1}{10} \\ &= 0.1 \end{aligned}$$

## Marginal Probability

What is the probability of an apple being picked, **regardless** of the box we selected?

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Marginal Probability

What is the probability of an apple being picked, **regardless** of the box we selected?

Marginal Probability of an Event

The proportion of times that this event happened out of the total number of trials.

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Marginal Probability

What is the probability of an apple being picked, **regardless** of the box we selected?

Marginal Probability of an Event

The proportion of times that this event happened out of the total number of trials.

Remem

Say that in 45 of the trials, we selected a blue box and an a

So, **irrespective of B**, an apple was selected  $45 + 1 =$

$$p(F = a) = \frac{55}{100} = \frac{11}{20}$$

## Conditional Probability

What is the probability of an apple being picked up, **given that** a red box was selected?

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Conditional Probability

What is the probability of an apple being picked up, **given that** a red box was selected?

Conditional Probability of an Event

The conditional probability of an event  $X$  with respect to an event  $Y$  is the proportion of outcomes in  $X$  that are also in  $Y$ .

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pro



## Conditional Probability

What is the probability of an apple being picked up, **given that** a red box was selected?

Conditional Probability of an Event

The conditional probability of an event  $X$  with respect to an event  $Y$  is the proportion of times  $X$  occurs **given that**  $Y$  occurs.

The trials  
apple was

We selected a red box **and** an apple 10 out of 100 times

We selected a red box (regardless of the fruit) 40 out of 100 times

$$p(F = a \text{ GIVEN } B = r) = p(F = a | B = r) = \frac{10}{40} = \frac{1}{4}$$

## Conditional Probability

What is the probability of an apple being picked up, **given that** a red box was selected?

### Conditional Probability of an Event

The conditional probability of an event  $X$  with respect to an event  $Y$  is the proportion of times  $X$  occurs **given that**  $Y$  occurs.

The trials  
apple was

We selected a red box **and** an apple 10 out of 100 times

We selected a red box (regardless of the fruit) 40 out of 100 times

$$p(F = a \text{ GIVEN } B = r) = p(F = a | B = r) = \frac{10}{40} = \frac{1}{4}$$

Can we write this in terms of the joint and marginal probabilities?

1 A General Communication System

# Assignment Project Exam Help

2 The Role of Uncertainty

3 Basic

<https://eduassistpro.github.io>

4 Relating Joint, Marginal and Conditional Prob

Add WeChat edu\_assist\_pr

5 Wrapping Up

## Joint, Marginal and Conditional Probabilities:

### A More General Formulation (1)

Consider the more general case of two random variables:

$X \in \{x_1, x_2, \dots, x_r\}$  and  $Y \in \{y_1, y_2, \dots, y_L\}$

# Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

$N$  : Total number of trials

# Joint, Marginal and Conditional Probabilities:

## A More General Formulation (1)

Consider the more general case of two random variables:

$X \in \{x_1, x_2, \dots, x_r\}$  and  $Y \in \{y_1, y_2, \dots, y_L\}$

# Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

$N$  : Total number of trials

$n_{ij}$  :  $\#(X = x_i, Y = y_j)$  =  $\#$  of times that  $x_i$  and  $y_j$  happen

# Joint, Marginal and Conditional Probabilities:

## A More General Formulation (1)

Consider the more general case of two random variables:

$X \in \{x_1, x_2, \dots, x_r\}$  and  $Y \in \{y_1, y_2, \dots, y_L\}$

# Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

$N$  : Total number of trials

$n_{ij}$  :  $\#(X = x_i, Y = y_j) = \#$  of times that  $x_i$  and  $y_j$  happen

$c_i$  :  $\#(X = x_i) = \sum_j n_{ij} = \#$  of times that  $x_i$  happens

# Joint, Marginal and Conditional Probabilities:

## A More General Formulation (1)

Consider the more general case of two random variables:

$X \in \{x_1, x_2, \dots, x_r\}$  and  $Y \in \{y_1, y_2, \dots, y_L\}$

# Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

$N$  : Total number of trials

$n_{ij}$  :  $\#(X = x_i, Y = y_j) = \#$  of times that  $x_i$  and  $y_j$  happen

$c_i$  :  $\#(X = x_i) = \sum_j n_{ij} = \#$  of times that  $x_i$  happens

$r_j$  :  $\#(Y = y_j) = \sum_i n_{ij} = \#$  of times that  $y_j$  happens

# Joint, Marginal and Conditional Probabilities:

## A More General Formulation (2)

Assignment Project Exam Help

<https://eduassistpro.github.io>

By definition

Add WeChat: edu\_assist\_pro

$$p(X = x_i, Y = y_j) = \frac{n_{ij}}{N}$$
$$p(X = x_i) = \frac{c_i}{N}$$

$$p(Y = y_j) = \frac{r_j}{N} \text{ (Marginal)}$$

$$p(Y = y_j | X = x_i) = \frac{n_{ij}}{c_i} \text{ (Conditional)}$$



# Joint, Marginal and Conditional Probabilities:

A More General Formulation (1)

## Assignment Project Exam Help

Bins and fr

<https://eduassistpro.github.io>

<b>Red</b>	30	

Verify the computations from previous section with  
Add WeChat edu\_assist\_pro

# Joint, Marginal and Conditional Probabilities:

A More General Formulation (3)

## Assignment Project Exam Help

<https://eduassistpro.github.io>

Observe

Add WeChat edu\_assist\_pro

$$\begin{aligned} p(X = x_i) &= \frac{\sum_j n_{ij}}{N} \\ &= \sum_j p(X = x_i, Y = y_j) \end{aligned}$$
$$\begin{aligned} p(Y = y_j | X = x_i) &= \frac{n_{ij}}{c_i} = \frac{n_{ij}}{N} \bigg/ \frac{c_i}{N} \\ &= p(X = x_i, Y = y_j) / p(X = x_i) \end{aligned}$$

# The Rules of Probability

Sum Rule / Marginalization :

$p(X = x_i) = \sum_j p(X = x_i, Y = y_j)$   
Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

# The Rules of Probability

Sum Rule / Marginalization :

$p(X = x_i) = \sum_j p(X = x_i, Y = y_j)$   
Assignment Project Exam Help

Product R

$\text{https://eduassistpro.github.io/}$

Add WeChat edu\_assist\_pr

# The Rules of Probability

Sum Rule / Marginalization :

$p(X = x_i) = \sum_j p(X = x_i, Y = y_j)$   
Assignment Project Exam Help

Product R

$\text{https://eduassistpro.github.io/}$

and by symmetry:

$p(Y = y_j, X = x_i) = p(X = x_i, Y = y_j)$   
Add WeChat edu\_assist\_pro

# The Rules of Probability

Sum Rule / Marginalization :

Assignment Project Exam Help

Product R

<https://eduassistpro.github.io>

and by symmetry:

Add WeChat edu\_assist\_pro

Therefore:

$$P(X = x_i) = \sum_j P(X = x_i, Y = y_j) = \sum_j P(X = x_i | Y = y_j) P(Y = y_j)$$

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

Assignment Project Exam Help

<https://eduassistpro.github.io>


Add WeChat edu\_assist\_pr



# Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr




marginal

# Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr



marginal

conditional

# Joint, Marginal and Conditional Probabilities:

An even More General Formulation

Assignment Project Exam Help

Given  $n$  random variables  $X_1, \dots, X_n$

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

# Joint, Marginal and Conditional Probabilities:

An even More General Formulation

Assignment Project Exam Help

Given  $n$  random variables  $X_1, \dots, X_n$

Chain Rule

<https://eduassistpro.github.io>

$$p(X_1, X_2) = p(X_1)p(X_2|X_1)$$

What are we

Add WeChat edu\_assist\_pro

# Joint, Marginal and Conditional Probabilities:

An even More General Formulation

## Assignment Project Exam Help

Given  $n$  random variables  $X_1, \dots, X_n$

Chain Rule

<https://eduassistpro.github.io>

$$p(X_1, X_2) = p(X_1)p(X_2|X_1)$$

What are we

$$p(X_1, X_2, X_3) = p(X_1, X_2)p(X_3|X_1, X_2)$$

Add WeChat [edu\\_assist\\_pro](https://eduassistpro.github.io)

# Joint, Marginal and Conditional Probabilities:

An even More General Formulation

## Assignment Project Exam Help

Given  $D$  random variables  $X_1, \dots, X_D$

<https://eduassistpro.github.io>

Chain Rule

$$p(X_1, X_2) = p(X_1)p(X_2|X_1)$$

What are we

$$p(X_1, X_2, X_3) = p(X_1)p(X_2|X_1)p(X_3|X_1, X_2)$$

$$p(X_1, \dots, X_D) = p(X_1)p(X_2|X_1)p(X_3|X_2, X_1) \dots p(X_D|X_1, \dots, X_{D-1})$$

1 A General Communication System

# Assignment Project Exam Help

2 The Role of Uncertainty

3 Basic

<https://eduassistpro.github.io>

4 Relating Joint, Marginal and Conditional Prob

Add WeChat edu\_assist\_pr

5 Wrapping Up

- General architecture for communication systems

- Wh

<https://eduassistpro.github.io>

- Probability theory: joint, marginal and condit

Add WeChat edu\_assist\_pr

- Reading: Mackay § 2.1 and § 2.2; Bishop § 1.2



## Exercise

Coming Back to our Original Example

Given:  $p(B = r) = 0.4$ ,  $p(B = b) = 0.6$

Assume the fruit are selected uniformly from each box

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

- Write down all conditional probabilities  $P(F|B)$
- Evaluate the overall probabilities  $P(F)$

# Assignment Project Exam Help

- More on joint, marginal and conditional distributions

- Wh <https://eduassistpro.github.io>

- What, if anything, does  $p(X = x, Y = y)$  tell you about  $p(Y = y|X = x)$ ?

Next time

# Assignment Project Exam Help

- Sign-up for tutorials open at 9am wednesday 25 July. Will offer **three** tuto

<https://eduassistpro.github.io>

- Class rep.

Add WeChat edu\_assist\_pr