



Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

School of Computer Science and En
University of New South W

10. LINEAR PROGRAMMING

Problem:

- You are given a list of food sources f_1, f_2, \dots, f_t ;

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Problem:

- You are given a list of food sources f_1, f_2, \dots, f_k ;
- for each source f_i you are given:

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Problem:

- You are given a list of food sources f_1, f_2, \dots, f_k ;
- for each source f_i you are given:

-

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Problem:

- You are given a list of food sources f_1, f_2, \dots, f_k ;
- for each source f_i you are given:

-

-

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Problem:

- You are given a list of food sources f_1, f_2, \dots, f_k ;
- for each source f_i you are given:

-

-

-

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Problem:

- You are given a list of food sources f_1, f_2, \dots, f_k ;
- for each source f_i you are given:

-

-

-

<https://eduassistpro.github.io>

- Your task: to find a combination of quantities of food

Add WeChat edu_assist_pr

Problem:

- You are given a list of food sources f_1, f_2, \dots, f_k ;
- for each source f_i you are given:

-

-

-

- Your task: to find a combination of quantities of food sources f_i such that:
 - the total number of calories in all of the chosen food sources is equal to the recommended daily value of 2000 calories

Problem:

- You are given a list of food sources f_1, f_2, \dots, f_k ;
- for each source f_i you are given:

-

-

-

<https://eduassistpro.github.io>

- Your task: to find a combination of quantities of food sources such that:
 - the total number of calories in all of the chosen food sources is at least the recommended daily value of 2000 calories
 - the total intake of each vitamin V_j is at least the recommended daily intake of w_j milligrams for all $1 \leq j \leq 13$;

Problem:

- You are given a list of food sources f_1, f_2, \dots, f_k ;
- for each source f_i you are given:

-

-

-

-

- Your task: to find a combination of quantities of food
 - the total number of calories in all of the chosen food, not exceeding the recommended daily value of 2000 calories
 - the total intake of each vitamin V_j is at least the recommended daily intake of w_j milligrams for all $1 \leq j \leq 13$;
 - the price of all food per day is as low as possible.

Linear Programming problems - Example 1 cont.

- To obtain the corresponding constraints let us assume that we take x_i grams of each food source f_i for $1 \leq i \leq n$. Then:

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming problems - Example 1 cont.

- To obtain the corresponding constraints let us assume that we take x_i grams of each food source f_i for $1 \leq i \leq n$. Then:
 - the total number of calories must satisfy

Assignment Project Exam Help

$$\sum_{i=1}^n x_i c_i = 2000$$

<https://eduassistpro.github.io>

Add WeChat edu_assist_pro

Linear Programming problems - Example 1 cont.

- To obtain the corresponding constraints let us assume that we take x_i grams of each food source f_i for $1 \leq i \leq n$. Then:
 - the total number of calories must satisfy

Assignment Project Exam Help

$$\sum_{i=1}^n x_i c_i = 2000$$

-

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming problems - Example 1 cont.

- To obtain the corresponding constraints let us assume that we take x_i grams of each food source f_i for $1 \leq i \leq n$. Then:
 - the total number of calories must satisfy

Assignment Project Exam Help

$$\sum_{i=1}^n x_i c_i = 2000$$

-

<https://eduassistpro.github.io>

- an implicit assumption is that all the quantities are non-negative numbers,

Add WeChat edu_assist_pro

$$x_i \geq 0,$$

Linear Programming problems - Example 1 cont.

- To obtain the corresponding constraints let us assume that we take x_i grams of each food source f_i for $1 \leq i \leq n$. Then:
 - the total number of calories must satisfy

Assignment Project Exam Help

$$\sum_{i=1}^n x_i c_i = 2000$$

•

<https://eduassistpro.github.io>

- an implicit assumption is that all the quantities are non-negative numbers,

Add WeChat edu_assist_pro

$$x_i \geq 0,$$

- Our goal is to minimise the objective function which is the total cost

$$y = \sum_{i=1}^n x_i p_i.$$

Linear Programming problems - Example 1 cont.

- To obtain the corresponding constraints let us assume that we take x_i grams of each food source f_i for $1 \leq i \leq n$. Then:
 - the total number of calories must satisfy

Assignment Project Exam Help

$$\sum_{i=1}^n x_i c_i = 2000$$

•

<https://eduassistpro.github.io>

- an implicit assumption is that all the quantities are non-negative numbers,

Add WeChat edu_assist_pro

$$x_i \geq 0,$$

- Our goal is to minimise the objective function which is the total cost

$$y = \sum_{i=1}^n x_i p_i.$$

- Note that all constraints and the objective function, are **linear**.

Problem:

- Assume now that you are politician and you want to make certain promises to the electorate which will ensure that your party will win in the forthcoming elections.

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming problems - Example 2

Problem:

- Assume now that you are politician and you want to make certain promises to the electorate which will ensure that your party will win in the forthcoming elections.
- You can promise that you will build

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming problems - Example 2

Problem:

- Assume now that you are politician and you want to make certain promises to the electorate which will ensure that your party will win in the forthcoming elections.
- You can promise that you will build
 - a certain number of bridges, each 3 billion a piece;

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming problems - Example 2

Problem:

- Assume now that you are politician and you want to make certain promises to the electorate which will ensure that your party will win in the forthcoming elections.
- You can promise that you will build
 - a certain number of bridges, each 3 billion a piece;
 -

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming problems - Example 2

Problem:

- Assume now that you are politician and you want to make certain promises to the electorate which will ensure that your party will win in the forthcoming elections.
- You can promise that you will build
 - a certain number of bridges, each 3 billion a piece;
 -
 -

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming problems - Example 2

Problem:

- Assume now that you are politician and you want to make certain promises to the electorate which will ensure that your party will win in the forthcoming elections.

- You can promise that you will build

- a certain number of bridges, each 3 billion a piece;

-

-

- You

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming problems - Example 2

Problem:

- Assume now that you are politician and you want to make certain promises to the electorate which will ensure that your party will win in the forthcoming elections.
- You can promise that you will build
 - a certain number of bridges, each 3 billion a piece;
 -
 -
- You
 - suburban votes and 9% of rural votes;

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming problems - Example 2

Problem:

- Assume now that you are politician and you want to make certain promises to the electorate which will ensure that your party will win in the forthcoming elections.

- You can promise that you will build

- a certain number of bridges, each 3 billion a piece;



- You



- suburban votes and 9% of rural votes;

- each rural airport you promise brings you no suburban votes and 15% of rural votes;

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming problems - Example 2

Problem:

- Assume now that you are politician and you want to make certain promises to the electorate which will ensure that your party will win in the forthcoming elections.

- You can promise that you will build

- a certain number of bridges, each 3 billion a piece;

-

-

- You

-

- suburban votes and 9% of rural votes;

- each rural airport you promise brings you no suburban votes and 15% of rural votes;

- each olympic swimming pool promised brings 3% of suburban votes and no rural votes.

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pro

Linear Programming problems - Example 2

Problem:

- Assume now that you are politician and you want to make certain promises to the electorate which will ensure that your party will win in the forthcoming elections.
- You can promise that you will build
 - a certain number of bridges, each 3 billion a piece;
 -
 -
- You can promise that you will build
 - a certain number of rural airports, each 3 billion a piece; brings 15% of rural votes and no suburban votes;
 - a certain number of olympic swimming pools, each 3 billion a piece; brings 3% of suburban votes and no rural votes.
- In order to win, you have to get at least 51% of each of the city, suburban and rural votes.

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pro

Linear Programming problems - Example 2

Problem:

- Assume now that you are politician and you want to make certain promises to the electorate which will ensure that your party will win in the forthcoming elections.
- You can promise that you will build
 - a certain number of bridges, each 3 billion a piece;
 -
 -
- You can promise that you will build
 - 1% of suburban votes and 9% of rural votes;
 - each rural airport you promise brings you no suburban votes and 15% of rural votes;
 - each olympic swimming pool promised brings you 3% of suburban votes and no rural votes.
- In order to win, you have to get at least 51% of each of the city, suburban and rural votes.
- You wish to win the election by cleverly making a promise that **appears** that it will blow as small hole in the budget as possible, i.e., that the total cost of your promises is as low as possible.

Linear Programming problems - Example 2

- We can let the number of bridges to be built be x_b , number of airports x_a and the number of swimming pools x_p .

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming problems - Example 2

- We can let the number of bridges to be built be x_b , number of airports x_a and the number of swimming pools x_p .
- We now see that the problem amounts to minimising the objective $y = 3x_b + 2x_a + x_p$ while making sure that the following constraints are satisfied

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming problems - Example 2

- We can let the number of bridges to be built be x_b , number of airports x_a and the number of swimming pools x_p .
- We now see that the problem amounts to minimising the objective $y = 3x_b + 2x_a + x_p$ while making sure that the following constraints are satisfied

$$0.05x_b + 0.12x_p \leq 0.51 \quad (\text{securing majority of city votes})$$

$$0.0 \leq x_b \leq 0.0 \quad (\text{airports})$$

$$0.0 \leq x_p \leq 0.0 \quad (\text{swimming pools})$$

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming problems - Example 2

- We can let the number of bridges to be built be x_b , number of airports x_a and the number of swimming pools x_p .

- We now see that the problem amounts to minimising the objective $y = 3x_b + 2x_a + x_p$ while making sure that the following constraints are satisfied

$$\begin{array}{rcll} 0.05x_b & + 0.12x_p & \geq 0.51 & \text{(securing majority of city votes)} \\ 0.0 & & & \text{s)} \\ 0.0 & & & \end{array}$$

- However, there is a very significant difference with

Linear Programming problems - Example 2

- We can let the number of bridges to be built be x_b , number of airports x_a and the number of swimming pools x_p .
- We now see that the problem amounts to minimising the objective $y = 3x_b + 2x_a + x_p$ while making sure that the following constraints are satisfied

$$0.05x_b + 0.12x_p \leq 0.51 \quad (\text{securing majority of city votes})$$

$$0.0 \leq x_b \leq 0.0 \quad (\text{airports})$$

$$0.0 \leq x_p \leq 0.0 \quad (\text{swimming pools})$$

- However, there is a very significant difference with the previous problem:
 - you can eat 150 grams of chocolate, but

Linear Programming problems - Example 2

- We can let the number of bridges to be built be x_b , number of airports x_a and the number of swimming pools x_p .
- We now see that the problem amounts to minimising the objective $y = 3x_b + 2x_a + x_p$ while making sure that the following constraints are satisfied

$$0.05x_b + 0.12x_p \leq 0.51 \quad (\text{securing majority of city votes})$$

$$0.0 \leq x_b \leq 1.0 \quad (\text{number of bridges})$$

$$0.0 \leq x_p \leq 1.0 \quad (\text{number of swimming pools})$$

- However, there is a very significant difference with the previous problem:
 - you can eat 1.56 grams of chocolate, but
 - you cannot promise to build 1.56 bridges, 2. swimming pools!

Linear Programming problems - Example 2

- We can let the number of bridges to be built be x_b , number of airports x_a and the number of swimming pools x_p .
- We now see that the problem amounts to minimising the objective

$y = 3x_b + 2x_a + x_p$ while making sure that the following constraints are satisfied

$$0.05x_b + 0.12x_p \leq 0.51 \quad (\text{securing majority of city votes})$$

$$0.0 \leq x_b \leq 1.0 \quad (\text{number of bridges})$$

$$0.0 \leq x_p \leq 1.0 \quad (\text{number of swimming pools})$$

- However, there is a very significant difference with the first example:
 - you can eat 150 grams of chocolate, but
 - you cannot promise to build 1.56 bridges, 2.56 swimming pools!
- The second example is an example of an **Integer Linear Programming problem**, which requires all the solutions to be integers.

Linear Programming problems - Example 2

- We can let the number of bridges to be built be x_b , number of airports x_a and the number of swimming pools x_p .
- We now see that the problem amounts to minimising the objective

$y = 3x_b + 2x_a + x_p$ while making sure that the following constraints are satisfied

$$0.05x_b + 0.12x_p \geq 0.51 \quad (\text{securing majority of city votes})$$

$$0.0 \leq x_b \leq 1.0 \quad (\text{number of bridges})$$

$$0.0 \leq x_p \leq 1.0 \quad (\text{number of swimming pools})$$

- However, there is a very significant difference with the first example:
 - you can eat 150 grams of chocolate, but
 - you cannot promise to build 1.56 bridges, 2.56 swimming pools!
- The second example is an example of an **Integer Linear Programming problem**, which requires all the solutions to be integers.
- Such problems are MUCH harder to solve than the “plain” Linear Programming problems whose solutions can be real numbers.

- In the **standard form** the *objective* to be maximised is given by

Assignment $\sum_{j=1}^n c_j x_j$ Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

- In the **standard form** the *objective* to be maximised is given by

Assignment $\sum_{j=1}^n c_j x_j$ Project Exam Help

- the

<https://eduassistpro.github.io>

$x_j \geq 0$
Add WeChat edu_assist_pro (2)

- In the **standard form** the *objective* to be maximised is given by

Assignment Project Exam Help

- the

<https://eduassistpro.github.io>

Add WeChat $x_j \geq 0$ edu_assist_pro (2)

- Let the boldface \mathbf{x} represent a (column) vector, $\mathbf{x} = \langle x_1 \dots x_n \rangle^T$.

Linear Programming problems

- In the **standard form** the *objective* to be maximised is given by

Assignment $\sum_{j=1}^n c_j x_j$ Project Exam Help

- the

<https://eduassistpro.github.io>

$j=1$

Add WeChat $x_j \geq 0$ edu_assist_pro (2)

- Let the boldface \mathbf{x} represent a (column) vector, $\mathbf{x} = \langle x_1 \dots x_n \rangle^T$.
- To get a more compact representation of linear programs we introduce a partial ordering on vectors $\mathbf{x} \in \mathbf{R}^n$ by $\mathbf{x} \leq \mathbf{y}$ if and only if the corresponding inequalities hold coordinate-wise, i.e., if and only if $x_j \leq y_j$ for all $1 \leq j \leq n$.

Assignment Project Exam Help

- Letting $\mathbf{c} = \langle c_1, \dots, c_n \rangle^T \in \mathbf{R}^n$ and $\mathbf{b} = \langle b_1, \dots, b_m \rangle^T \in \mathbf{R}^m$, and letting A be the matrix $A = (a_{ij})$ of size $m \times n$, we get that the above problem can be formulated simply as:

-
-

<https://eduassistpro.github.io>

and

Add WeChat ^xedu_assist_pr

Assignment Project Exam Help

- Letting $\mathbf{c} = \langle c_1, \dots, c_n \rangle^T \in \mathbf{R}^n$ and $\mathbf{b} = \langle b_1, \dots, b_m \rangle^T \in \mathbf{R}^m$, and letting A be the matrix $A = (a_{ij})$ of size $m \times n$, we get that the above problem can be formulated simply as:

-
-

<https://eduassistpro.github.io>

and

- Thus, to specify a Linear Programming optimis^x have to provide a triplet $(A, \mathbf{b}, \mathbf{c})$; Add WeChat edu_assist_pr

Assignment Project Exam Help

- Letting $\mathbf{c} = \langle c_1, \dots, c_n \rangle^T \in \mathbb{R}^n$ and $\mathbf{b} = \langle b_1, \dots, b_m \rangle^T \in \mathbb{R}^m$, and letting A be the matrix $A = (a_{ij})$ of size $m \times n$, we get that the above problem can be formulated simply as:

-
-

<https://eduassistpro.github.io>

and

- Add WeChat ^xedu_assist_pro
- Thus, to specify a Linear Programming optimis
 - This is the usual form which is accepted by most standard LP solvers.

- The value of the objective for any value of the variables which makes the constraints satisfied is called a *feasible solution* of the LP problem.

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

- The value of the objective for any value of the variables which makes the constraints satisfied is called a *feasible solution* of the LP problem.

Assignment Project Exam Help

- Equality constraints of the form $\sum_{i=1}^n a_{ij}x_i = b_j$ can be replaced by two inequalities: $\sum_{i=1}^n a_{ij}x_i \geq b_j$ and $\sum_{i=1}^n a_{ij}x_i \leq b_j$; thus, we can assume that all constraints are inequalities.

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

- The value of the objective for any value of the variables which makes the constraints satisfied is called a *feasible solution* of the LP problem.

Assignment Project Exam Help

- Equality constraints of the form $\sum_{i=1}^n a_{ij}x_i = b_j$ can be replaced by two inequalities: $\sum_{i=1}^n a_{ij}x_i \geq b_j$ and $\sum_{i=1}^n a_{ij}x_i \leq b_j$; thus, we can assume that all constraints are inequalities.

- In ge
does
vari

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

- The value of the objective for any value of the variables which makes the constraints satisfied is called a *feasible solution* of the LP problem.

Assignment Project Exam Help

- Equality constraints of the form $\sum_{i=1}^n a_{ij}x_i = b_j$ can be replaced by two inequalities: $\sum_{i=1}^n a_{ij}x_i \geq b_j$ and $\sum_{i=1}^n a_{ij}x_i \leq b_j$; thus, we can assume that all constraints are inequalities.

- In ge
does
vari
- Ho
vari

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

- The value of the objective for any value of the variables which makes the constraints satisfied is called a *feasible solution* of the LP problem.

- Equality constraints of the form $\sum_{i=1}^n a_{ij}x_i = b_j$ can be replaced by two inequalities: $\sum_{i=1}^n a_{ij}x_i \leq b_j$ and $\sum_{i=1}^n a_{ij}x_i \geq b_j$; thus, we can assume that all constraints are inequalities.

- In ge
does
vari

- Ho
vari

- This poses no problem, because each occurrence of variable x_j can be replaced by the expression x_j^* are new variables satisfying the constraints

- The value of the objective for any value of the variables which makes the constraints satisfied is called a *feasible solution* of the LP problem.

- Equality constraints of the form $\sum_{i=1}^n a_{ij}x_i = b_j$ can be replaced by two inequalities: $\sum_{i=1}^n a_{ij}x_i \leq b_j$ and $\sum_{i=1}^n a_{ij}x_i \geq b_j$; thus, we can assume that all constraints are inequalities.

- In ge
does
vari

- Ho
vari

- This poses no problem, because each occurrence of variable x_j can be replaced by the expression x_j^* are new variables satisfying the constraints

- If $\mathbf{x} = (x_1, \dots, x_n)$ is a vector, we let $|\mathbf{x}| = (|x_1|, \dots, |x_n|)$. Problems are naturally translated into constraints of the form $|\mathbf{Ax}| \leq \mathbf{b}$. This also poses no problem because we can replace such constraints with two linear constraints: $\mathbf{Ax} \leq \mathbf{b}$ and $-\mathbf{Ax} \leq \mathbf{b}$ because $|x| \leq y$ if and only if $x \leq y$ and $-x \leq y$.

- Standard Form: maximize $\mathbf{c}^T \mathbf{x}$ subject to $A\mathbf{x} \leq \mathbf{b}$ and $\mathbf{x} \geq \mathbf{0}$.

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

- Standard Form: maximize $\mathbf{c}^T \mathbf{x}$ subject to $A\mathbf{x} \leq \mathbf{b}$ and $\mathbf{x} \geq \mathbf{0}$.
- Any vector \mathbf{x} which satisfies the two constraints is called a *feasible solution*, regardless of what the corresponding objective value $\mathbf{c}^T \mathbf{x}$ might be.

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming - Standard Form

- Standard Form: maximize $\mathbf{c}^T \mathbf{x}$ subject to $A\mathbf{x} \leq \mathbf{b}$ and $\mathbf{x} \geq \mathbf{0}$.
- Any vector \mathbf{x} which satisfies the two constraints is called a *feasible solution*, regardless of what the corresponding objective value $\mathbf{c}^T \mathbf{x}$ might be.
- As an example, let us consider the following optimisation problem

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming - Standard Form

- Standard Form: maximize $\mathbf{c}^T \mathbf{x}$ subject to $A\mathbf{x} \leq \mathbf{b}$ and $\mathbf{x} \geq \mathbf{0}$.
- Any vector \mathbf{x} which satisfies the two constraints is called a *feasible solution*, regardless of what the corresponding objective value $\mathbf{c}^T \mathbf{x}$ might be.
- As an example, let us consider the following optimisation problem

$$\begin{array}{ll} \text{maximize} & z(x_1, x_2, x_3) = 3x_1 + x_2 + 2x_3 \end{array} \quad (3)$$

subj

$$\begin{array}{ll} & x_1 \leq 36 \quad (4) \\ & x_2 \leq 24 \quad (5) \end{array}$$

Add WeChat edu_assist_pr

Linear Programming - Standard Form

- Standard Form: maximize $\mathbf{c}^T \mathbf{x}$ subject to $A\mathbf{x} \leq \mathbf{b}$ and $\mathbf{x} \geq \mathbf{0}$.
- Any vector \mathbf{x} which satisfies the two constraints is called a *feasible solution*, regardless of what the corresponding objective value $\mathbf{c}^T \mathbf{x}$ might be.
- As an example, let us consider the following optimisation problem

$$\begin{array}{ll} \text{maximize} & z(x_1, x_2, x_3) = 3x_1 + x_2 + 2x_3 \quad (3) \\ \text{subj} & \end{array}$$

$$\begin{array}{ll} & x_1 \leq 36 \quad (4) \\ & x_2 \leq 24 \quad (5) \end{array}$$

- How large can the value of the objective be, without violating the constraints?

Linear Programming - Standard Form

- Standard Form: maximize $\mathbf{c}^T \mathbf{x}$ subject to $A\mathbf{x} \leq \mathbf{b}$ and $\mathbf{x} \geq \mathbf{0}$.
- Any vector \mathbf{x} which satisfies the two constraints is called a *feasible solution*, regardless of what the corresponding objective value $\mathbf{c}^T \mathbf{x}$ might be.
- As an example, let us consider the following optimisation problem

$$\begin{array}{ll} \text{maximize} & z(x_1, x_2, x_3) = 3x_1 + x_2 + 2x_3 \\ \text{subj} & \end{array} \quad (3)$$

$$\begin{array}{ll} & 3x_1 + 2x_2 + 8x_3 \leq 36 \\ & 3x_1 + 3x_2 + 8x_3 \leq 24 \end{array} \quad \begin{array}{l} (4) \\ (5) \end{array}$$

- How large can the value of the objective be, without violating the constraints?
- If we add inequalities (4) and (5), we get

$$3x_1 + 3x_2 + 8x_3 \leq 54 \quad (8)$$

Linear Programming - Standard Form

- Standard Form: maximize $\mathbf{c}^T \mathbf{x}$ subject to $A\mathbf{x} \leq \mathbf{b}$ and $\mathbf{x} \geq \mathbf{0}$.
- Any vector \mathbf{x} which satisfies the two constraints is called a *feasible solution*, regardless of what the corresponding objective value $\mathbf{c}^T \mathbf{x}$ might be.
- As an example, let us consider the following optimisation problem

$$\begin{array}{ll}\text{maximize} & z(x_1, x_2, x_3) = 3x_1 + x_2 + 2x_3 \\ \text{subj} & \end{array} \quad (3)$$

$$\begin{array}{ll} & 3x_1 + 3x_2 \leq 54 \\ & 2x_1 + x_2 \leq 24 \end{array} \quad \begin{array}{l} (4) \\ (5) \end{array}$$

- How large can the value of the objective be, without violating the constraints?
 - If we add inequalities (4) and (5), we get
- $$3x_1 + 3x_2 + 8x_3 \leq 54 \quad (8)$$
- Since all variables are constrained to be non-negative, we are assured that

$$3x_1 + x_2 + 2x_3 \leq 3x_1 + 3x_2 + 8x_3 \leq 54$$

maximize: $z(x_1, x_2, x_3) = 3x_1 + x_2 + 2x_3$ (3)

with constraints: $x_1 + x_2 + 3x_3 \leq 30$ (4)

$2x_1 + 2x_2 + 5x_3 \leq 24$ (5)

$4x_1 + x_2 + 2x_3 \leq 36$ (6)

Assignment Project Exam Help

- <https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming - Standard Form

maximize: $z(x_1, x_2, x_3) = 3x_1 + x_2 + 2x_3$ (3)

with constraints: $x_1 + x_2 + 3x_3 \leq 30$ (4)

$$2x_1 + 2x_2 + 5x_3 \leq 24 \quad (5)$$

$$4x_1 + x_2 + 2x_3 \leq 36 \quad (6)$$

- The $z(x_1, x_2, x_3)$
- Can $y_1, y_2, y_3 \geq 0$ to be used to for a linear combination of

$$y_1(x_1 + x_2 + 3x_3) = 3$$

$$y_2(2x_1 + 2x_2 + 5x_3) = 24$$

$$y_3(4x_1 + x_2 + 2x_3) \leq 36y_3$$

Linear Programming - Standard Form

$$\text{maximize:} \quad z(x_1, x_2, x_3) = 3x_1 + x_2 + 2x_3 \quad (3)$$

$$\text{with constraints:} \quad x_1 + x_2 + 3x_3 \leq 30 \quad (4)$$

$$2x_1 + 2x_2 + 5x_3 \leq 24 \quad (5)$$

$$4x_1 + x_2 + 2x_3 \leq 36 \quad (6)$$

- Then, we can use the constraints (4), (5), and (6) to express $z(x_1, x_2, x_3)$ in terms of x_1, x_2, x_3 .
 - Can we use the constraints (4), (5), and (6) to express $z(x_1, x_2, x_3)$ in terms of x_1, x_2, x_3 ?
- $y_1, y_2, y_3 \geq 0$ to be used to for a linear combination of

$$\begin{aligned} y_1(x_1 + x_2 + 3x_3) &= 30y_1 \\ y_2(2x_1 + 2x_2 + 5x_3) &= 24y_2 \\ y_3(4x_1 + x_2 + 2x_3) &= 36y_3 \end{aligned}$$

- Then, summing up all these inequalities and factoring, we get

$$x_1(y_1 + 2y_2 + 4y_3) + x_2(y_1 + 2y_2 + y_3) + x_3(3y_1 + 5y_2 + 2y_3) \leq 30y_1 + 24y_2 + 36y_3$$

Linear Programming - Standard Form

$$\text{maximize:} \quad z(x_1, x_2, x_3) = 3x_1 + x_2 + 2x_3 \quad (3)$$

$$\text{with constraints:} \quad x_1 + x_2 + 3x_3 \leq 30 \quad (4)$$

$$2x_1 + 2x_2 + 5x_3 \leq 24 \quad (5)$$

$$4x_1 + x_2 - 2x_3 \leq 36 \quad (6)$$

$$x_1, x_2, x_3 \geq 0 \quad (7)$$

- So w

$$x_1(y_1 + 2y_2$$

$$\text{https://eduassistpro.github.io}$$

$$x_1 - 24y_2 + 36y_3 = 0$$

Add WeChat edu_assist_pr

Linear Programming - Standard Form

$$\text{maximize:} \quad z(x_1, x_2, x_3) = 3x_1 + x_2 + 2x_3 \quad (3)$$

$$\text{with constraints:} \quad x_1 + x_2 + 3x_3 \leq 30 \quad (4)$$

$$2x_1 + 2x_2 + 5x_3 \leq 24 \quad (5)$$

$$4x_1 + x_2 - 2x_3 \leq 36 \quad (6)$$

$$x_1, x_2, x_3 \geq 0 \quad (7)$$

- So w

$$x_1(y_1 + 2y_2$$

- If we c
and y_3 so that:

<https://eduassistpro.github.io>

Add WeChat [edu_assist_pr](#)

then

$$3x_3 + x_2 + 2x_3 \leq x_1(y_1 + 2y_2 + 4y_3) + x_2(y_1 + 2y_2 + y_3) + x_3(3y_1 + 5y_2 + 2y_3)$$

Linear Programming - Standard Form

$$\text{maximize:} \quad z(x_1, x_2, x_3) = 3x_1 + x_2 + 2x_3 \quad (3)$$

$$\text{with constraints:} \quad x_1 + x_2 + 3x_3 \leq 30 \quad (4)$$

$$2x_1 + 2x_2 + 5x_3 \leq 24 \quad (5)$$

$$4x_1 + x_2 - 2x_3 \leq 36 \quad (6)$$

$$x_1, x_2, x_3 \geq 0 \quad (7)$$

- So w

$$x_1(y_1 + 2y_2$$

- If we c
and y_3 so that:

<https://eduassistpro.github.io>

Add WeChat [edu_assist_pr](#)

$$\begin{aligned} & y_1 + 2y_2 + 4y_3 \\ & y_1 + 2y_2 + y_3 \\ & 3y_1 + 5y_2 + 2y_3 \end{aligned}$$

then

$$3x_3 + x_2 + 2x_3 \leq x_1(y_1 + 2y_2 + 4y_3) + x_2(y_1 + 2y_2 + y_3) + x_3(3y_1 + 5y_2 + 2y_3)$$

Combining this with (9) we get:

$$30y_1 + 24y_2 + 36y_3 \geq 3x_1 + x_2 + 2x_3 = z(x_1, x_2, x_3)$$

Linear Programming - Standard Form

- Consequently, in order to find as tight upper bound for our objective $z(x_1, x_2, x_3)$ of the problem P :

$$\text{maximize:} \quad z(x_1, x_2, x_3) = 3x_1 + x_2 + 2x_3 \quad (3)$$

$$\text{with constraints:} \quad x_1 + x_2 + 3x_3 \leq 30 \quad (4)$$

$$2x_1 + 2x_2 + 5x_3 \leq 24 \quad (5)$$

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming - Standard Form

- Consequently, in order to find as tight upper bound for our objective $z(x_1, x_2, x_3)$ of the problem P :

$$\text{maximize:} \quad z(x_1, x_2, x_3) = 3x_1 + x_2 + 2x_3 \quad (3)$$

$$\text{with constraints:} \quad x_1 + x_2 + 3x_3 \leq 30 \quad (4)$$

$$2x_1 + 2x_2 + 5x_3 \leq 24 \quad (5)$$

we have

$$\text{minimise:} \quad z^*(y_1, y_2, y_3) = 30 \quad (10)$$

with constraints

$$\begin{aligned} y_1 + y_2 + y_3 &\geq 30 \\ y_1, y_2, y_3 &\geq 0 \end{aligned} \quad (14)$$

Linear Programming - Standard Form

- Consequently, in order to find as tight upper bound for our objective $z(x_1, x_2, x_3)$ of the problem P :

$$\text{maximize:} \quad z(x_1, x_2, x_3) = 3x_1 + x_2 + 2x_3 \quad (3)$$

$$\text{with constraints:} \quad x_1 + x_2 + 3x_3 \leq 30 \quad (4)$$

$$2x_1 + 2x_2 + 5x_3 \leq 24 \quad (5)$$

we have <https://eduassistpro.github.io>

$$\text{minimise:} \quad z^*(y_1, y_2, y_3) = 30 \quad (10)$$

with constraints

$$\begin{aligned} y_1 & & y_2 & & y_3 & \geq \\ y_1, y_2, y_3 & \geq 0 \end{aligned} \quad (14)$$

then $z^*(y_1, y_2, y_3) = 30y_1 + 24y_2 + 36y_3 \geq 3x_1 + x_2 + 2x_3 = z(x_1, x_2, x_3)$
will be a tight upper bound for $z(x_1, x_2, x_3)$

- The new problem P^* is called the *dual problem* for the problem P .

- Let us now repeat the whole procedure with P^* in place of P , i.e., let us find the dual program $(P^*)^*$ of P^* .

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming - Standard Form

- Let us now repeat the whole procedure with P^* in place of P , i.e., let us find the dual program $(P^*)^*$ of P^* .
- We are now looking for $z_1, z_2, z_3 \geq 0$ to multiply inequalities (11)-(13) and obtain

Assignment Project Exam Help

$$z_1(y_1 + 2y_2 + 4y_3) \geq 3z_1$$
$$z_2(y_1 + 2y_2 + y_3) \geq z_2$$

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Linear Programming - Standard Form

- Let us now repeat the whole procedure with P^* in place of P , i.e., let us find the dual program $(P^*)^*$ of P^* .
- We are now looking for $z_1, z_2, z_3 \geq 0$ to multiply inequalities (11)-(13) and obtain

$$\begin{aligned} z_1(y_1 + 2y_2 + 4y_3) &\geq 3z_1 \\ z_2(y_1 + 2y_2 + y_3) &\geq z_2 \end{aligned}$$

- Su

$$y_1(z_1 + z_2) + 2y_2(z_1 + z_2) + y_3(4z_1 + z_2) \geq z_1 + z_2 + 2z_3$$

<https://eduassistpro.github.io>

(15)

Add WeChat edu_assist_pr

Linear Programming - Standard Form

- Let us now repeat the whole procedure with P^* in place of P , i.e., let us find the dual program $(P^*)^*$ of P^* .
- We are now looking for $z_1, z_2, z_3 \geq 0$ to multiply inequalities (11)-(13) and obtain

$$\begin{aligned} z_1(y_1 + 2y_2 + 4y_3) &\geq 3z_1 \\ z_2(y_1 + 2y_2 + y_3) &\geq z_2 \end{aligned}$$

- So

$$y_1(z_1 + z_2) + 2y_2(z_1 + z_2) + 4y_3z_1 + y_3z_2 \geq 3z_1 + z_2 \quad (15)$$

- If we choose multipliers z_1, z_2, z_3 so that

$$z_1 + z_2 + 3z_3 = 3 \quad (16)$$

$$2z_1 + 2z_2 + 5z_3 = 5 \quad (17)$$

$$4z_1 + z_2 + 2z_3 \leq \quad (18)$$

Linear Programming - Standard Form

- Let us now repeat the whole procedure with P^* in place of P , i.e., let us find the dual program $(P^*)^*$ of P^* .
- We are now looking for $z_1, z_2, z_3 \geq 0$ to multiply inequalities (11)-(13) and obtain

$$\begin{aligned} z_1(y_1 + 2y_2 + 4y_3) &\geq 3z_1 \\ z_2(y_1 + 2y_2 + y_3) &\geq z_2 \end{aligned}$$

- So

$$y_1(z_1 + z_2) + y_2(2z_1 + 2z_2) + y_3(4z_1 + z_2 + 2z_3) \geq 3z_1 + z_2 \quad (15)$$

- If we choose multipliers z_1, z_2, z_3 so that

$$z_1 + z_2 + 3z_3 = 3 \quad (16)$$

$$2z_1 + 2z_2 + 5z_3 = 1 \quad (17)$$

$$4z_1 + z_2 + 2z_3 \leq 0 \quad (18)$$

we will have:

$$y_1(z_1 + z_2 + 3z_3) + y_2(2z_1 + 2z_2 + 5z_3) + y_3(4z_1 + z_2 + 2z_3) \leq 3y_1 + 2y_2 + 3y_3$$

Linear Programming - Standard Form

- Let us now repeat the whole procedure with P^* in place of P , i.e., let us find the dual program $(P^*)^*$ of P^* .
- We are now looking for $z_1, z_2, z_3 \geq 0$ to multiply inequalities (11)-(13) and obtain

$$\begin{aligned} z_1(y_1 + 2y_2 + 4y_3) &\geq 3z_1 \\ z_2(y_1 + 2y_2 + y_3) &\geq z_2 \end{aligned}$$

- So

$$y_1(z_1 + z_2) + z_2 + 2z_3$$

<https://eduassistpro.github.io>

- If we choose multipliers z_1, z_2, z_3 so that

$$z_1 + z_2 + 3z_3 = 3 \quad (16)$$

Add WeChat [edu_assist_pro](https://eduassistpro.github.io)

$$2z_1 + 2z_2 + 5z_3 = 5 \quad (17)$$

$$4z_1 + z_2 + 2z_3 \leq \quad (18)$$

we will have:

$$y_1(z_1 + z_2 + 3z_3) + y_2(2z_1 + 2z_2 + 5z_3) + y_3(4z_1 + z_2 + 2z_3) \leq 30y_1 + 24y_2 + 36y_3$$

- Combining this with (15) we get

$$3z_1 + z_2 + 2z_3 \leq 30y_1 + 24y_2 + 36y_3$$

- Consequently, finding the dual program $(P^*)^*$ of P^* amounts to maximising the objective $3z_1 + z_2 + 2z_3$ subject to the constraints

$$\begin{aligned} z_1 + z_2 + 3z_3 &\leq 30 \\ 2z_1 + 2z_2 + 5z_3 &\leq 24 \end{aligned}$$

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

- Consequently, finding the dual program $(P^*)^*$ of P^* amounts to maximising the objective $3z_1 + z_2 + 2z_3$ subject to the constraints

$$\begin{aligned} z_1 + z_2 + 3z_3 &\leq 30 \\ 2z_1 + 2z_2 + 5z_3 &\leq 24 \end{aligned}$$

Assignment Project Exam Help

- But n exactly our starting program P . Thus, the dual program $(P^*)^*$ for program P^* is just P itself, i.e., $(P^*)^* = P$.

Add WeChat edu_assist_pr

- Consequently, finding the dual program $(P^*)^*$ of P^* amounts to maximising the objective $3z_1 + z_2 + 2z_3$ subject to the constraints

$$\begin{aligned} z_1 + z_2 + 5z_3 &\leq 30 \\ 2z_1 + 2z_2 + 5z_3 &\leq 24 \end{aligned}$$

Assignment Project Exam Help

<https://eduassistpro.github.io>

- But not exactly our starting program P . Thus, the dual program $(P^*)^*$ for program P^* is just P itself, i.e., $(P^*)^* = P$.

- So, at the first sight, looking for the multipliers is not much, because it only reduced a maximisation problem to a hard minimisation problem.

Add WeChat edu_assist_pro

- Consequently, finding the dual program $(P^*)^*$ of P^* amounts to maximising the objective $3z_1 + z_2 + 2z_3$ subject to the constraints

$$\begin{aligned} z_1 + z_2 + 5z_3 &\leq 30 \\ 2z_1 + 2z_2 + 5z_3 &\leq 24 \end{aligned}$$

Assignment Project Exam Help

<https://eduassistpro.github.io>

- But not exactly our starting program P . Thus, the dual program $(P^*)^*$ for program P^* is just P itself, i.e., $(P^*)^* = P$.

Add WeChat edu_assist_pro

- So, at the first sight, looking for the multipliers is not so much, because it only reduced a maximisation problem to a hard minimisation problem.
- It is now useful to remember how we proved that the Ford - Fulkerson Max Flow algorithm in fact produces a **maximal flow**, by showing that it terminates only when we reach the capacity of a **minimal cut**.

Linear Programming - primal/dual problem forms

- The original, *primal* Linear Program P and its *dual* Linear Program can be easily described in the most general case:

$$\begin{aligned} P : \text{ maximize } & z(\mathbf{x}) = \sum_{j=1}^n c_j x_j, \\ \text{subject to the constraints } & a_{ij} x_j \leq b_i; \quad 1 \leq i \leq m \end{aligned}$$

<https://eduassistpro.github.io>

Add WeChat edu_assist_pro

$$y_1, \dots, y_m \geq 0,$$

Linear Programming - primal/dual problem forms

- The original, *primal* Linear Program P and its *dual* Linear Program can be easily described in the most general case:

$$\begin{aligned} P : \text{maximize} \quad & z(\mathbf{x}) = \sum_{j=1}^n c_j x_j, \\ \text{subject to the constraints} \quad & a_{ij} x_j \leq b_i; \quad 1 \leq i \leq m \end{aligned}$$

<https://eduassistpro.github.io>

Add WeChat edu_assist_pro

or, in matrix form,

$$\begin{aligned} P : \text{maximize} \quad & z(\mathbf{x}) = \mathbf{c}^T \mathbf{x}, \text{ subject to the constraints } \mathbf{Ax} \leq \mathbf{b} \text{ and } \mathbf{x} \geq 0; \\ P^* : \text{minimize} \quad & z^*(\mathbf{y}) = \mathbf{b}^T \mathbf{y}, \text{ subject to the constraints } \mathbf{A}^T \mathbf{y} \geq \mathbf{c} \text{ and } \mathbf{y} \geq 0. \end{aligned}$$

Weak Duality Theorem

- Recall that any vector \mathbf{x} which satisfies the two constraints, $A\mathbf{x} \leq \mathbf{b}$ and $\mathbf{x} \geq 0$ is called a *feasible solution*, regardless of what the corresponding objective value $\mathbf{c}^T \mathbf{x}$ might be.

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Weak Duality Theorem

- Recall that any vector \mathbf{x} which satisfies the two constraints, $A\mathbf{x} \leq \mathbf{b}$ and $\mathbf{x} \geq 0$ is called a *feasible solution*, regardless of what the corresponding objective value $\mathbf{c}^T \mathbf{x}$ might be.

- Theorem** If $\mathbf{x} = \langle x_1, \dots, x_n \rangle$ is any basic feasible solution for P and $\mathbf{y} = \langle y_1, \dots, y_m \rangle$ is any basic feasible solution for P^* , then

n

n

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Weak Duality Theorem

- Recall that any vector \mathbf{x} which satisfies the two constraints, $A\mathbf{x} \leq \mathbf{b}$ and $\mathbf{x} \geq 0$ is called a *feasible solution*, regardless of what the corresponding objective value $\mathbf{c}^T \mathbf{x}$ might be.

Theorem If $\mathbf{x} = \langle x_1, \dots, x_n \rangle$ is any basic feasible solution for P and $\mathbf{y} = \langle y_1, \dots, y_m \rangle$ is any basic feasible solution for P^* , then

n

n

Pro
we ca

<https://eduassistpro.github.io>

respectively,

P^*

P to obtain

$$z(\mathbf{x}) = \sum_{j=1}^n c_j x_j \leq \sum_{j=1}^n \left(\sum_{i=1}^m a_{ij} y_i \right) x_j = \sum_{i=1}^m y_i \left(\sum_{j=1}^n a_{ij} x_j \right) = \sum_{i=1}^m y_i b_i = z^*(\mathbf{y})$$

Weak Duality Theorem

- Recall that any vector \mathbf{x} which satisfies the two constraints, $A\mathbf{x} \leq \mathbf{b}$ and $\mathbf{x} \geq 0$ is called a *feasible solution*, regardless of what the corresponding objective value $\mathbf{c}^T \mathbf{x}$ might be.

Theorem If $\mathbf{x} = \langle x_1, \dots, x_n \rangle$ is any basic feasible solution for P and $\mathbf{y} = \langle y_1, \dots, y_m \rangle$ is any basic feasible solution for P^* , then

n

n

Pro
we ca

respectively,

P^*

P to obtain

$$z(\mathbf{x}) = \sum_{j=1}^n c_j x_j \leq \sum_{j=1}^n \left(\sum_{i=1}^m a_{ij} y_i \right) x_j = \sum_i y_i \left(\sum_{j=1}^n a_{ij} x_j \right) = \sum_i y_i b_i = z^*(\mathbf{y})$$

- Thus, the value of (the objective of P^* for) any feasible solution of P^* is an upper bound for the set of all values of (the objective of P for) all feasible solutions of P , and

Weak Duality Theorem

- Recall that any vector \mathbf{x} which satisfies the two constraints, $A\mathbf{x} \leq \mathbf{b}$ and $\mathbf{x} \geq 0$ is called a *feasible solution*, regardless of what the corresponding objective value $\mathbf{c}^T \mathbf{x}$ might be.

Theorem If $\mathbf{x} = \langle x_1, \dots, x_n \rangle$ is any basic feasible solution for P and $\mathbf{y} = \langle y_1, \dots, y_m \rangle$ is any basic feasible solution for P^* , then

$$\mathbf{c}^T \mathbf{x} \leq \mathbf{b}^T \mathbf{y}$$


Proof
we can

respectively,
 P^* to obtain

$$z(\mathbf{x}) = \sum_{j=1}^n c_j x_j \leq \sum_{j=1}^n \left(\sum_{i=1}^m a_{ij} y_i \right) x_j = \sum_{i=1}^m y_i \left(\sum_{j=1}^n a_{ij} x_j \right) = \sum_{i=1}^m y_i b_i = z^*(\mathbf{y})$$

- Thus, the value of (the objective of P^* for) any feasible solution of P^* is an upper bound for the set of all values of (the objective of P for) all feasible solutions of P , and
- every feasible solution of P is a lower bound for the set of feasible solutions for P^* .

Assignment Project Exam Help




<https://eduassistpro.github.io>

- Thus, if we find a feasible solution for P which is equal to a feasible solution to P^* , such solution must be the maximum objective of P and the minimal feasible value of P^* .

Add WeChat edu_assist_pro


Assignment Project Exam Help



<https://eduassistpro.github.io>

- Thus, if we find a feasible solution for P which is equal to a feasible solution to P^* , such solution must be the maximum objective of P and the minimal feasible value of P^* .
- If we use a search procedure to find an optimal solution, when to stop: when such a value is also a feasible solution.


Assignment Project Exam Help



<https://eduassistpro.github.io>

- Thus, if we find a feasible solution for P which is equal to a feasible solution to P^* , such solution must be the maximum objective of P and the minimal feasible value of P^* .
- If we use a search procedure to find an optimal solution, when to stop: when such a value is also a feasible solution.
- This is why the most commonly used LP solving method, the SIMPLEX method, produces optimal solution for P , because it stops at a value of the primal objective which is also a value of the dual objective.

Assignment Project Exam Help



<https://eduassistpro.github.io>

- Thus, if we find a feasible solution for P which is equal to a feasible solution to P^* , such solution must be the maximum objective of P and the minimal feasible value of P^* .
- If we use a search procedure to find an optimal solution, when to stop: when such a value is also a feasible solution.
- This is why the most commonly used LP solving method, the SIMPLEX method, produces optimal solution for P , because it stops at a value of the primal objective which is also a value of the dual objective.
- See the Lecture Notes for the details and an example of how the SIMPLEX algorithm runs.

Assignment Project Exam Help

Five siste
is playing
Helen, th

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr