COMP4161 T3/2022 Advanced Topics in Software Verification

Assignment 1

This assignment starts on Tuesday 20th September 2022 and is due on Tuesday 27th September 2022 6pm. We will accept plain text (.txt) files, PDF (.pdf) files, and Isabelle theory (.thy) files. The assignment is take-home. This does NOT mean you can work in groups. Each submission is personal. For more information, see the plagiarism policy: https://student.unsw.edu.au/plagiarism Submit using give on a CSE machine:

```
give cs4161 a1 files ...
```

For example:

give cs4161 a1 a1.thy a1.pdf

Types (15 marks) ent Project Exam Help (a) Provide the most general type for the term λx y z. z x (a y y). Show a type derivation

tree to justify your

Each node of the typing rule to the typing rule to

Under which contexts is the term type correct? (10 marks)

(b) Find a term that the few typhat edu assist pro

```
('a \Rightarrow 'a \Rightarrow 'b \Rightarrow 'c) \Rightarrow 'a \Rightarrow 'b \Rightarrow 'c
```

(You don't need to provide a type derivation, just the term).

(5 marks)

2 λ -Calculus (23 marks)

Recall the encoding of booleans and booleans operations in lambda calculus seen in the lecture:

```
true
                  \lambda x y. x
false
                  \lambda x y. y
ifthen
                  \lambda z \times y. z \times y
                  \lambda x. ifthen x false true
not
```

- (a) Define (in Isabelle) xor, the exclusive OR operator, using the definitions of ifthen and not (3 marks).
- (b) Show by beta reduction (by hand, not using Isabelle) that:

```
xor =_{\beta} \lambda x y. x (y false true) y
then show by beta reduction that:
xor false y =_{\beta} y.
```

and

xor true $y =_{\beta} not y$.

Each step should be a single beta reduction or definition unfolding. Alpha conversion is allowed (14 marks).

(c) Prove the above 3 lemmas in Isabelle, using unfold and refl. Explain (informally) what the refl theorem states and explain why it can be used to prove the lemmas (6 marks).

3 Propositional Logic (35 marks)

Prove each of the following statements (after stating them in Isabelle for (c) and (f)), using only the proof methods

rule, erule, assumption, cases, frule, drule, rule_tac, erule_tac, frule_tac, drule_tac, rename_tac, and case_tac;

and using only the proof rules

impI, impE, conjI, conjE, disjI1, disjI2, disjE, notI, notE, iffI, iffE, iffD1, iffD2,
ccontr, classical, FalseE, TrueI, conjunct1, conjunct2, and mp.

You do not need to use all of these methods and rules.

- (a) $A \lor B \lor A \longrightarrow B \lor A$ (3 marks)
- (b) (¬ Assignment Project Exam Help (3 marks
- (c) "Saying that if Ali saying that they c (5 marks)
- (d) (A A B V C) https://eduassistpro.github.io/arks)

$$(e) \neg P \land Q \longrightarrow \neg (R \land P) \land (R \longrightarrow Q)$$
 (7 marks)

(f) "If either it is not runing of the very an uniform that is also raining." (f) "If either it is not runing of the very an uniform that is also raining."

(g)
$$(((f \longrightarrow g) \land h \longrightarrow f) \longrightarrow g) = ((f \longrightarrow g) \land (g \lor h))$$
 (7 marks)

4 Higher-Order Logic (27 marks)

Prove each of the following statements (after stating them in Isabelle for (d)), using only the proof methods and proof rules stated in the previous question, plus any of the following proof rules:

allI, allE, exI, and exE.

You do not need to use all of these methods and rules. You may use rules proved in earlier parts of the question when proving later parts.

(a)
$$(\forall x. \neg P x) = (\nexists x. P x)$$
 (4 marks)

(b)
$$(\forall x. B x) \lor (\forall y. A y) \longrightarrow (\forall x y. A y \lor B x)$$
 (4 marks)

(c)
$$(\forall x \ y. \ A \ y \lor B \ x) \longrightarrow (\forall x. \ B \ x) \lor (\forall y. \ A \ y)$$
 (7 marks)

- (d) "If any proposition is true then the value True is the same as the value False." (4 marks)
- (e) $((\exists x. A x) \longrightarrow \neg C) \longrightarrow (\forall x. B x \longrightarrow A x) \longrightarrow (\exists x. B x) \longrightarrow \neg C$ (4 marks)

(f)
$$(\forall x. \neg R \times x) \longrightarrow \neg (\forall x y. \neg R \times y \longrightarrow R \times x)$$
 (4 marks)