

Numerical Optimisation: Assignment Project Exam Help

<https://eduassistpro.github.io>

`f.rullan@cs.ucl.ac.uk`

Add WeChat `edu_assist_pro`

Department of Computer Science
Centre for Medical Image Computing
Centre for Inverse Problems
University College London

Lecture 1

Assignment Project Exam Help

$$\min_{x \in \mathbb{R}^n} f(x) \quad \text{subject to} \quad \begin{array}{ll} c_i(x) = 0, & i \in \mathcal{E} \\ c_i(x) \geq 0, & i \in \mathcal{I} \end{array}$$

- <https://eduassistpro.github.io>

$i \in \mathcal{I}$ inequality constraints.

- $x \in \mathbb{R}^n$: optimisation variable

Optimal solution x^* has the smallest value of f among all x which satisfy the constraints.

Example: geodesics

Geodesics are the shortest surface paths between two points.

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Figure: <https://academo.org/demos/geodesics/>

A very short and incomplete early history

Source <http://www.mitrikitti.fi/opthist.html>

- **Antiquity: geometrical optimisation problems**

300 BC Euclid considers the minimal distance between a point and a line, and proves that a square has the greatest area among the rectangles with given total length of edges



<https://eduassistpro.github.io>

smallest variation of volume w.r.t. barrel p

¹

Early version of the secretary problem (opt problem) when he started to look for a new wife

1636 P. Fermat shows that at the extreme po

of a function vanishes. In 1657 Fermat shows that light travels between two points in minimal time.

¹<http://www.maa.org/press/periodicals/convergence/kepler-the-volume-of-a-wine-barrel-solving-the-problem-of-maxima-wine-barrel-design>

Source <http://www.mit-ikitti.fi/opthist.html>

Assignment Project Exam Help

- Calculus of Variations

<https://eduassistpro.github.io>

1696 Johann and Jacob Bernoulli study Br

problem, calculus of variations is born

1740 L. Euler's publication begins the reser

theory of calculus of variations

A very short and incomplete early history cont.

Source <http://www.mitrikitti.fi/opthist.html>

- **Least squares**

1806 A. M. Legendre presents the least square method, which also J.C.F. Gauss claims to have invented. Legendre made contributions in the field of CoV, too



<https://eduassistpro.github.io>

1939 L.V. Kantorovich presents LP-mod

for solving it. In 1975 Kantorovich and T.C. K

the Nobel memorial price in economics for t

on LP-problem

1947 G. Dantzig, who works for US air-force, presents the Simplex method for solving LP-problems, von Neumann establishes the theory of duality for LP-problems

- 2 factories, F_i
- 12 retail outlets, R_j
- each factory F_i can produce up to a_i tones of a certain

- <https://eduassistpro.github.io>

- the cost of shipping of one tone of the compound to R_j is c_{ij}

Add WeChat edu_assist_pr

Goal: what is the optimal amount to ship from each factory to each outlet which satisfies demand at minimal cost.

Assignment Project Exam Help

12

<https://eduassistpro.github.io>

$$x_{ij} \geq b_j, j = 1 \dots 12$$

$i=1$

Add WeChat [edu_assist_pro](https://eduassistpro.github.io)



Linear programming problem because the objective function and all constraints are linear.

Assignment Project Exam Help

$$\begin{aligned} \min & f(x, y) \\ \text{subject to} & -\frac{1}{2}y + \frac{1}{2}x - \frac{1}{2} \geq 0 \end{aligned}$$

<https://eduassistpro.github.io>

Add WeChat edu_assist_pro



A set $\mathbb{S} \subset \mathbb{R}^n$ is **convex** if for any two points $x, y \in \mathbb{S}$ the line segment connecting them lies entirely in \mathbb{S}

$$\alpha x + (1 - \alpha)y \in \mathbb{S}, \quad \forall \alpha \in [0, 1].$$

Examples:




Figure: (a) unit ball $\{x \in \mathbb{R}^n : \|x\|_p \leq 1\}$, $p \geq 1$; (b) polyhedron $\{x \in \mathbb{R}^n : Ax = b, Cx \leq d\}$

A function f is **convex** if

- its domain \mathbb{S} is a convex set,

- for any two points $x, y \in \mathbb{S}$,

$$f(\alpha x + (1 - \alpha)y) \leq \alpha f(x) + (1 - \alpha)f(y), \quad \alpha \in [0, 1].$$

A function

<https://eduassistpro.github.io/>

A function f is **concave** if $-f$ is convex.

Examples:

- linear function $f(x) = c^T x + \alpha$, where $c \in \mathbb{R}^n$, $\alpha \in \mathbb{R}$
- convex quadratic function $f(x) = x^T H x$, where $H \in \mathbb{R}^{n \times n}$ symmetric positive (semi)definite

Assignment Project Exam Help

- **convex** vs **non-convex**
- **smooth** vs **non-smooth**

-

- <https://eduassistpro.github.io>
-

- **local** vs **global**
- **stochastic** vs **deterministic**
- discrete vs **continuous**

Add WeChat edu_assist_pro

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a smooth function for which we can evaluate f and its derivatives at any given point $x \in \Omega \subseteq \mathbb{R}^n$.

Unconstraint optimisation problem

Assignment Project Exam Help

$$\min f(x). \quad (1)$$

A point <https://eduassistpro.github.io>

$f(x^*) \leq f(x), \forall x$
Add WeChat edu_assist_pr

A point x^* is a **local minimiser** if

$$\exists \mathcal{N}(x^*) : f(x^*) \leq f(x), \forall x \in \mathcal{N}(x^*),$$

$\mathcal{N}(y)$ is a neighbourhood of y (an open set which contains y).

A point x^* is a **strict (or strong) local minimiser** if

$$(x^*) : f(x^*) < f(x), \quad x \in (x^*), \quad x \neq x^*.$$

Exa

- <https://eduassistpro.github.io>
- $f(x) = (x - 2)^4$: $x^* = 2$ is a strict local
global one)
- $f(x) = \cos(x)$: $x^* = \pi + 2k\pi, k \in \mathbb{Z}$
minimisers (but not strict global on

A point x^* is an **isolated local minimiser** if

Assignment Project Exam Help

$\exists \delta(x^*) > 0$ such that x^* is the only local minimiser in $N_\delta(x^*)$.

Some

<https://eduassistpro.github.io>

has a strict local minimiser at $x^* = 0$ but it

is not an isolated local minimiser. **Add WeChat: edu_assist_pro**

minimisers: $f(x_j) = x_j^4 > 0 = f(0)$ for $x_j \neq 0$, $j \rightarrow \infty$.

However, all isolated local minimiser are strict.

Unconstraint minimisation



Difficulties with global minimisation:

$$f(x) = (\cos(20\pi x) + 2)|x|$$

has a unique global minimiser $x^* = 0$, but the algorithms usually get trapped into one of the many local minima.

For co
mini

<https://eduassistpro.github.io>



Add WeChat edu_assist_pro

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuously differentiable. Then for $p \in \mathbb{R}^n$ we have

Assignment Project Exam Help

for some $t \in (0, 1)$.

If mor

<https://eduassistpro.github.io>

0

and

Add WeChat edu_assist_pr

$$f(x + p) = f(x) + \nabla f(x)^T p + \frac{1}{2} p^T \nabla^2 f(x + tp) p,$$

fore some $t \in (0, 1)$.

Theorem [1st order necessary condition]

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuously differentiable in an open neighbourhood of a **local minimiser** x^* , then $\nabla f(x^*) = 0$.

Proof: [by contradiction]

Suppose that $\nabla f(x^*) \neq 0$ and define $p = -\nabla f(x^*)$. Note that

$p^T \nabla f(x^*) = -\|\nabla f(x^*)\|^2 < 0$ near

<https://eduassistpro.github.io>

By Taylor's theorem, for any $\bar{t} \in (0, T]$

$$f(x^* + \bar{t}p) = f(x^*) + \bar{t}p^T \nabla f(x^*) + o(\bar{t})$$

Hence $f(x^* + \bar{t}p) < f(x^*)$ for all $\bar{t} \in (0, T]$, and we have found a direction leading away from x^* along which f decreases which is in contradiction with x^* being a local minimiser.

Assignment Project Exam Help

We ca

By Th
a stati

<https://eduassistpro.github.io>

Add WeChat edu_assist_pr

Theorem [2nd order necessary condition]

If x^* is a **local minimiser** of f and $\nabla^2 f$ exists and is continuous in an open neighbourhood of x^* , then $\nabla f(x^*) = 0$ and $\nabla^2 f(x^*)$ is **positive semidefinite**.

Proof: [by contradiction]

By Th

$\nabla f(x^*) = 0$.

Assu

vect

conti

$p^T \nabla^2 f(x^* + tp)p < 0$ for all $t \in [0, T]$

By Taylor theorem we have for any \bar{t}

$$f(x^* + \bar{t}p) = f(x^*) + \bar{t}p^T \nabla f(x^*) + \frac{1}{2} \bar{t}^2 p^T \nabla^2 f(x^* + \bar{t}p)p < f(x^*).$$

We have found a decrease direction for f away from x^* which contradicts x^* being a local minimiser.

Theorem [2nd order sufficient condition]

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ with $\nabla^2 f$ continuous in an open neighbourhood of x^* . If $\nabla f(x^*) = 0$ and $\nabla^2 f(x^*)$ is **positive definite**, then x^* is a **strict local minimiser** of f .

Proof:

Because the Hessian $\nabla^2 f$ is continuous and positive definite at x^* , we can

define

For a

<https://eduassistpro.github.io>

$$\begin{aligned} f(x^* + p) &\stackrel{(2)}{=} f(x^*) + p^T \nabla f(x^*) \\ &= f(x^*) + \frac{1}{2} p^T \nabla^2 f(x^*) p \end{aligned} \quad (3)$$

for some $t \in (0, 1)$.

Furthermore, $x^* + tp \in B_2(x^*, r)$ thus $p^T \nabla^2 f(x^* + tp) p > 0$ and therefore $f(x^* + p) > f(x^*)$.

Assignment Project Exam Help

2nd order sufficient condition guarantees a stronger statement than the necessary conditions (strict local minimiser).

A strict

<https://eduassistpro.github.io>

$$f(x) = x^2, \quad f(x) = 4x^2, \quad f(x) = 12x^2$$

$x^* = 0$ is a strict local minimiser while

the necessary but not the sufficient conditions

Add WeChat [edu_assist_pro](#)

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex, any local minimiser x^* is also a global minimiser of f . If, f is also differentiable, then any stationary point x^* is a global minimiser.

Proof:

Supp

*

$\exists z \in$

z i.e.

x^* and

<https://eduassistpro.github.io>

$$\mathcal{L}(x^*, z) = \{x : x = \lambda z + (1 -$$

by convexity of f we have

Add WeChat edu_assist_pr

$$f(x) \leq \lambda f(z) + (1 - \lambda)f(x^*) < f(x^*).$$

For any neighbourhood $\mathcal{N}(x^*) \cap \mathcal{L}(x^*, z) \neq \emptyset$, hence $\exists x \in \mathcal{N}(x^*) : f(x) < f(x^*)$ and x^* is not a local minimiser.

Proof: cont.

For the second part, we suppose that x^* is not global minimiser.
For all z chosen as before by convexity of f it follows

$$\begin{aligned} & \lim_{\lambda \rightarrow 0} \frac{\lambda f(z)}{\lambda f(z) + (1-\lambda)f(x^*)} \\ & \leq \lim_{\lambda \rightarrow 0} \frac{\lambda f(z)}{\lambda f(z) + (1-\lambda)f(x^*)} \\ & = f(z) - f(x^*) \end{aligned}$$

Hence $\nabla f(x^*) \neq 0$ and x^* is not a stationary point.