

Numerical Optimisation
Constraint optimisation:

Penalty and augmented Lagrangian methods

Assignment Project Exam Help

<https://eduassistpro.github.io>

f.rullan@cs.uc

Add WeChat edu_assist_pro

Department of Computer Science
Centre for Medical Image Computing

Centre for Inverse Problems
University College London

Lecture 14

Constraint optimization problem

Assignment Project Exam Help

$$\min_{x \in \mathbb{R}^n} f(x)$$

$$\text{subject to } f_i(x) \leq 0, \quad i = 1, \dots, m,$$

Con constraints.

Idea: Minimise a merit function $Q(x)$:
Some minimisers of $Q(x; \mu)$ approach the constraints as μ approach some set \mathcal{M} .

Benefit: reformulation as an unconstraint problem.

Consider a problem with equality constraints

$$\begin{aligned} & \min_{x \in \mathcal{D} \subset \mathbb{R}^n} f(x) \\ \text{subject to } & h_i(x) = 0, \quad i = 1, \dots, p. \end{aligned}$$

The m

<https://eduassistpro.github.io>

$$Q(x; \mu) := f(x) + \frac{\mu}{2} \|h(x)\|^2,$$

where $\mu > 0$ is the *penalty parameter*.

Idea: choose a sequence $\{\mu_k\}$: $\mu_k \rightarrow \infty$ as $k \rightarrow \infty$,
i.e. increasingly penalise the constraint, and compute the sequence
 $\{x_k\}$ of (approximate) minimisers of $Q(x; \mu_k)$.

Let $\{x_k\}$ be the sequence of approximate minimisers of $Q(x; \mu_k)$, such that $\|\nabla_x Q(x_k; \mu_k)\| \leq \tau_k$, x^* be the limit point of $\{x_k\}$ as $\tau_k \rightarrow 0$.
the sequences of the penalty parameters $\mu_k \rightarrow \infty$ and tolerances

Assignment Project Exam Help

-

- <https://eduassistpro.github.io>

$\nabla h_i(x^*)$ are linearly independent, then x^* is a KKT point for (COP:E), and we have that

Add WeChat [edu_assist_pro](https://eduassistpro.github.io)

$$\lim_{k \rightarrow \infty} \mu_k h_i(x_k) = \nu_i^*,$$

where ν^* is the multiplier vector that satisfies the KKT conditions for (COP:E).

Proof:

$$\nabla_x Q(x_k; \mu_k) = \nabla f(x_k) + \sum_{i=1}^p \mu_k h_i(x_k) \nabla h_i(x_k) \quad (1)$$

Assignment Project Exam Help

From the convergence criterium $\|\nabla_x Q(x_k; \mu_k)\| \leq \tau_k$ (using the ineq

<https://eduassistpro.github.io>

$$\|\sum_{i=1}^p \frac{\nabla h_i(x_k)}{\mu_k}\|$$

As $k \rightarrow \infty$: $\tau_k \rightarrow 0$, $\|\nabla f(x_k)\| \rightarrow \|\nabla f(x^\star)\|$

$$\sum_{i=1}^p h_i(x^\star) \nabla h_i(x^\star) = 0.$$

- i) If $h_i(x^*) \neq 0, i = 1, \dots, p$ then $\nabla h_i(x^*)$ are linearly dependent which implies that x^* is a stationary point of $\|h(x)\|^2$.
- ii) If $\nabla h_i(x^*), i = 1, \dots, p$ are linearly independent, $h_i(x^*) = 0$ and x^* is primarily feasible i.e. satisfies the second KKT condition. It remains to show that the ‘dual feasibility’ (the first KKT condition) is satisfied.

Case ii

Intu

As k

<https://eduassistpro.github.io>

$$\mathcal{L}(x^*; \nu^*) = f(x^*) + \sum_{i=1}^p \nu_i^* h_i(x^*) \quad (2)$$

Add WeChat edu_assist_pro

and $\nabla_x Q(x^k)$ its derivative i.e. the “dual feasibility” condition

$$\nabla_x \mathcal{L}(x^*; \nu^*) = \nabla f(x^*) + \sum_{i=1}^p \nu_i^* \nabla h_i(x^*). \quad (3)$$

Rearranging (1) and denoting $A(x)^T := \nabla h_i(x_k)$, $i = 1, \dots, p$ and $\nu^k := \mu_k h(x_k)$ we obtain

$$A(x_k)^T \nu^k = -\nabla f(x_k) + \nabla_x Q(x_k; \mu_k), \quad \|\nabla_x Q(x_k; \mu_k)\| \leq \tau_k.$$

For large enough k the matrix $A(x_k)$ has full row rank and hence the above overdetermined system has the unique solution

Taki <https://eduassistpro.github.io>

$\lim_{k \rightarrow \infty} \nu^k = \nu^* = -(A(x^*)^T A(x^*))^{-1} A(x^*)^T b$

and the same in (1) yields the “dual feasibility” con

$$\nabla f(x^*) + A(x^*)^T \nu^* = 0.$$

Hence, x^* is the KKT point with unique Lagrange multiplier ν^* .

Example



$$\begin{array}{ll} \min & x_1 + x_2 \\ \text{subject to} & x_1^2 + x_2^2 - 2 = 0. \end{array}$$

Solution: $(-1, -1)^T$.

Qua

$$-\frac{x_1^2 + x_2^2 - 2}{2}.$$

<https://eduassistpro.github.io>



Assignment Project Exam Help

$$\min -5x_1^2 + x_2^2$$

Solu <https://eduassistpro.github.io>

Quadratic penalty function: $Q(x; \mu) =$ $\begin{matrix} 2 & 2 & \mu & 2 \end{matrix}$.

$Q(x; \mu)$ is unbounded for $\mu < 10$.

Add WeChat edu_assist_pro

The iterates would diverge. Unfortunately, a co

III-conditioning of Hessian

Newton step: $\nabla_{xx}^2 Q(x; \mu_k) p_n = -\nabla_x Q(x; \mu_k)$

$\nabla_{xx}^2 Q(x; \mu_k) = \nabla^2 f(x) + \sum_{i=1}^p \mu_k h_i(x) \nabla^2 h_i(x) + \mu_k \nabla h(x) \nabla h(x)^T$
 $\approx \nu_i$
=: $A(x)^T$

If $x \in$

<https://eduassistpro.github.io>

As $\mu_k \rightarrow \infty$ the Hessian is dominated by the second term and hence increasingly ill-conditioned.

Add WeChat edu_assist_pro

Alternative formulation avoids ill-conditioning

$$\begin{bmatrix} \nabla^2 f(x) + \sum_{i=1}^p \mu_k h_i(x) \nabla^2 h_i(x) & A(x)^T \\ A(x) & \mu_k^{-1} I \end{bmatrix} \begin{bmatrix} p_n \\ \zeta \end{bmatrix} = \begin{bmatrix} -\nabla_x Q(x; \mu_k) \\ 0 \end{bmatrix}.$$

Still, if $\mu_k h_i(x)$ is not a good enough approximation to ν^* , inadequate quadratic model yields inadequate search direction p_n .

For general constraint problems including equality and inequality constraints, the quadratic penalty function can be defined as

<https://eduassistpro.github.io>

where $[y]^- := \max\{y, 0\}$

Note: Q may be less smooth than the objective a functions e.g. $f_1(x) = x_1 \geq 0$, then max second derivate and so does Q .

Add WeChat `edu_assist_pro`

- μ_k can be chosen adaptively based on the difficulty of minimising the penalty function in each iteration i.e. when minimising $Q(x; \mu_k)$ is expensive, choose μ_{k+1} moderately

$(x; \mu_k)$

<https://eduassistpro.github.io>

- There is no guarantee that $\|\nabla_x Q(x; \mu_k)\| - \tau_k$ will be satisfied. Practical implementations need to increase μ (and possibly recompute the initial value) until the constraint violation is not decreasing fast enough. If the iterations appear diverging.

Add WeChat edu_assist_pro

- When only equality constraints are present, $Q(x; \mu_k)$ is smooth and algorithms for unconstraint optimisation can be used, however $Q(x; \mu_k)$ becomes more difficult to minimise as μ_k becomes large unless special techniques are used. In

<https://eduassistpro.github.io>

direction due to inadequacy of quadratic approximation.

- Choice of initial point e.g. warm start
Add WeChat edu_assist_pro

Nonsmooth penalty functions

Some penalty functions are *exact* i.e. for certain choices of penalty parameters, minimisation w.r.t. x yields the exact minimiser of f .

To be exact the function has to be nonsmooth.

Assignment Project Exam Help

$$Q_1(x, \mu) := f(x) + \mu \sum_{i=1}^m |h_i(x)| + \mu \sum_{i=1}^m [h_i(x)]^+,$$

where

Let <https://eduassistpro.github.io>

order necessary conditions with Lagrange multipliers $\star \star$

x^* is a local minimiser of $Q_1(x; \mu)$ for all $\mu > 0$

If moreover the 2nd order sufficient conditions hold

then x^* is a strict local minimiser of Q_1

Let \hat{x} be a stationary point of the penalty function $Q_1(x; \mu)$ for all $\mu > \hat{\mu} > 0$. Then, if \hat{x} is feasible for (COP), it satisfies KKT conditions. If \hat{x} is not feasible for (COP), it is an infeasible stationary point.

Example revisited



$$\min \quad x_1 + x_2$$

$$\text{subject to} \quad x_1^2 + x_2^2 - 2 = 0.$$

Solution: $(-1, -1)^T$.

ℓ_1 pe

<https://eduassistpro.github.io>



Augmented Lagrangian

Reduces ill-conditioning by introducing explicit Lagrange multiplier estimates into the function to be minimised.

Can preserve smoothness. Can be implemented using standard unconstrained (or bound constrained) optimization.

Assignment Project Exam Help

Motivation: The minimisers x_k of $Q(x; \mu_k)$ do not quite satisfy the fe

<https://eduassistpro.github.io>

Obviously, in the limit $\mu_k \rightarrow \infty$, $h_i(x)$

systematic perturbation for moderate values of

Add WeChat edu_assist_pro

Augmented Lagrangian:

$$\mathcal{L}_A(x, \nu; \mu) := f(x) + \sum_{i=1}^p \nu h_i(x) + \frac{\mu}{2} \sum_{i=1}^p h_i^2(x).$$

Update of Lagrange multiplier estimate

Optimality condition for the unconstraint minimiser of
 $\mathcal{L}_A(x, \nu^k; \mu_k)$

Assignment Project Exam Help

$$0 \approx \nabla_x \mathcal{L}_A(x_k, \nu^k, \mu_k) = \nabla f(x_k) + \sum_{i=1}^p [\nu_i^k + \mu_k h_i(x_k)] \nabla h_i(x_k).$$

Opti

<https://eduassistpro.github.io>

$$0 \approx \nabla_x \mathcal{L}(x_k, \nu^*) = \nabla f(x_k) \quad *$$

Add WeChat edu_assist_pro

Comparison yields (an update scheme for

$$\nu^* \approx \nu^k + \mu_k h_i(x_k), \quad i = 1, \dots, p$$

as from $h_i(x_k) = \frac{1}{\mu_k}(\nu_i^* - \nu_i^k)$, $i = 1, \dots, p$ we see that if ν^k is close to ν^* the infeasibility goes to 0 faster than $1/\mu_k$.

Example revisited

$$\begin{aligned} & \min \quad x_1 + x_2 \\ \text{subject to} \quad & x_1^2 + x_2^2 - 2 = 0. \end{aligned}$$

Assignment Project Exam Help

Aug
 $\mathcal{L}(x,$





Figure: $\nu = 0.4$

Convergence

Let x^* be a local minimiser of (COP:E) at which the constraint gradients are linearly independent and which satisfies the 2nd order sufficient conditions with Lagrange multipliers ν^* . Then for all $u \geq \bar{U} > 0$, x^* is a strict local minimiser of $\mathcal{L}_A(x, \nu^*, h)$. Furthermore, there exist $\delta, \epsilon, M > 0$ such that for all ν^k, μ_k satis

• <https://eduassistpro.github.io>, has a

unique solution x_k and it holds

• Add WeChat `edu_assist_pr`

• it holds

$$\|\nu^{k+1} - \nu^*\| \leq M\|\nu^k - \nu^*\|/\mu_k,$$

where $\nu^{k+1} = \nu^k + \mu_k h(x_k)$.

• the matrix $\nabla_{xx}^2 \mathcal{L}_A(x_k, \nu^k; \mu_k)$ is positive definite and the constraint gradients $\nabla h_i(x_k), i = 1, \dots, p$ are linearly independent.

Assignment Project Exam Help

- Bound constraint formulation: convert inequality constraints into equality constraints using slack variables

<https://eduassistpro.github.io>

gradient algorithm

Add WeChat `edu_assist_pro`

where $P(\cdot; l, u)$ projects on the box $[l, u]$.

- **Linearly constraint formulation:** transform into equality constraint problem and linearise constraints

$\min F_k(x)$, subject to $f_i(x_k) + \nabla f_i^T(x_k)(x - x_k) = 0, \quad l \leq x \leq u.$

Choose F_k as

Assignment Project Exam Help

$$F_k(x) = f(x) + \sum_{i=1}^m \nu_i^k \bar{f}_i^k(x),$$

<https://eduassistpro.github.io>

$i \quad i \quad i \quad k \quad i \quad k \quad k$

Preferred choice (larger convergence radius)

Add WeChat edu_assist_pro

$$F_k(x) = f(x) + \sum_{i=1}^m \nu_i^k \bar{f}_i^k \quad \bar{2} \quad i$$

- **Unconstraint formulation:** obtain unconstraint formulation using smooth approximation to feasibility set indicator function.