

# Assignment Project Exam Help

Machine learning lecture slides

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

Regression I: Linear regression

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

- ▶ Statistical model for regression
- ▶ College GPA example
- ▶ Ordinary least squares for linear regression
- ▶ The expected mean squared error

Assignment Project Exam Help

- ▶
- ▶
- ▶
- ▶

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

Figure 1: Galton board

## Real-valued predictions

- ▶ Example: Galton board
- ▶ Physical model: hard
- ▶ Statistical model: final position of ball is random
  - ▶ Normal (Gaussian) distribution with mean  $\mu$  and variance  $\sigma^2$

<https://eduassistpro.github.io>

- ▶ Goal: predict final position accurately, i.e.  
(also called squared error)

$$(\text{prediction} - \text{outcome})^2$$

- ▶ Outcome is random, so look at expected squared loss (also called mean squared error)

## Optimal prediction for mean squared error

- ▶ Predict  $\hat{y} \in \mathbb{R}$ ; true final position is  $Y$  (random variable) with mean  $\mathbb{E}(Y) = \mu$  and variance  $\text{var}(Y) = \mathbb{E}[(Y - \mathbb{E}(Y))^2] = \sigma^2$ .
- ▶ Square error is  $(\hat{y} - Y)^2$ .
- ▶ Bias-variance decomposition:

<https://eduassistpro.github.io>

$$y - \mu \quad \sigma^2.$$

Add WeChat [edu\\_assist\\_pro](#)

- ▶ This is true for any random variable assumption.
- ▶ So optimal prediction is  $\hat{y} = \mu$ .
- ▶ When parameters are unknown, can estimate from related data, ...
- ▶ Can also do an analysis of a plug-in prediction ...

## Statistical model for regression

- ▶ Setting is same as for classification except:
  - ▶ Label is real number, rather than  $\{0, 1\}$  or  $\{1, 2, \dots, K\}$
  - ▶ Care about squared loss, rather than whether prediction is correct
  - ▶ Mean squared error of  $f$ :

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Optimal prediction function for regression

- If  $(X, Y)$  is random test example, then

optimal prediction function is

Assignment Project Exam Help

$$f^*(x) = \mathbb{E}[Y \mid X = x]$$

function  
<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr



## Test MSE (1)

- ▶ Just like in classification, we can use test data to estimate  $\text{mse}(\hat{f})$  for a function  $\hat{f}$  that depends only on training data.

▶ IID model

$(X_1, Y_1), \dots, (X_n, Y_n), (X'_1, Y'_1), \dots, (X'_m, Y'_m), (X, Y)$  are iid

<https://eduassistpro.github.io>

- ▶ Predictor  $\hat{f}$  is based only on training exa
- ▶ Hence, **test examples are independent** (important!)
- ▶ We would like to estimate  $\text{mse}(\hat{f})$

## Test MSE (2)

- ▶ Test MSE  $\text{mse}(\hat{f}, T) = \frac{1}{m} \sum_{i=1}^m (\hat{f}(X'_i) - Y'_i)^2$
- ▶ By law of large numbers,  $\text{mse}(\hat{f}, T) \rightarrow \text{mse}(\hat{f})$  as  $m \rightarrow \infty$

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Example: College GPA

- ▶ Data from 750 Dartmouth students' College GPA
  - ▶ Mean: 2.46
  - ▶ Standard deviation: 0.746
- ▶ Assume this data is iid sample from the population of Dartmouth students (false)

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Predicting College GPA from HS GPA (1)

- ▶ Students represented in data have High School (HS) GPA
  - ▶ Maybe HS GPA is predictive of College GPA?

▶ Data:  $S := ((x_1, y_1) \dots, (x_n, y_n))$   
▶  $x_i$  is HS GPA of  $i$ -th student

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr




Figure 3: Plot of College GPA vs HS G

# Predicting College GPA from HS GPA (2)

- ▶ First attempt:

- ▶ Define intervals of possible HS GPAs:

$(0.00, 0.25]$ ,  $(0.25, 0.50]$ ,  $(0.50, 0.75]$ , ...

ge

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pro

$$\hat{f}(x) := \begin{cases} \hat{\mu}_{(0.25, 0.50]} \\ \hat{\mu}_{(0.50, 0.75]} \end{cases}$$

- ▶ (What to do about an interval  $I$  that contains no student's HS GPA?)




Figure 4: Plot of mean College GPA vs binne

## Predicting College GPA from HS GPA (3)

- Define

$$\text{mse}(f, S) := \frac{1}{|S|} \sum_{(x, y) \in S} (f(x) - y)^2,$$

the mean squared error of predictions made by  $f$  on examples

<https://eduassistpro.github.io>

$$\text{mse}(\hat{f}, S) = 0.376$$

$\sqrt{\text{mse}(\hat{f}, S)} = 0.613 < 0.710$  (Add WeChat [edu\\_assist\\_pro](#))

- Piece-wise constant function  $\hat{f}$  is an improvement over the constant function (i.e., just predicting the mean 2.46 for all  $x$ )!



## Predicting College GPA from HS GPA (4)

- ▶ But  $\hat{f}$  has some quirks.
- ▶ E.g., those with HS GPA between 2.50 and 2.75 are predicted to have a lower College GPA than those with HS GPA between 2.25 and 2.50.
- ▶ E.g., something unusual with the student who has HS GPA of




Figure 5: Plot of mean College GPA vs binned HS GPA

## Least squares linear regression (1)

- Suppose we'd like to only consider functions with a specific functional form, e.g., a linear function:

Assignment Project Exam Help

$$f(x) = mx + \theta$$

<https://eduassistpro.github.io>

$m$

prediction of College GPA.

Add WeChat edu\_assist\_pr

## Least squares linear regression (2)

- ▶ What is the linear function with smallest MSE on  $(x_1, y_1), \dots, (x_n, y_n) \in \mathbb{R} \times \mathbb{R}$ ? This is the problem of

least squares linear regression

Assignment Project Exam Help

- ▶ Find  $(m, \theta) \in \mathbb{R}^2$  to minimize

<https://eduassistpro.github.io>

- ▶ Also called ordinary least squares \_\_\_\_

Add WeChat edu\_assist\_pr




Figure 6: Plot of least squares linear regression

# Computing OLS (1)

- Derivatives equal zero conditions (normal equations):

$$\frac{\partial}{\partial \theta} \left\{ \frac{1}{n} \sum_{i=1}^n (mx_i + \theta - y_i)^2 \right\} = \frac{2}{n} \sum_{i=1}^n (mx_i + \theta - y_i) = 0$$

$$\frac{\partial}{\partial m} \left\{ \frac{1}{n} \sum_{i=1}^n (mx_i + \theta - y_i)^2 \right\} = \frac{2}{n} \sum_{i=1}^n x_i (mx_i + \theta - y_i) = 0.$$

- Define

$$\bar{x} := \frac{1}{n} \sum_{i=1}^n x_i, \quad \bar{y} := \frac{1}{n} \sum_{i=1}^n y_i$$

$$\overline{xy} := \frac{1}{n} \sum_{i=1}^n x_i y_i, \quad \bar{y} := \frac{1}{n} \sum_{i=1}^n y_i$$

so system can be re-written as

$$\bar{x}m + \theta = \bar{y}$$

$$\bar{x}^2 m + \bar{x} \theta = \overline{xy}.$$

- Write in matrix notation:

Assignment  $\begin{bmatrix} 1 & 1 \\ \frac{x}{2} & x \end{bmatrix} \begin{bmatrix} \hat{m} \\ \hat{\theta} \end{bmatrix} = \begin{bmatrix} \bar{y} \\ \overline{xy} \end{bmatrix}$  Exam Help

<https://eduassistpro.github.io>

- Solution:  $(\hat{m}, \hat{\theta}) \in \mathbb{R}^2$  given by

Add WeChat edu\_assist\_pro

$$\hat{m} := \frac{\overline{xy} - \bar{x} \cdot \bar{y}}{\overline{x^2} - \bar{x}^2}, \quad \hat{\theta} := - \frac{\overline{xy} - \bar{x} \cdot \bar{y}}{\overline{x^2} - \bar{x}^2}$$

## Computing OLS (3)

- ▶ Catch: The above solution only makes sense if  $\overline{x^2} - \bar{x}^2 \neq 0$ , i.e., the variance of the  $x_i$ 's is non-zero.

Assignment Project Exam Help

<https://eduassistpro.github.io>

- ▶ If  $\overline{x^2} - \bar{x}^2 = 0$ , then the matrix defining the equations is singular.

Add WeChat edu\_assist\_pro

- ▶ In general, “derivative equals zero” is only a necessary condition for a solution to be optimal; not necessarily a sufficient condition!

Assignment Project Exam Help

<https://eduassistpro.github.io>

- ▶ **Theorem** Every solution to the normal eq is an optimal solution to the least squares linear r

Add WeChat: edu\_assist\_pro



## Decomposition of expected MSE (1)

- ▶ Two different functions of HS GPA for predicting College GPA.
  - ▶ What makes them different?
  - ▶ We care about prediction of College GPA for student we haven't seen before based on their HS GPA.

Assignment Project Exam Help

- ▶
- ▶
- ▶

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Decomposition of expected MSE (2)

Assignment Project Exam Help

$$\begin{aligned}\mathbb{E}[\text{mse}(\hat{f})] &= \mathbb{E}[\mathbb{E}[(f(X) - Y)^2 \mid \hat{f}]] \\ &= \mathbb{E}[\mathbb{E}[(f(X) - \mathbb{E}[Y \mid X])^2 \mid \hat{f}]] \\ &= \mathbb{E}[\text{var}(Y \mid X) + \mathbb{E}[(f(X) - \mathbb{E}[Y \mid X])^2 \mid \hat{f}]] \\ &= \mathbb{E}[\text{var}(Y \mid X) + \text{var}(\hat{f}(X) \mid X) + \mathbb{E}[(f(X) - \mathbb{E}[Y \mid X] - \hat{f}(X) + \hat{f}(X))^2 \mid X]] \\ &= \underbrace{\mathbb{E}[\text{var}(Y \mid X)]}_{\text{unavoidable error}} + \underbrace{\mathbb{E}[\text{var}(\hat{f}(X) \mid X)]}_{\text{variability of } \hat{f}} + \underbrace{\mathbb{E}[\mathbb{E}[(f(X) - \mathbb{E}[Y \mid X] - \hat{f}(X) + \hat{f}(X))^2 \mid X]]}_{\text{approximation error of } \hat{f}}\end{aligned}$$

<https://eduassistpro.github.io>

Add WeChat: edu\_assist\_pro

## Decomposition of expected MSE (3)

- ▶ First term quantifies inherent unpredictability of  $Y$  (even after seeing  $X$ )
- ▶ Second term measures the “variability” of  $\hat{f}$  due to the random nature of training data. Depends on:

<https://eduassistpro.github.io>

- ▶ Third term quantifies how well a function  $p$  fitting procedure can approximate the reg after removing the “variability” of

## Multivariate linear regression (1)

- ▶ For Dartmouth data, also have SAT Score for all students.
  - ▶ Can we use both predictor variables (HS GPA and SAT Score) to get an even better prediction of College GPA?
  - ▶ Binning approach: instead of a 1-D grid (intervals), consider a 2-D grid (squares).

<https://eduassistpro.github.io>

for some  $(m_1, m_2) \in \mathbb{R}^2$  and  $\theta$

Add WeChat edu\_assist\_pr

## Multivariate linear regression (2)

- The general case: a (homogeneous) linear function  $f: \mathbb{R}^d \rightarrow \mathbb{R}$  of the form

Assignment Project Exam Help  
 $f(x) = x^T w$   
for some  $w \in \mathbb{R}^d$ .

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Multivariate ordinary least squares (1)

- What is the linear function with smallest MSE on  $(x_1, y_1), \dots, (x_n, y_n) \in \mathbb{R}^d \times \mathbb{R}$ ?

► Find  $w \in \mathbb{R}^d$  to minimize

$$\frac{1}{n} \sum_{i=1}^n (w^\top x_i - y_i)^2$$

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Multivariate ordinary least squares (2)

- In matrix notation:

Assignment Project Exam Help

where

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pro

- If we put vector  $v \in \mathbb{R}^d$  in the context, it is treated as a column vector by default!
- If we want a row vector, we write

- Therefore

$$Aw - b = \frac{1}{\sqrt{n}} \begin{bmatrix} x_1^\top w - y_1 \\ \vdots \\ x_n^\top w - y_n \end{bmatrix}$$

# Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pro

Figure 7: Geometric picture of least squares linear regression



## Multivariate normal equations (1)

- ▶ Like the one-dimensional case, optimal solutions are characterized by a system of linear equations (the “derivatives equal zero” conditions) called the normal equations:

$$\frac{\partial}{\partial w_d} (w)$$

$$\frac{\partial}{\partial w_d} \left( \sum_{i=1}^n (y_i - \hat{y}_i)^2 \right)$$

which is equivalent to

$$A^T A w =$$

## Multivariate normal equations (2)

- ▶ If  $A^T A$  is non-singular (i.e., invertible), then there is a unique solution given by

Assignment Project Exam Help

$$\hat{w} := (A^T A)^{-1} A^T b.$$



<https://eduassistpro.github.io>



Add WeChat edu\_assist\_pr

**Theorem:** Every solution to the normal eq  
optimal solution to the least squares linear r

## Algorithm for least squares linear regression

- ▶ How to solve least squares linear regression problem?
  - ▶ Just solve the normal equations, a system of  $d$  linear equations in  $d$  unknowns.
  - ▶ Time complexity (naïve) of Gaussian elimination algorithm:  $O(d^3)$ .

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

# Classical statistics view of OLS (1)

- ▶ Normal linear regression model

- ▶ Model training examples  $(X_1, Y_1), \dots, (X_n, Y_n)$  as iid random variables taking values in  $\mathbb{R}^d \times \mathbb{R}$  where

$$Y \mid X = x \sim \mathcal{N}(x^\top w, \sigma^2)$$

- ▶ <https://eduassistpro.github.io>  
problem of finding the maximum likelihood

Add WeChat edu\_assist\_pr

## Classical statistics view of OLS (2)

- ▶ Suppose your data really does come from a distribution in this statistical model, say, with parameters  $w$  and  $\sigma^2$ .

Assignment Project Exam Help

- ▶ Then the function with smallest MSE is the linear function  $f^*(x) = x^T w$ , and its MSE is  $\text{mse}(f^*) = \sigma^2$ .

- ▶ So estimating  $w$  is a sensible idea! (Plug-in principle...)

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Statistical learning view of OLS (1)

- ▶ IID model:  $(X_1, Y_1), \dots, (X_n, Y_n), (X, Y) \sim_{\text{iid}} P$  are iid random variables taking values in  $\mathbb{R}^d \times \mathbb{R}$ 
  - ▶  $(X, Y)$  is the (unlabeled) 'test' example
- ▶ Goal: find a (linear) function  $w \in \mathbb{R}^d$  with small MSE

- ▶ <https://eduassistpro.github.io>  
 $w \in \mathbb{R}^d$ ,  
since it is an expectation (e.g., integral) with respect to unknown distribution  $P$

Add WeChat edu\_assist\_pro

## Statistical learning view of OLS (2)

- ▶ However, we have an iid sample  $S := ((X_1, Y_1), \dots, (X_n, Y_n))$ .
- ▶ We swap out  $P$  in the definition of  $\text{mse}(f)$ , and replace it with the empirical distribution  $P_n$ .

Assignment Project Exam Help

<https://eduassistpro.github.io> on the

$i$ -th training example.

- ▶ Resulting objective function is

Add WeChat edu\_assist\_pro

$$\mathbb{E}[(\tilde{X}^\top w - \tilde{Y})^2] = \frac{1}{n} \sum_{i=1}^n (X_i^\top w - Y_i)^2$$

where  $(\tilde{X}, \tilde{Y}) \sim P_n$ .

## Statistical learning view of OLS (3)

- ▶ In some circles:

- ▶ (True/population) risk of  $w$ :  $\mathcal{R}(w) := \mathbb{E}[(X^\top w - Y)^2]$

- ▶ Empirical risk of  $w$ :  $\hat{\mathcal{R}}(w) := \frac{1}{n} \sum_{i=1}^n (X_i^\top w - Y_i)^2$

- ▶ This is another instance of the plug-in principle!

e

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr



## Statistical learning view of OLS (4)

- ▶ This is not specific to linear regression; also works for other types of functions, and also other types of prediction problems, including classification.
- ▶ For classification:

<https://eduassistpro.github.io>

- ▶ Procedure that minimizes empirical risk:

*Empirical risk minimization (ERM)*

Add WeChat edu\_assist\_pro

## Upgrading linear regression (1)

- ▶ Make linear regression more powerful by being creative about features

Assignment Project Exam Help

- ▶ We are forced to do this if  $x$  is not already provided as a vector of numbers

- ▶ ion  $\varphi$

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Upgrading linear regression (2)

- ▶ Examples:

- ▶ Affine feature expansion, e.g.,  $\varphi(x) = (1, x)$ , to accommodate intercept

- ▶ Standardization, e.g.,  $\varphi(x) = (x - \mu)/\sigma$  where  $(\mu, \sigma^2)$  are (estimates of) the mean and variance of the feature value  $x$

<https://eduassistpro.github.io>

- ▶ Polynomial expansion, e.g.,

- $\varphi(x) = (1, x_1, \dots, x_d, x_1^2, \dots, x_d^2)$

- ▶ Headless neural network,  $\varphi(x) =$

- $N: \mathbb{R}^d \rightarrow \mathbb{R}^k$  is a map computed by a in neural network

- ▶ (Later, we'll talk about how to “learn”  $N$ .)

## Example: Taking advantage of linearity

- ▶ Example:  $y$  is health outcome,  $x$  is body temperature
  - ▶ Physician suggests relevant feature is (square) deviation from normal body temperature  $(x - 98.6)^2$
  - ▶ What if you didn't know the magic constant 98.6? (Apparently it is wrong in the US anyway)

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Example: Binning features

- ▶ Dartmouth data example, where we considered intervals for the HS GPA variable:

**Assignment Project Exam Help**  
 $(0.00, 0.25], (0.25, 0.50], (0.50, 0.75], \dots$

- ▶  $\text{ear}$
- ▶ <https://eduassistpro.github.io>
- ▶  $\varphi(x)^\top w = w_j$  if  $x$  is in the  $j$ -th i

**Add WeChat edu\_assist\_pr**

## Effect of feature expansion on expected MSE

$$\mathbb{E}[\text{mse}(\hat{f})] = \underbrace{\mathbb{E}[\text{var}(Y | X)]}_u + \underbrace{\mathbb{E}[\text{var}(\hat{f}(X) | X)]}_{\text{approximation error}} + \underbrace{\mathbb{E}[(\mathbb{E}[\hat{f}(X) | X] - \mathbb{E}[Y | X])^2]}_{\text{error of } \hat{f}}$$

- ▶ <https://eduassistpro.github.io>  
(approximation error)
- ▶ But maybe at the cost of increasing the second (variability)

## Performance of OLS (1)

- ▶ Study in context of IID model
- ▶  $(X_1, Y_1), \dots, (X_n, Y_n), (X, Y)$  are iid, and assume  $\mathbb{E}[XX^T]$  is invertible (WLOG).
- ▶ Let  $w^*$  denote the minimizer of  $\text{mse}(w)$  over all  $w \in \mathbb{R}^a$ .

Assignment Project Exam Help

<https://eduassistpro.github.io>

- ▶ How much larger is  $\text{mse}(\hat{w})$  comp

Add WeChat edu\_assist\_pr

## Performance of OLS (2)

- **Theorem:** In the IID model, the OLS solution  $\hat{w}$  satisfies

**Assignment Project Exam Help**

as  $n \rightarrow \infty$ , where  $W = \mathbb{E}[XX^\top]^{-1/2}X$  and  $\varepsilon = Y - X^\top w^*$ .

- <https://eduassistpro.github.io>

**Add WeChat edu\_assist\_pro**

which is more typically written as

$$\mathbb{E}[\text{mse}(\hat{w})] \rightarrow \left(1 + \frac{d}{n}\right) \text{mse}(w^*).$$



- ▶ Write  $A = \begin{bmatrix} \uparrow & & \uparrow \\ a_1 & \cdots & a_d \\ \downarrow & & \downarrow \end{bmatrix}$ 
  - ▶  $a_j \in \mathbb{R}^n$  is  $j$ -th column of  $A$
  - ▶ Span of  $a_1, \dots, a_d$  is  $\text{range}(A)$ , a subspace of  $\mathbb{R}^n$

Assignment Project Exam Help

▶ <https://eduassistpro.github.io>ding

Add WeChat edu\_assist\_pr




Figure 8: Orthogonal projection of  $b$  onto  $\text{range}(A)$

## Linear algebraic view of OLS (2)

- ▶ Solution  $\hat{b}$  is orthogonal projection of  $b$  onto  $\text{range}(A)$

- ▶  $\hat{b}$  is unique

- ▶ Residual  $b - \hat{b}$  is orthogonal to  $\hat{b}$

- ▶ To get  $w$  from  $\hat{b}$ , solve  $Aw = \hat{b}$  for  $w$ .

- ▶ If  $\text{rank}(A) < d$  (always the case if  $n < d$ ), then infinitely-many

- ▶ <https://eduassistpro.github.io>  $n \geq d$ ,

Add WeChat edu\_assist\_pr

## Over-fitting (1)

- In the IID model, over-fitting is the phenomenon where the true risk is much worse than the empirical risk.

Assignment Project Exam Help

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pr

## Over-fitting (2)

- ▶ Example:

- ▶  $\varphi(x) = (1, x, x^2, \dots, x^k)$ , degree- $k$  polynomial expansion

- ▶ Dimension is  $d = k + 1$

- ▶ Any function of  $\leq k + 1$  points can be interpolated by polynomial of degree  $\leq k$

<https://eduassistpro.github.io>

Add WeChat edu\_assist\_pro




Figure 9: Polynomial interpolation

- ▶ Recall plug-in principle
  - ▶ Want to minimize risk with respect to (unavailable)  $P$ ; use  $P_n$  instead
- ▶ What if we can't regard data as iid from  $P$ ?

$\frac{1}{n}$   $\frac{1}{n}$

<https://eduassistpro.github.io>

- ▶ How to implement plug-in principle?

Add WeChat edu\_assist\_pr