

Supplementary materials for ‘Unusually warm winter seasons may compromise the performance of current phenology models - Predicting bloom dates in young apple trees with **PhenoFlex**’

Eduardo Fernandez^{a*}, Katja Schiffers^a, Carsten Urbach^b and Eike Luedeling^a

^a Department of Horticultural Science, Institute of Crop Science and Resource Conservation (INRES), Auf dem Hügel 6, D-53121 Bonn, Germany

^b Helmholtz Institut für Strahlen- und Kernphysik (HISKP), Nußalle 14-16, D-53115 Bonn, Germany

Introduction

Phenology models are crucial tools for assessing the impacts of climate change on forestry, ecology and agriculture. In the article ‘Unusually warm winter seasons may compromise the performance of current phenology models - Predicting bloom dates in young apple trees with **PhenoFlex**’, we assessed the performance of the recently proposed **PhenoFlex** phenology modelling framework (Luedeling et al. 2021) in response to different characteristics of the calibration data. In this document, we provide supplementary materials to the main document. Among these, we highlight the specific set of parameters used in each of the ten model runs as well as the results of our supplementary validation of one of the versions of the framework used in the study (i.e. Phenoflex_{excluded}).

Detailed description of the experimental design

In this study, we used a subset of phenology and weather records reported on in an earlier study by Fernandez et al. (2021). Here, we explain the experimental design and show environmental conditions used in the study. During two consecutive winters, we exposed young apple trees to several environments at different times (Fig. S1).



Figure S1: Conditions used in the experiment during the first year

By frequently transferring the trees among environments, we were able to generate 59 experimental seasons and evaluate the phenology of the trees after the exposure to these conditions (Fig. S2).

*Corresponding author: Eduardo Fernández (efernand@uni-bonn.de)





Figure S2: Experimental seasons used in the study. The first 5 experimental seasons represent the marginal experimental seasons excluded in the respective version of the analysis. In the Environment legend, CKA stands for Campus Klein-Altendorf and ETFE stands for ethylene-tetrafluoroethylene copolymer

The experimental setup allowed us to generate a substantial temperature variation during the two winter seasons as well as conditions that may have been marginal for apple trees to overcome dormancy as discussed in the main manuscript. Most of the time over the experiment, the mean temperature in those 5 marginal seasons was in the upper part of the temperature range identified among experimental seasons (Fig. S3).



Parameters used in each model run

The `phenologyFitter()` function from the `chillR` package (Luedeling 2021), which is needed to fit the 12 model parameters to data, requires an estimated set of parameters (i.e. `par.guess`) as well as the `lower` and `upper` bounds for these values. We initialized the model fitting procedure with a common set of parameters for both versions of the analysis (i.e. `PhenoFlexall` and `PhenoFlexexcluded`).

First model run

```
# Set the initial parameters (wide ranges)
# Parameter <- c(yc, zc, s1, Tu, EO, E1, A0, A1, Tf, Tc, Tb, slope)
lower      <- c(20, 100, 0.1, 0, 3000.0, 9000.0, 6000.0, 5.00e+13, 0, 0, 0, 0.05)
par.guess   <- c(40, 190, 0.5, 25, 3372.8, 9900.3, 6319.5, 5.94e+13, 4, 36, 4, 1.60)
upper      <- c(80, 500, 1.0, 30, 4000.0, 10000.0, 7000.0, 6.00e+13, 10, 40, 10, 50.00)
```

We introduced these parameters to the fitter function for each version of the analysis. Note that the specific values for the parameters and all scripts developed in this study are deposited in a public online repository (https://github.com/EduardoFernandezC/phenoflex_exp_data). The first fitting procedure resulted in `RMSE` values of 7.2 days for `PhenoFlexall` and 4.13 days for `PhenoFlexexcluded`. In the second round of the fitting procedure, we

adjusted the parameter inputs.

Second model run

```
# Set the parameters using the results from the previous run

# Version PhenoFlex_all (pheno_fit_v1_r1$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v1_r2 <- c(45, 290, 0.00, 18.0, 3000.0, 9000.0, 6000.0, 5.20e+13, 0.0, 29.0, 0.0, 0.05)
par_v1_r2 <- c(59, 326, 0.16, 22.5, 3309.9, 9900.1, 6295.7, 5.94e+13, 6.1, 39.9, 8.6, 14.60)
upper_v1_r2 <- c(65, 380, 0.50, 30.0, 4000.0, 10000.0, 7000.0, 6.20e+13, 10.0, 44.0, 12.0, 30.00)

# Same for version PhenoFlex_excluded (pheno_fit_v2_r1$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v2_r2 <- c(25, 340, 0.00, 17.0, 3000.0, 9000.0, 6000.0, 5.20e+13, 0.0, 30.0, 0.0, 1.00)
par_v2_r2 <- c(35, 387, 0.03, 21.8, 3371.0, 9900.8, 6283.6, 5.94e+13, 5.1, 40.0, 6.3, 40.00)
upper_v2_r2 <- c(40, 420, 0.50, 32.0, 4000.0, 10000.0, 7000.0, 6.20e+13, 10.0, 42.0, 10.0, 50.00)
```

In the second round of the fitting procedure, PhenoFlex_{all} reached an RMSE of 6.94 days, whereas PhenoFlex_{excluded} reached an RMSE of 3.91 days. In the third round of the fitting procedure, we adjusted the parameter inputs as follow:

Third model run

```
# Set the parameters using the results from the previous run

# Version PhenoFlex_all (pheno_fit_v1_r2$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v1_r3 <- c(48, 290, 0.00, 15.0, 3000.0, 9000.0, 6000.0, 5.20e+13, 0.0, 24.0, 0.0, 0.05)
par_v1_r3 <- c(58, 332, 0.39, 22.1, 3310.5, 9900.3, 6344.8, 5.94e+13, 7.2, 35.0, 8.8, 18.20)
upper_v1_r3 <- c(68, 370, 0.65, 32.0, 4000.0, 10500.0, 7000.0, 6.20e+13, 11.0, 44.0, 11.0, 35.00)

# Same for PhenoFlex_excluded (pheno_fit_v2_r2$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v2_r3 <- c(20, 350, 0.00, 18.0, 3000.0, 9000.0, 6000.0, 5.20e+13, 0.0, 30.0, 0.0, 1.00)
par_v2_r3 <- c(34, 406, 0.20, 21.4, 3371.4, 9901.3, 6214.0, 5.94e+13, 1.5, 42.0, 6.1, 10.70)
upper_v2_r3 <- c(50, 450, 0.55, 35.0, 4000.0, 10500.0, 7000.0, 6.20e+13, 10.0, 46.0, 10.0, 35.00)
```

After the third round of the fitting procedure, RMSE values decreased to 6.63 days for PhenoFlex_{all} and 3.91 days for PhenoFlex_{excluded}. In the fourth round of the fitting procedure, we used the following set of parameters:

Fourth model run

```
# Set the parameters using the results from the previous run

# Version PhenoFlex_all (pheno_fit_v1_r3$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v1_r4 <- c(45, 290, 0.00, 15.0, 3000.0, 9000.0, 5500.0, 5.20e+13, 0.0, 20.0, 0.0, 1.00)
par_v1_r4 <- c(60, 339, 0.54, 23.0, 3310.3, 9901.5, 6346.3, 5.94e+13, 7.2, 30.7, 7.6, 7.30)
upper_v1_r4 <- c(70, 370, 0.75, 32.0, 4000.0, 10300.0, 7000.0, 6.20e+13, 12.0, 40.0, 12.0, 20.00)

# Same for version 2 (pheno_fit_v2_r3$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v2_r4 <- c(28, 360, 0.00, 17.0, 3000.0, 9300.0, 5800.0, 5.30e+13, 0.0, 35.0, 0.0, 10.00)
par_v2_r4 <- c(34, 396, 0.20, 21.6, 3371.0, 9901.3, 6214.0, 5.94e+13, 1.5, 46.0, 5.9, 11.12)
upper_v2_r4 <- c(44, 440, 0.45, 30.0, 4000.0, 10500.0, 6800.0, 6.50e+13, 10.0, 48.0, 12.0, 40.00)
```

In this round of the fitting procedure, PhenoFlex_{all} reached an RMSE of 6.59 days, whereas PhenoFlex_{excluded} reached an RMSE of 3.65 days.

Fifth model run

```
# Set the parameters using the results from the previous run
# Version PhenoFlex_all (pheno_fit_v1_r4$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v1_r5 <- c(50, 330, 0.00, 18.0, 3000.0, 9600.0, 6000.0, 5.50e+13, 2.0, 30.0, 2.0, 0.50)
par_v1_r5 <- c(60, 354, 0.47, 22.8, 3310.3, 9901.5, 6344.3, 5.94e+13, 7.2, 36.0, 7.6, 7.63)
upper_v1_r5 <- c(75, 370, 0.55, 28.0, 4000.0, 10300.0, 7000.0, 6.50e+13, 10.0, 42.0, 12.0, 15.00)

# Same for version PhenoFlex (pheno_fit_v2_r4$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v2_r5 <- c(27, 360, 0.00, 18.0, 3000.0, 9500.0, 5800.0, 5.50e+13, 0.0, 35.0, 1.0, 1.00)
par_v2_r5 <- c(34, 383, 0.21, 22.5, 3371.0, 9901.3, 6213.9, 5.94e+13, 1.5, 45.6, 5.9, 1.08)
upper_v2_r5 <- c(37, 400, 0.55, 32.0, 4000.0, 10300.0, 6800.0, 6.30e+13, 8.0, 48.0, 10.0, 25.00)
```

In this round of the fitting procedure, PhenoFlex_{all} reached an RMSE of 6.57 days, whereas PhenoFlex_{excluded} reached an RMSE of 3.56 days.

Sixth model run

```
# Set the parameters using the results from the previous run
# Version PhenoFlex_all (pheno_fit_v1_r5$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v1_r6 <- c(40, 290, 0.00, 18.0, 2900.0, 9000.0, 5700.0, 5.30e+13, 0.0, 20.0, 0.0, 0.01)
par_v1_r6 <- c(60, 343, 0.32, 25.1, 3310.3, 9901.6, 6342.9, 5.94e+13, 5.6, 33.6, 7.0, 2.24)
upper_v1_r6 <- c(80, 390, 0.55, 32.0, 3900.0, 10500.0, 6900.0, 6.80e+13, 12.0, 48.0, 12.0, 20.00)

# Same for version PhenoFlex_excluded (pheno_fit_v2_r5$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v2_r6 <- c(25, 330, 0.00, 18.0, 3000.0, 9200.0, 5700.0, 5.40e+13, 0.05, 40.0, 0.0, 0.50)
par_v2_r6 <- c(35, 361, 0.15, 22.4, 3371.0, 9901.3, 6213.8, 5.94e+13, 1.45, 48.0, 5.9, 1.10)
upper_v2_r6 <- c(40, 390, 0.35, 30.0, 4000.0, 10800.0, 6900.0, 6.60e+13, 10.00, 50.0, 10.0, 20.00)
```

In this round of the fitting procedure, PhenoFlex_{all} reached an RMSE of 6.41 days, whereas PhenoFlex_{excluded} reached an RMSE of 3.55 days.

Seventh model run

```
# Set the parameters using the results from the previous run
# Version PhenoFlex_all (pheno_fit_v1_r6$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v1_r7 <- c(55, 325, 0.00, 20.0, 3000.0, 9000.0, 5500.0, 5.50e+13, 0.0, 30.0, 2.0, 0.00)
par_v1_r7 <- c(61, 334, 0.53, 25.0, 3310.3, 9901.6, 6342.7, 5.94e+13, 6.4, 33.9, 7.0, 1.57)
upper_v1_r7 <- c(65, 345, 0.65, 30.0, 4000.0, 10500.0, 6600.0, 6.50e+13, 10.0, 40.0, 10.0, 10.00)

# Same for version PhenoFlex_excluded (pheno_fit_v2_r6$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v2_r7 <- c(30, 365, 0.000, 20.0, 3000.0, 9500.0, 5800.0, 5.50e+13, 0.0, 35.0, 0.0, 1.00)
par_v2_r7 <- c(34, 380, 0.160, 26.4, 3371.0, 9901.3, 6214.2, 5.94e+13, 1.3, 41.2, 3.3, 1.12)
upper_v2_r7 <- c(40, 385, 0.225, 30.0, 4000.0, 10500.0, 6800.0, 6.40e+13, 6.0, 45.0, 8.0, 20.00)
```

In this round of the fitting procedure, PhenoFlex_{all} reached an RMSE of 6.05 days, whereas PhenoFlex_{excluded} reached an RMSE of 3.46 days.

Eighth model run

```
# Set the parameters using the results from the previous run
# Version PhenoFlex_all (pheno_fit_v1_r7$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v1_r8 <- c(55, 315, 0.30, 22.0, 2800.0, 9500.0, 5900.0, 5.60e+13, 2.0, 26.0, 2.0, 0.00)
par_v1_r8 <- c(62, 336, 0.52, 27.1, 3310.3, 9901.7, 6395.4, 5.94e+13, 6.5, 30.2, 5.6, 1.39)
upper_v1_r8 <- c(70, 355, 0.70, 32.0, 3800.0, 10200.0, 6700.0, 6.40e+13, 9.0, 36.0, 8.0, 15.00)

# Same for version PhenoFlex_excluded (pheno_fit_v2_r7$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v2_r8 <- c(28, 345, 0.01, 20.0, 2800.0, 9600.0, 5800.0, 5.60e+13, 0.0, 38.0, 0.0, 7.00)
par_v2_r8 <- c(35, 375, 0.13, 26.4, 3371.0, 9901.3, 6214.4, 5.94e+13, 1.3, 44.7, 3.0, 1.80)
upper_v2_r8 <- c(40, 405, 0.30, 30.2, 3700.0, 10300.0, 6800.0, 6.30e+13, 8.0, 52.0, 8.0, 30.00)
```

In this round of the fitting procedure, PhenoFlex_{all} reached an RMSE of 5.96 days, whereas PhenoFlex_{excluded} reached an RMSE of 3.45 days.

Ninth model run

```
# Set the parameters using the results from the previous run
# Version PhenoFlex_all (pheno_fit_v1_r8$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v1_r9 <- c(55, 300, 0.50, 20.0, 3000.0, 9500.0, 5500.0, 5.00e+13, 1.0, 20.0, 2.0, 0.00)
par_v1_r9 <- c(63, 315, 0.70, 27.9, 3310.3, 9901.7, 6396.2, 5.94e+13, 6.5, 27.9, 5.7, 1.39)
upper_v1_r9 <- c(75, 325, 0.85, 35.0, 4000.0, 10500.0, 6900.0, 7.00e+13, 11.0, 37.0, 10.0, 25.00)

# Same for version PhenoFlex_excluded (pheno_fit_v2_r8$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v2_r9 <- c(23, 325, 0.00, 16.0, 3000.0, 9500.0, 5700.0, 5.60e+13, 0.0, 40.0, 0.0, 1.00)
par_v2_r9 <- c(34, 357, 0.14, 26.8, 3371.0, 9901.2, 6214.1, 5.94e+13, 1.4, 51.3, 3.6, 8.45)
upper_v2_r9 <- c(43, 375, 0.50, 36.0, 3800.0, 10500.0, 6700.0, 6.30e+13, 10.0, 55.0, 10.0, 20.00)
```

In this round of the fitting procedure, PhenoFlex_{all} reached an RMSE of 5.94 days, whereas PhenoFlex_{excluded} reached an RMSE of 3.37 days.

Tenth model run

```
# Set the parameters using the results from the previous run
# Version PhenoFlex_all (pheno_fit_v1_r9$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v1_r10 <- c(58, 305, 0.15, 20.0, 2800.0, 9500.0, 5800.0, 5.70e+13, 0.0, 20.0, 0.0, 0.00)
par_v1_r10 <- c(63, 319, 0.85, 27.9, 3310.4, 9901.6, 6396.2, 5.94e+13, 6.5, 27.9, 5.6, 1.39)
upper_v1_r10 <- c(68, 335, 1.05, 35.0, 3700.0, 10400.0, 6600.0, 6.20e+13, 12.0, 34.0, 10.0, 15.00)

# Same for version PhenoFlex_excluded (pheno_fit_v2_r9$model_fit$par)
# Parameter <- c(yc, zc, s1, Tu, E0, E1, A0, A1, Tf, Tc, Tb, slope)
lower_v2_r10 <- c(28, 360, 0.050, 20.0, 2900.0, 9500.0, 5800.0, 5.70e+13, 0.0, 45.0, 0.0, 0.00)
par_v2_r10 <- c(34, 371, 0.130, 25.0, 3371.0, 9901.3, 6214.6, 5.94e+13, 1.7, 53.3, 4.0, 3.17)
upper_v2_r10 <- c(38, 380, 0.325, 30.0, 3700.0, 10400.0, 6600.0, 6.20e+13, 10.0, 58.0, 10.0, 15.00)
```

In this round of the fitting procedure, PhenoFlex_{all} reached an RMSE of 5.94 days, whereas PhenoFlex_{excluded} reached an RMSE of 3.37 days. Note that the RMSE values obtained in this run did not improve compared to the results of run number 9. For this reason, we used the parameters estimated in the ninth run as final values.

Supplementary validation of PhenoFlex_{excluded}

As mentioned in the main text, we assessed the performance of the PhenoFlex_{excluded} version when used to predict bloom dates in the marginal seasons. Following this approach, we aimed to test the viability of extrapolating the modelling framework to conditions outside the calibration data set. The results of this analysis demonstrated the importance of explicitly testing the predictive accuracy of the model under environmental conditions outside the range of the calibration data. When comparable temperature conditions are used for calibration and validation, the model is – despite potentially incorrect assumptions – sufficiently flexible to be well fitted to data and shows good projection quality. However, when validation data contain conditions outside the calibration range, the conceivably inaccurate model structure cannot be compensated by parameter fitting and finds expression in high residual errors (Fig. S1). In this validation, RMSE and RPIQ reached values of 61.2 days and 0.3, respectively. It should be noted, however, that these results are greatly influenced by one outlier showing 127 days of error (observed versus predicted). When removing the outlier from the marginal seasons, the prediction error ranged between 9.3 and 44.4 days (Fig. S1).




Figure S4: Validation of the PhenoFlex extsubscriptexcluded version using the five marginal seasons excluded from the calibration data set. Whereas the open blue circles represent the experimental seasons used for calibration, the filled red circles represent the five marginal seasons used for validating the framework. The vertical lines in the marginal season dots represent the uncertainty estimated by bootstrapping.

References

- Fernandez, E, P Krefting, A Kunz, H Do, E Fadon, and E Luedeling. 2021. "Boosting Statistical Delineation of Chill and Heat Periods in Temperate Fruit Trees Through multi-environment Observations." *Agricultural and Forest Meteorology* 310, 108652. <https://doi.org/10.1016/j.agrformet.2021.108652>.
- Luedeling, Eike. 2021. *chillR: Statistical Methods for Phenology Analysis in Temperate Fruit Trees*.
- Luedeling, Eike, Katja Schiffers, Till Fohrmann, and Carsten Urbach. 2021. "PhenoFlex - an Integrated Model to Predict Spring Phenology in Temperate Fruit Trees." *Agricultural and Forest Meteorology* 307: 108491. <https://doi.org/10.1016/j.agrformet.2021.108491>.