

Intersection between cylinder and ray

Eduardo Neville (314667), Elias Boschung (315707), Natalja Sagel (362089)

March 2023

1 Finding the Implicit function of a cylinder

We have that for a ray and a cylinder to coincide, the following equality must hold:

$$o + t\vec{d} = c + r \begin{pmatrix} \cos \theta \\ \sin \theta \\ z \end{pmatrix}$$

We will this have to solve for both t and θ . This becomes too complicated and we will have to find a better solution for it. We can use the implicit representation of a cylinder.

Let the axis of the cylinder be a line going through its center $\vec{x}_0 = (x_0, y_0, z_0)$ and in the direction of $\vec{v} = (a, b, c)$ with radius R . Then we can write the equation of the cylinder as the set of all points $\vec{x} = (x, y, z)$ satisfying:

$$|\vec{x} - \vec{x}_0|^2 = R^2 + [(\vec{x} - \vec{x}_0) \cdot \vec{v}]^2$$

where $z \in [z_0 - h, z_0 + h]$.

We have that our $\vec{x} = o + t\vec{d}$ such that

$$0 = R^2 + \left[(o + t\vec{d} - \vec{x}_0) \cdot \vec{v} \right]^2 - |o + t\vec{d} - \vec{x}_0|^2$$

Given that $|\vec{d}| = 1$, solving for t gives us the quadratic equation:

$$0 = ((\vec{d} \cdot \vec{v})^2 - 1)t^2 + 2 \left[((o - \vec{x}_0) \cdot \vec{v})(\vec{d} \cdot \vec{v}) - (o - \vec{x}_0) \cdot \vec{d} \right] t + ((o - \vec{x}_0) \cdot \vec{v})^2 - |o - x_0|^2 + R^2$$