

1. MILESTONE I

In this section we consider the evolution of the uniform background of the universe in the Lambda-Cold-Dark-Matter (Λ CDM) model which is considered to be the present day standard model of cosmology. The main goal of this section is to solve the unperturbed background evolution of the universe by using known cosmological parameters supplied by [?] and comparing this to observational supernova data from [?].

1. Theory

1. Friedmann-Lemaître-Robertson-Walker

In this milestone we approximate the universe to be completely isotropic and homogeneous over all scales. For large scales this has been shown to be a reasonable assumption from observations of the CMB power spectra [?]. The universe must then satisfy a maximally symmetric metric. With some symmetry arguments, we arrive at an exact solution for the metric called the Friedmann-Lemaître-Robertson-Walker (FLRW) metric. In flat space, $k = 0$, and Cartesian coordinates it is given by:

$$g_{\mu\nu} = \text{diag} [-1, a^2(t), a^2(t), a^2(t)],$$

where $a(t)$ is the scale factor, defined to be 1 today, and we are using the mostly plus metric signature $(-, +, +, +)$.

To make use of the line element above we consider the Einstein field equations (EFE):

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G T_{\mu\nu}, \quad (1)$$

With the isotropy requirement, any stress-energy tensors which we will consider must be invariant under spatial rotations. To consider the general allowed form of the stress-energy tensor we decompose it as such:

$$T_{\mu\nu}(x) = \begin{bmatrix} T_{00}(x) & T_{0i}(x) \\ T_{0i}(x) & \sigma_{ij}(x) \end{bmatrix},$$

As usual, the $T_{00} = \rho$ component is to be interpreted as the energy density in a non-boosted frame. The requirement of homogeneity then imposes that one should not have any energy flux, as such finds $T_{0i} = 0$. The isotropy requirement imposes that the matrix σ must satisfy $R^T \sigma(x) R = \sigma(x)$ for all x and rotation matrices R . The latter then implies that $[R, \sigma(x)] = 0$. As the generators of the $SO(3)$ group form an algebra, Schurs' Lemma then states that this is satisfied iff $\sigma(x) \propto I \implies \sigma(x) = p(x)I$ for some function $p(x)$. Furthermore the homogeneity requirement imposes translation invariance, meaning that $p(x) = p(x + x_0)$. These requirements, a comoving observer, i.e. an observer in

the rest frame of the universe, then implies that we may uniquely describe the universe as a perfect liquid with:

$$T_{\mu\nu} = (\rho + p)u_\mu u_\nu + g_{\mu\nu}p,$$

where p is the pressure. As such, from the FRWL metric and the EFE one can derive the **Friedmann equations**:

$$\left(\frac{\dot{a}}{a}\right)^2 \equiv H^2 = \frac{8\pi G}{3}\rho, \quad (2)$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}(\rho + 3p), \quad (3)$$

where ρ and p are, as explained, the energy density and pressure respectively for a perfect liquid.

2. Λ CDM Model

The Λ CDM model is derived by assuming; that we have some family of weakly interacting massive particles (WIMPs) which we call cold-dark-matter (CDM) and a cosmological constant, Λ , which accounts for the present day acceleration of the universe, which will be the dark energy in this article. The name CDM comes from assuming that they are massive enough to not be relativistic and hence cold. The first Friedmann equation (2) can be rewritten such that the time dependence of the Hubble factor $H \equiv \dot{a}/a$ is given by:

$$H^2 = H_0^2 \sum_i \Omega_{i0} a^{-3(1+\omega_i)}, \quad (4)$$

where $\omega_i \equiv p_i/\rho_i$ is the equation of state for the i -th particle type, $\Omega_i \equiv \rho_i/\rho_C$ is the relative energy density of the i -th particle type, ρ_i is the energy density corresponding to the i -th particle type and $\sum_i \rho_i = \rho_C$ is the critical energy density required to have a flat universe which we observe today [?]. We have that $\omega_B = 0$, $\omega_R = 1/3$ and $\omega_\Lambda = -1$ for baryons, relativistic particles (photons and massless neutrinos) and dark energy (DE) respectively. Simply plugging in the various particle types in (4) we have:

$$H = H_0 \sqrt{\Omega_{M0} a^{-3} + \Omega_{R0} a^{-4} + \Omega_{K0} a^{-2} + \Omega_{\Lambda0}}, \quad (5)$$

$$\Omega_{M0} \equiv \Omega_{B0} + \Omega_{CDM0}, \quad \Omega_{R0} \equiv \Omega_{\gamma0} + \Omega_{\nu0},$$

where Ω_{B0} , Ω_{CDM0} , $\Omega_{\gamma0}$, $\Omega_{\nu0}$ and $\Omega_{\Lambda0}$ are then the present day relative densities of baryonic matter (electrons & protons), cold dark matter, radiation, neutrinos and dark energy respectively. The term $\Omega_{K0} = -kc^2/H_0$ denotes the curvature of the universe and encapsulates how said curvature affects expansion rates and energy densities. With the requirement for a flat universe we have that $\sum_i \Omega_i = 1$ where i refers to all non-curvature energy types. Since Ω_B , Ω_{CDM} and Ω_γ , Ω_ν give the same contribution to the time evolution of the Hubble parameter, we will often bundle these together as “matter” and “radiation” (or “relativistic”) particles respectively as was done in (5).

The other Friedmann equation (3), together with each particles equation of state, is then used to describe how each component evolves with time:

$$\begin{aligned}\Omega_B(a) &= \frac{\Omega_{B0}}{a\mathcal{H}^2/H_0^2}, & \Omega_{\text{CDM}}(a) &= \frac{\Omega_{\text{CDM}0}}{a\mathcal{H}^2/H_0^2}, \\ \Omega_\gamma(a) &= \frac{\Omega_{\gamma0}}{a^2\mathcal{H}^2/H_0^2}, & \Omega_\nu(a) &= \frac{\Omega_{\nu0}}{a^2\mathcal{H}^2/H_0^2}, \\ \Omega_K(a) &= \frac{\Omega_{K0}}{\mathcal{H}^2/H_0^2}, & \Omega_\Lambda(a) &= \frac{\Omega_{\Lambda0}}{H^2/H_0^2},\end{aligned}\quad (6)$$

where $\mathcal{H} \equiv aH$ is the **conformal Hubble factor**. Two of the six density parameters follow from the observed temperature of the CMB and are given by:

$$\Omega_{\gamma0} = \frac{8\pi^3 G}{45H_0^2} \frac{(k_B T_{\text{CMB}0})^4}{\hbar^3 c^5}, \quad (7)$$

$$\Omega_{\nu0} = \Omega_{\gamma0} N_{\text{eff}} \cdot \frac{7}{8} \left(\frac{4}{11} \right)^{4/3}, \quad (8)$$

where $T_{\text{CMB}0}$ is the temperature of CMB photons today and $N_{\text{eff}} = 3.046$ is the effective number of massless neutrinos. [?]

The Λ CDM model is then fully determined by these cosmological observables which come from 2018 Planck [?]:

$$\begin{aligned}\frac{H_0 \text{ Mpc}}{100 \text{ km/s}} &\equiv h = 0.67, & N_{\text{eff}} &= 3.046, \\ \Omega_{B0} &= 0.05, & \Omega_{\text{CDM}0} &= 0.267, \\ \Omega_{K0} &= 0.00, & T_{\text{CMB}0} &= 2.7255.\end{aligned}\quad (9)$$

3. Time and Distance Measurements in Cosmology

Since the expansion of the universe is strictly positive w.r.t. time, the scale factor $a(t)$ is an injective function and can thus be used as a time measurement. In this paper we will, for computational purposes, be using the variable $x \equiv \ln(a) \implies a = e^x$.

Again, due to the expansion of the universe, one can also measure how much the wavelength of a photon released at initial time t_i has stretched before it reaches us at final time t_f , commonly known as the **redshift**:

$$z = e^{x(t_f) - x(t_i)} - 1.$$

A derivation can be found in [?]. Since we are mainly interested in the redshift w.r.t. today then $x(t_f) = 0$. Thus the formula which will be used in this article is $z = e^{-x(t_i)} - 1$.

Further we consider the **horizon**, i.e. the “distance” massless particles may have travelled since the big bang. Since the universe is expanding, this will be appreciably larger than ct . Considering a time t_1 , light would have travelled a distance $d_1 > ct_1$. We can then consider what t_1 must be such that this becomes an equality. For this

we define the **conformal time** $d\eta = dt/a$ (or conformal distance with the appropriate insertion of c) which can be rewritten as $\eta(t) \equiv \int_0^t \frac{c}{a} dt'$ and we get the differential equation:

$$\frac{d\eta}{dx} = \frac{c}{\mathcal{H}}, \quad (10)$$

along with the initial condition $\eta(-\infty) = 0$, which can be solved analytically in the radiation dominated era. We will then use the analytical solution to approximate that at some very early time, x_{start} such that we have the new initial condition $\eta(x_{\text{start}}) = c/\mathcal{H}(x_{\text{start}})$ where we will solve onwards numerically.

A useful distance measure is the **comoving distance** χ which is derived by the FLRW metric:

$$\chi = \int_{t_i}^{t_0} dt \frac{c}{a} = \int_1^a da' \frac{c}{a'^2 H} = \int_0^z dz' \frac{c}{H} = \eta - \eta_0.$$

Considering a light-like path with the FLRW metric in curved space one arrives at the **proper distance** r :

$$r = \begin{cases} \chi \frac{\sin(\sqrt{|\Omega_{K0}|} H_0 \chi / c)}{\sqrt{|\Omega_{K0}|} H_0 \chi / c}, & \Omega_{K0} < 0 \text{ “Closed”} \\ \chi, & \Omega_{K0} = 0 \text{ “Flat”} \\ \chi \frac{\sinh(\sqrt{|\Omega_{K0}|} H_0 \chi / c)}{\sqrt{|\Omega_{K0}|} H_0 \chi / c}, & \Omega_{K0} > 0 \text{ “Open”} \end{cases}.$$

We then consider the angular **diameter distance** $d_A = \Delta s / \Delta \theta$ where Δs is the objects physical size and $\Delta \theta$ its angular size as viewed from Earth. Once again from the FRLW metric, now in spherical coordinates:

$$ds^2 = -c^2 dt^2 + a^2(dr^2 + r^2 d\theta^2 + \sin^2 \theta d\phi^2),$$

one can find that this can be expressed in terms of our cosmological parameters as $d_A = ar$. The **luminosity distance** d_L is defined from $d_L = \sqrt{\frac{L}{4\pi F}}$ where L is the intrinsic luminosity and F is the measured flux. As implied by its name, this quantity is related to d_A by $d_L = d_A/a^2 = r/a$.

At last we consider the **cosmic time** t in our x coordinate:

$$t(x) = \int_0^a \frac{da}{aH} = \int_{-\infty}^0 \frac{dx}{H(x)}.$$

From this we get to the ODE:

$$\frac{dt}{dx} = \frac{1}{H}, \quad (11)$$

with the initial condition $t(-\infty) = 0$. This initial condition states that cosmic time begins once the big bang happened. As before, we make an analytical approximation which gives us the initial condition $t(x_{\text{start}}) = \frac{1}{2H(x_{\text{start}})}$ and solved numerically from here.

4. Markov-Chain Monte Carlo Method

In this milestone one of the major computational method which will be used is the Markov Chain Monte Carlo (MCMC) method. The particular algorithm we use is the Metropolis algorithm which generates a Markov chain, where in each step, the chain corresponds to a proposed set of model parameters, which in our case are h, Ω_{M0} and Ω_{K0} . These parameters are sampled from a probability distribution guided by certain priors, i.e. accepted ranges, where data outside of these priors are omitted. The algorithm then eventually converges to the posterior distribution of the parameters given the observed data. This can then be used to constrain the cosmological parameters, in which we decide to focus on h, Ω_{CDM0} and Ω_{M0} . A χ^2 -test is then done by the assumption that the measurements are Gaussian distributed and uncorrelated between different redshift. The likelihood function is then given by $L \propto e^{-\chi^2/2}$ where:

$$\chi^2(h, \Omega_{M0}, \Omega_{K0}) = \sum_{i=1}^N [d_L(z_i, h, \Omega_{M0}, \Omega_{K0}) - d_L^{\text{obs}}]^2 / \sigma_i^2.$$

Here σ_i is the standard deviation for the i -th data point and N is the total number of data points. The set of parameters with the highest likelihood is our best-fit model, which also happen to be the values which minimize χ^2 . To check that this best-fit is in fact a good fit one can consider a set of data points which all lie exactly 1σ away from the observed values. As such the expression in the sum simply reduces to 1. Thus when summing over N terms equal to 1 we would have that $\chi^2/N = 1$. Hence we will consider a best-fit to be a good fit if $\chi^2 \sim N$. For the case when $\chi^2 \ll N$ then we over-fit the model, meaning we may be capturing random noise and fluctuations in the data, whilst if $\chi^2 \gg N$ then we simply have a bad correspondence with the data and/or underlying assumptions (such as the assumption that the measurements are Gaussian).

2. Implementation details

1. Solving the ODEs

To implement some numerical tools to solve for the background we first implement the data from (9) and computed the derived quantities $\Omega_{\gamma 0}$ and $\Omega_{\nu 0}$ with (7,8), and $\Omega_{\Lambda 0} = 1 - \Omega_{\text{rest}0}$ where “rest” refers to all the other Ω_i .

Next we implemented an ordinary differential equation (ODE) solver to solve the ODEs for $\eta(x)$ and $t(x)$. This was done by first setting up the respective differential equation (10,11) with its corresponding initial condition, then an ODE solver was used to get a result which was then splined. The ODE solver and splining programs were already provided in the template.

Further we computed $H(x)$ from (5) which was used together with the various relations from the theory section to compute $\mathcal{H}, \frac{d\mathcal{H}}{dx}, \frac{d^2\mathcal{H}}{dx^2}, \Omega_i, r, d_A, d_L, \chi$ and T_{CMB} all as a function of our time variable x . Note that the expressions for $\frac{d\mathcal{H}}{dx}$ and $\frac{d^2\mathcal{H}}{dx^2}$ were found analytically from 5. The calculated data was then written to data files to be analyzed in Python.

2. MCMC fit

We then compared our numerical data for the luminosity distance d_L to observational data from [?], containing $N = 31$ data points, by performing an MCMC fit. The upper and lower priors which were used are $\{1.5, 1, 1\}$ and $\{0.5, 0, -1\}$ for h, Ω_M and Ω_K respectively. The algorithm then computes an initial χ^2 with some randomly generated parameters. It repeats this process and checks whether the new χ_{new}^2 is less than the old χ_{old}^2 . If this is the case then it accepts this sample and starts again. However if this previous test is not the case we then perform a random check $e^{-(\chi_{\text{new}}^2 - \chi_{\text{old}}^2)/2} > u$ where u is number drawn from a uniform distribution in the interval $[0, 1]$. If it passes this test then χ_{new}^2 is accepted and the process carries on. This additional test is the main part of the Metropolis algorithm and is done to capture the full posterior probability distribution function (PDF) of the various parameters. As before, the data was printed to a data file and imported to Python.

3. Python

x -values corresponding to radiation-matter- and matter-DE-equality were found by finding running a script over the data to find where the absolute value of the difference $|\Omega_i - \Omega_j| = 0$. In reality they were never quite 0 due to running over a discrete index, thus we simply set some low threshold, e.g. $5 \cdot 10^{-5}$. To find when the universe accelerates we note that:

$$\mathcal{H} = \frac{\dot{a}}{a} a = \dot{a} \implies \frac{d\mathcal{H}}{dx} = \frac{d\mathcal{H}}{dt} \frac{dt}{dx} = \ddot{a} \frac{dt}{dx},$$

and from (11) we then have:

$$\frac{d\mathcal{H}}{dx} \equiv \mathcal{H}' = \frac{\ddot{a}}{H}.$$

Since H is strictly positive we can then look at when \mathcal{H}' changes from negative to positive. So again a Python script which runs over the indices of \mathcal{H} were used to find which x -values this sign change happened, which corresponds to where the universe began to accelerate.

The MCMC-fit was then analyzed and a scatterplot in the $\Omega_{\Lambda}\Omega_M$ -plane was created with the $(1\sigma, 2\sigma)$ confidence regions $\chi^2 - \chi_{\text{min}}^2 < (3.53, 8.02)$ respectively. These particular values were found from common χ^2 distribution tables with $k = 3$ degrees of freedom. A histogram of the

PDF from the accepted samples of h along with a Gaussian fit was along with a comparison of our theoretical d_L vs. the real supernova data were also plotted.

3. Tests of data

Before considering the results it is reasonable to conduct some sanity checks on the data to make sure everything was implemented correctly. To check that the data is consistent with the analytical expressions we consider the following quantities analytically in the different regimes:

$$\frac{\mathcal{H}'}{\mathcal{H}}, \quad \frac{1}{\mathcal{H}} \frac{d^2 \mathcal{H}}{dx^2} \equiv \frac{\mathcal{H}''}{\mathcal{H}}, \quad \frac{1}{c} \eta \mathcal{H}.$$

1. \mathcal{H}' and \mathcal{H}''

In the radiation dominated era where we approximate $\Omega_R \approx 1 \implies \Omega_{\text{rest}} \approx 0$. Then by using (5) we find:

$$H_R(x) \approx H_0, \quad \mathcal{H}_R(x) \approx H_0 e^{-x}, \\ \frac{\mathcal{H}'_R}{\mathcal{H}_R} \approx -1, \quad \frac{\mathcal{H}''_R}{\mathcal{H}_R} \approx 1,$$

and similarly using the approximations $\Omega_M \approx 1$ and $\Omega_\Lambda \approx 1$ respectively, we get:

$$H_M(x) \approx H_0 e^{x/2}, \quad \mathcal{H}_M(x) \approx H_0 e^{-x/2}, \\ \frac{\mathcal{H}'_M}{\mathcal{H}_M} \approx -1/2, \quad \frac{\mathcal{H}''_M}{\mathcal{H}_M} \approx 1/4, \\ H_\Lambda(x) \approx H_0, \quad \mathcal{H}_\Lambda(x) \approx H_0 e^x, \\ \frac{\mathcal{H}'_\Lambda}{\mathcal{H}_\Lambda} \approx \frac{\mathcal{H}''_\Lambda}{\mathcal{H}_\Lambda} \approx 1.$$

Plotting these assumptions with the data we have FIG. 1. We see that there is a reasonable agreement with the analytical approximations in the given regimes. As we will see in later, $\Omega_M \approx 1$ is a relatively poor approximation compared to the others, hence larger a deviation is to be expected.

2. Conformal Time

Analytical approximations for the conformal time requires a little more effort. For the radiation dominated era we can solve (10) analytically:

$$\eta_R(x) = \int_{-\infty}^x \frac{c}{\mathcal{H}_R} dx' \approx \int_{-\infty}^x \frac{c}{H_0} e^{x'} dx' = \frac{c}{H_0} e^x.$$

Now we approximate this radiation dominated epoch to end abruptly at some time x_1 such that we can write the




FIG. 1. \mathcal{H}'/\mathcal{H} and $\mathcal{H}''/\mathcal{H}$ compared to analytical approximations in the various regimes where the left and right dash-dotted vertical lines signify radiation-matter equality and matter-dark energy equality respectively.

conformal time in the matter dominated epoch as:

$$\eta_M(x) \approx \eta_R(x_1) + \int_{x_1}^x \frac{c}{\mathcal{H}_M} dx' \\ \approx \frac{c}{H_0} \left(e^{x_1} + \int_{x_1}^x e^{x'/2} dx' \right) \\ = \frac{c}{H_0} (e^{x_1} - 2e^{x_1/2} + 2e^{x/2}).$$

Again we assume that the matter dominated epoch ends abruptly at some time x_2 such that we can approximate:

$$\eta_\Lambda(x) \approx \eta_M(x_2) + \int_{x_2}^\infty \frac{c}{\mathcal{H}_\Lambda} dx' \\ \approx \frac{c}{H_0} \left(e^{x_1} - 2e^{x_1/2} + 2e^{x_2/2} + \int_{x_2}^x e^{-x'} dx' \right) \\ = \frac{c}{H_0} (e^{x_1} - 2e^{x_1/2} + 2e^{x_2/2} + e^{-x_2} - e^{-x}).$$


Thus we find that for the various regimes we have:

$$\eta_R \mathcal{H}_R / c \approx 1, \\ \eta_M \mathcal{H}_M / c \approx (e^{x_1} - 2e^{x_2/2} + 2e^{x/2}) e^{-x/2}, \\ \eta_\Lambda \mathcal{H}_\Lambda / c \approx (e^{x_1} - 2e^{x_1/2} + 2e^{x_2/2} + e^{-x_2} - e^{-x}) e^x.$$

Using the x values corresponding to $\Omega_R = \Omega_M$ and $\Omega_M = \Omega_\Lambda$ as the time where the respective epochs start and end we get the upper graph in FIG. 2. Clearly this approximation will be nothing close to exact due to the assumption that the epochs end abruptly when in reality the epoch's change continuously and relatively slowly. Since η depends on how long the previous epochs lasted we can see that this heavily affects the final epoch.

Overall the data seems to be consistent with the analytical expressions, even if the conformal time is a rather difficult parameter to define a robust approximation for.

FIG. 2. Numerical $\eta\mathcal{H}/c$ compared to analytical approximations in the various regimes where epochs are approximated to abruptly once $\Omega_i = \Omega_j$.



4. Results

Now that the data has been stress tested to show that they are at least valid in certain regions we can go on to look at some of the results.

1. Time Evolution of Cosmic Parameters

We first consider the time evolution of the various relative energy densities Ω_i as a function of the scale factor $a(t)$ as given in FIG. 3.




FIG. 3. Time evolution of the density parameters Ω_i as a function of the scale factor a .

Here we can clearly see the different regimes of the various quantities. The early universe is in a complete radiation domination at early times which implies that the universe expands $a(t) \propto t^{1/2}$. The universe then relatively slowly switches over to becoming matter dominated as time passes. For a short period of time we have complete matter domination, implying an expansion $a(t) \propto t^{2/3}$. Then dark energy starts to dominate at later times, causing the universe to begin to acceler-

ate, until we get the known result of today where dark energy accounts for roughly 68% of the total energy in the universe today. It then further predicts that in the future, dark energy domination will continue to grow as time passes.

The time evolution of the conformal Hubble factor $\mathcal{H}(x)$, cosmic time $t(x)$ and conformal distance $\eta(x)/c$ is given in FIG. 4. The notable acceleration at later times in the \mathcal{H} plot can be seen by when the derivative of \mathcal{H} changes sign as mentioned previously. One can also see that this same phenomena affects both the cosmic time and conformal distance inversely at the same time. This is due to them both being inversely proportional to the conformal Hubble factor.




FIG. 4. Time evolution of the conformal Hubble factor \mathcal{H} , cosmic time t and conformal distance η/c as a function of x .

Next we consider the values of $x, z, t, \eta, \Omega_M, \Omega_\Lambda$ and Ω_R

at various important events. A summary of these events is given in TABLE I.

Here we can see that radiation-matter-equality happened roughly 51000 years after the big bang. Further, the universe begins to accelerate roughly 7.7 billion years after the big bang. One may notice that this happens $2\Omega_\Lambda \approx \Omega_M$ and $\Omega_{\text{others}} \approx 0$. We can check this analytically by considering these approximation. We have that $\Omega_{\text{tot}} \approx 3\Omega_\Lambda$ and hence $\rho_{\text{tot}} = 3\rho_\Lambda$. Further we have that $p_{\text{tot}} = p_\Lambda + p_M = p_\Lambda$ due to $p_M = 0$ from its equation of state. Inserting this into the Friedman equation (3) with the relevant equation of state we get:

$$\begin{aligned} \frac{\ddot{a}}{a} &\approx -\frac{4\pi G}{3}(3\rho_\Lambda + 3(p_\Lambda + p_M)) \\ &= -\frac{4\pi G}{3}(3\rho_\Lambda - 3\rho_\Lambda) = 0 \end{aligned}$$

Now, since a is strictly a positive number then, mathematically, all that remains to check whether this is not when deceleration starts or a terrace point. To ensure this one can just check some arbitrarily close points such as $\Omega_\Lambda = 1/3 \pm \epsilon$ and $\Omega_M = 2/3 \mp \epsilon$ where ϵ is some tiny positive number.

Next in the table we have $t(x=0)$ which corresponds to the age of the universe today. [?] estimates this to be 13.78 ± 0.20 Gly which is reasonably close to our value of 13.86 Gly. Similarly $\eta(x=0)$ shows us the actual size of the observable universe today. [?] states the expected conformal distance of the universe is roughly 14 Gpc which corresponds to 45.7 Gly, again in reasonable agreement with the data.

2. Supernova Comparison

Further we compare our numerical data to data collected from supernova observations [?] by; comparing the luminosity distance is given in FIG. 5 and making a posterior probability distribution function (PDF) of the Hubble parameter H_0 in FIG. 6. We note that the found $\chi^2_{\text{min}} = 29.3$ for the MCMC fit, which then yields that for the points accepted within the 1σ region follow $\chi^2 \sim N$ to a very good approximation, suggesting that we have a good fit.

The luminosity distance from the observational data is seemingly always lower than our data. This phenomena is known as the **Hubble tension** which is a widely known problem in today's standard model of cosmology [?]. Direct data, such as supernova data, seemingly always prefers lower values for luminosity distance, and thus, higher values for h that can be seen in FIG. 6. Indirect data, such as the CMB, however prefer higher values for d_L and thus lower values h with much smaller error-bars. This is an ongoing issue which suggests that the Λ CDM model is not a complete model and must be modified. The posterior PDF of the Hubble parameter today H_0 shows that the Gaussian fit is centered at roughly

70.1 km/s/Mpc. As mentioned, this is a large discrepancy from $H_0 = 67$ km/s/Mpc which we got from the fiducial cosmology but still considered a common result from direct data.




FIG. 5. Luminosity distance over redshift z , plotted against redshift z for both; our numerical data, and observational data from Betoule et. al. [?] with error-bars.




FIG. 6. Posterior PDF of the Hubble parameter H_0 compared to the fiducial value $H_0 = 67$ km/s/Mpc.


Next we consider the scatterplot in the $\Omega_\Lambda\Omega_M$ -plane shown in FIG. 7. The scatter-plot helps us visualize the degeneracy between the density parameters, i.e. that many different combinations of Ω_M and Ω_Λ can yield similar observational results. This leads to rather large degenerate regions in the parameter space. In this figure we can see that the flat universe constraint significantly reduces this degeneracy by lowering the allowed parameters by multiple orders of magnitude. The figure clearly shows that $\Omega_{\Lambda 0} = 0$ is completely excluded given the observational data as this would be far outside the 2σ constraint. The best-fit value for $\Omega_{K0} = 0.067$ tells us that the universe is seemingly quite flat, but that there is indeed some curvature. To test out whether this might just be an insensitivity to the Ω_{K0} parameter the code was run again with a large initial value Ω_{K0} . The result did not seem to change whatsoever and thus this implies

TABLE I. Results for cosmological parameters at various important events from numerical data.

Event	x	$z(x)$	$t(x)$ [Gly]	$\eta(x)/c$ [Gly]	Ω_M	Ω_Λ	Ω_{Rel}
Radiation-Matter Equality	-8.132	3401	$5.106 \cdot 10^{-5}$	0.368	0.500	$2.737 \cdot 10^{-11}$	0.500
Universe Accelerates	-0.486	0.626	7.761	38.55	0.666	0.334	$3.183 \cdot 10^{-4}$
Matter-DE Equality	-0.256	0.292	10.38	42.33	0.500	0.500	$1.900 \cdot 10^{-4}$
Universe Today	0.000	0.000	13.86	46.29	0.317	0.683	$9.320 \cdot 10^{-5}$

that our model is insensitive to the curvature of the universe.

Degeneracy between dark energy and matter density parameters in cosmology

FIG. 7. Supernova data with 1σ and 2σ constraints from MCMC fits.

5. Summary & Conclusion

To summarize this section, we numerically solved the ODEs for η and t past the radiation domination. We then

computed $H(x)$ which gave us access to all the other relevant cosmological parameters. Further we checked that the numerical data corresponds to analytical approximations in the different regimes. Next we introduced all the most relevant data in a graphed form in FIG 4-6 such that important events and values were easily visualized. The most relevant values for the most important events in our background cosmology were summarized in TABLE I. Finally, an MCMC fit with observational data from supernova events and the numerical data in the $\Omega_\Lambda\Omega_M$ -plane was made to consider some important aspects of our model, in particular that we cannot consider a model without dark energy.

In conclusion, the data all seems to correspond quite well with observational data and previous calculations from more sophisticated methods.