

DRP F26 - Coxeter Group

Note Taker: **Edward Chang**

Email: *edward.chang@uwaterloo.ca*

Updated: 2026-09-18

Contents

1	Introduction	2
---	--------------	---


2026-09-14

1 Introduction

Def: Group

A **group** $(G, \bullet)$ where G is a set and $\bullet : G \times G \rightarrow G$ is a binary operator on the set, and it follows that:

- i) (Associativity): $\forall a, b, c \in G, (a \bullet b) \bullet c = a \bullet (b \bullet c)$
- ii) (Identity element): $\exists 1 \in G : \forall a \in G a \bullet 1 = 1 \bullet a = a$
- iii) (Inverse element): $\forall a \in G \exists a^{-1} \in G : a \bullet a^{-1} = a^{-1} \bullet a = 1$



Conjecture:

If mirrors a and b are at a $\theta = \frac{2\pi}{n}$ angle, the group generated by R_a, R_b is finite.

Idea:

- If $\theta \neq \frac{2\pi}{n}$ for any $n \in \mathbb{N}$, we don't get a finite partition of the planes.
- There is some relation between transformations generated by reflections and chambers.

Conjecture:

$$|G| = \text{number of chambers.}$$

Note

Root vec

$$R_a = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$